



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 04:43 AM GMT

PDB ID : 2NXZ  
Title : HIV-1 gp120 Envelope Glycoprotein (T257S, S334A, S375W) Complexed with CD4 and Antibody 17b  
Authors : Zhou, T.; Xu, L.; Dey, B.; Hessel, A.J.; Van Ryk, D.; Xiang, S.H.; Yang, X.; Zhang, M.Y.; Zwick, M.B.; Arthos, J.; Burton, D.R.; Dimitrov, D.S.; Sodroski, J.; Wyatt, R.; Nabel, G.J.; Kwong, P.D.  
Deposited on : 2006-11-20  
Resolution : 2.04 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1192 (2.04-2.04)
Clashscore	102246	1269 (2.04-2.04)
Ramachandran outliers	100387	1258 (2.04-2.04)
Sidechain outliers	100360	1258 (2.04-2.04)
RSRZ outliers	91569	1194 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	317	
2	B	184	
3	C	214	
4	D	229	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	A	730	-	-	-	X
5	NAG	A	734	-	-	-	X
5	NAG	A	789	-	-	-	X
5	NAG	A	839	-	-	-	X
6	SUC	D	4000	-	-	-	X
7	EDO	C	4002	-	-	-	X
8	HEZ	A	4005	-	-	-	X

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 7913 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENVELOPE GLYCOPROTEIN GP120.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	306	Total	C	N	O	S	0	1	0
			2375	1492	413	450	20			

- Molecule 2 is a protein called T-cell surface glycoprotein CD4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	181	Total	C	N	O	S	0	1	0
			1418	889	249	276	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1000	MET	-	INITIATING METHIONINE	UNP P01730

- Molecule 3 is a protein called ANTIBODY 17B, LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	214	Total	C	N	O	S	0	0	0
			1646	1028	282	331	5			

- Molecule 4 is a protein called ANTIBODY 17B, HEAVY CHAIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	222	Total	C	N	O	S	0	1	0
			1682	1065	283	329	5			

- Molecule 5 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is SUGAR (SUCROSE) (three-letter code: SUC) (formula: C<sub>12</sub>H<sub>22</sub>O<sub>11</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	D	1	Total	C	O	0	0
			23	12	11		

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	C	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is HEXANE-1,6-DIOL (three-letter code: HEZ) (formula:  $C_6H_{14}O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 9 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula:  $C_3H_8O$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			4	3	1		

- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	190	Total 190	O 190	0	0
10	B	87	Total 87	O 87	0	0
10	C	124	Total 124	O 124	0	0
10	D	162	Total 162	O 162	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ENVELOPE GLYCOPROTEIN GP120




- Molecule 2: T-cell surface glycoprotein CD4



- Molecule 3: ANTIBODY 17B, LIGHT CHAIN



- Molecule 4: ANTIBODY 17B, HEAVY CHAIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.92Å 88.13Å 197.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.38 – 2.04 35.38 – 2.04	Depositor EDS
% Data completeness (in resolution range)	87.1 (35.38-2.04) 87.2 (35.38-2.04)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.29 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.206 , 0.244 0.202 , 0.239	Depositor DCC
$R_{free}$ test set	3578 reflections (5.36%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.1	Xtriage
Anisotropy	0.110	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 45.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 70311 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7913	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: IPA, SUC, NAG, HEZ, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.34	0/2427	0.48	0/3292
2	B	0.29	0/1443	0.51	1/1945 (0.1%)
3	C	0.61	3/1683 (0.2%)	0.53	1/2288 (0.0%)
4	D	0.33	0/1727	0.51	0/2353
All	All	0.41	3/7280 (0.0%)	0.51	2/9878 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	2192	LYS	CE-NZ	18.08	1.94	1.49
3	C	2192	LYS	CG-CD	9.19	1.83	1.52
3	C	2192	LYS	CD-CE	6.69	1.68	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2192	LYS	CB-CG-CD	-5.49	97.33	111.60
2	B	1037	LEU	CA-CB-CG	5.45	127.84	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2375	0	2317	27	0
2	B	1418	0	1444	16	0
3	C	1646	0	1590	23	0
4	D	1682	0	1647	18	0
5	A	182	0	169	1	0
6	D	23	0	22	4	0
7	A	8	0	12	0	0
7	C	4	0	6	1	0
8	A	8	0	14	3	0
9	B	4	0	6	1	0
10	A	190	0	0	2	1
10	B	87	0	0	3	0
10	C	124	0	0	0	0
10	D	162	0	0	3	0
All	All	7913	0	7227	86	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 86 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:2192:LYS:CG	3:C:2192:LYS:CD	1.83	1.52
3:C:2192:LYS:CE	3:C:2192:LYS:NZ	1.94	1.29
4:D:3215:HIS:HB3	4:D:3220:THR:HG22	1.34	1.07
4:D:3046:GLU:OE2	4:D:3063[B]:HIS:CE1	2.23	0.91
4:D:3046:GLU:OE2	4:D:3063[B]:HIS:HE1	1.55	0.89

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:A:4133:HOH:O	10:A:4133:HOH:O[4_556]	1.55	0.65

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/317 (96%)	284 (94%)	16 (5%)	3 (1%)	19	8
2	B	180/184 (98%)	173 (96%)	7 (4%)	0	100	100
3	C	212/214 (99%)	203 (96%)	9 (4%)	0	100	100
4	D	219/229 (96%)	214 (98%)	5 (2%)	0	100	100
All	All	914/944 (97%)	874 (96%)	37 (4%)	3 (0%)	46	36

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	464	GLU
1	A	462	ASN
1	A	398	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/280 (97%)	265 (98%)	6 (2%)	60	53
2	B	165/166 (99%)	155 (94%)	10 (6%)	23	13
3	C	184/184 (100%)	176 (96%)	8 (4%)	35	27
4	D	188/193 (97%)	180 (96%)	8 (4%)	35	27
All	All	808/823 (98%)	776 (96%)	32 (4%)	38	29

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	1109	LEU
3	C	2090	GLN
4	D	3165	VAL
3	C	2073	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
3	C	2107	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	1033	GLN
2	B	1040	GLN
4	D	3043	GLN
1	A	462	ASN
3	C	2102	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
7	EDO	A	4001	-	3,3,3	0.43	0	2,2,2	0.42	0
7	EDO	A	4003	-	3,3,3	0.47	0	2,2,2	0.43	0
8	HEZ	A	4005	-	7,7,7	0.26	0	6,6,6	0.59	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	A	588	1	14,14,15	0.50	0	15,19,21	0.59	0
5	NAG	A	697	1	14,14,15	0.44	0	15,19,21	0.65	0
5	NAG	A	730	1	14,14,15	0.51	0	15,19,21	0.63	0
5	NAG	A	734	1	14,14,15	0.55	0	15,19,21	0.65	0
5	NAG	A	741	1	14,14,15	0.57	0	15,19,21	0.74	0
5	NAG	A	762	1	14,14,15	0.58	0	15,19,21	0.87	0
5	NAG	A	776	1	14,14,15	0.55	0	15,19,21	0.76	0
5	NAG	A	789	1	14,14,15	0.53	0	15,19,21	0.71	0
5	NAG	A	795	1	14,14,15	0.55	0	15,19,21	1.06	0
5	NAG	A	839	1	14,14,15	0.52	0	15,19,21	0.68	0
5	NAG	A	886	1	14,14,15	0.52	0	15,19,21	0.77	0
5	NAG	A	892	1	14,14,15	0.55	0	15,19,21	0.57	0
5	NAG	A	948	1	14,14,15	0.51	0	15,19,21	0.78	0
9	IPA	B	4004	-	3,3,3	1.42	1 (33%)	3,3,3	1.16	0
7	EDO	C	4002	-	3,3,3	0.48	0	2,2,2	0.41	0
6	SUC	D	4000	-	24,24,24	0.41	0	36,36,36	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	A	4001	-	-	0/1/1/1	0/0/0/0
7	EDO	A	4003	-	-	0/1/1/1	0/0/0/0
8	HEZ	A	4005	-	-	0/5/5/5	0/0/0/0
5	NAG	A	588	1	-	0/6/23/26	0/1/1/1
5	NAG	A	697	1	-	0/6/23/26	0/1/1/1
5	NAG	A	730	1	-	0/6/23/26	0/1/1/1
5	NAG	A	734	1	-	0/6/23/26	0/1/1/1
5	NAG	A	741	1	-	0/6/23/26	0/1/1/1
5	NAG	A	762	1	-	0/6/23/26	0/1/1/1
5	NAG	A	776	1	-	0/6/23/26	0/1/1/1
5	NAG	A	789	1	-	0/6/23/26	0/1/1/1
5	NAG	A	795	1	-	0/6/23/26	0/1/1/1
5	NAG	A	839	1	-	0/6/23/26	0/1/1/1
5	NAG	A	886	1	-	0/6/23/26	0/1/1/1
5	NAG	A	892	1	-	0/6/23/26	0/1/1/1
5	NAG	A	948	1	-	0/6/23/26	0/1/1/1
9	IPA	B	4004	-	-	0/0/0/0	0/0/0/0
7	EDO	C	4002	-	-	0/1/1/1	0/0/0/0
6	SUC	D	4000	-	-	0/12/51/51	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	B	4004	IPA	C3-C2	-2.02	1.33	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	4005	HEZ	3	0
5	A	795	NAG	1	0
9	B	4004	IPA	1	0
7	C	4002	EDO	1	0
6	D	4000	SUC	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	306/317 (96%)	0.57	32 (10%) 8 9	7, 24, 56, 87	0
2	B	181/184 (98%)	0.37	14 (7%) 16 19	11, 29, 51, 61	0
3	C	214/214 (100%)	0.01	1 (0%) 91 93	6, 22, 50, 60	0
4	D	222/229 (96%)	0.05	7 (3%) 51 58	7, 18, 47, 69	0
All	All	923/944 (97%)	0.28	54 (5%) 26 29	6, 23, 53, 87	0

The worst 5 of 54 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	399	THR	7.8
1	A	129	ALA	6.6
1	A	462	ASN	6.0
2	B	1001	LYS	5.4
1	A	89	VAL	5.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
8	HEZ	A	4005	8/8	0.71	0.35	9.63	41,43,44,45	0
5	NAG	A	730	14/15	0.77	0.51	8.29	58,63,65,65	0
5	NAG	A	734	14/15	0.75	0.27	8.09	41,45,47,47	0
5	NAG	A	839	14/15	0.54	0.44	3.33	42,47,48,48	0
7	EDO	C	4002	4/4	0.86	0.17	2.82	32,33,33,33	0
6	SUC	D	4000	23/23	0.81	0.26	2.57	39,41,41,42	0
5	NAG	A	789	14/15	0.91	0.21	2.43	29,31,33,33	0
5	NAG	A	948	14/15	0.74	0.23	1.12	37,39,41,42	0
5	NAG	A	795	14/15	0.91	0.17	0.34	23,26,28,28	0
5	NAG	A	776	14/15	0.89	0.15	0.15	31,34,36,37	0
7	EDO	A	4001	4/4	0.98	0.12	-0.06	16,17,18,18	0
5	NAG	A	762	14/15	0.92	0.14	-0.49	21,25,27,28	0
5	NAG	A	741	14/15	0.78	0.43	-	43,47,48,48	0
5	NAG	A	886	14/15	0.87	0.13	-	30,33,34,35	0
7	EDO	A	4003	4/4	0.93	0.19	-	31,31,32,32	0
5	NAG	A	697	14/15	0.74	0.46	-	44,48,49,49	0
9	IPA	B	4004	4/4	0.90	0.20	-	31,31,31,31	0
5	NAG	A	588	14/15	0.47	0.51	-	36,49,52,52	0
5	NAG	A	892	14/15	0.68	0.44	-	38,42,43,44	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.