



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 04:44 AM GMT

PDB ID : 2NZ9  
Title : Crystal structure of botulinum neurotoxin type A complexed with monoclonal antibody AR2  
Authors : Stevens, R.C.; Arndt, J.W.  
Deposited on : 2006-11-22  
Resolution : 3.79 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1317 (4.10-3.50)
Clashscore	102246	1458 (4.10-3.50)
Ramachandran outliers	100387	1397 (4.10-3.50)
Sidechain outliers	100360	1392 (4.10-3.50)
RSRZ outliers	91569	1325 (4.10-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1295	 61% 31% 5% ..
1	B	1295	 63% 30% 5% ..
2	C	218	 60% 33% 6% .
2	E	218	 62% 30% 6% ..
3	D	224	 63% 27% 5% ..

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	224	 A horizontal bar chart showing the quality of chain F. The bar is divided into three segments: green (63%), yellow (29%), and red (8%). The green segment is labeled '63%' and the yellow segment is labeled '29%'. The red segment is the smallest and is followed by three dots '...'.

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 26985 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Botulinum neurotoxin type A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1267	Total	C	N	O	S	0	0	0
			10200	6541	1691	1937	31			
1	B	1267	Total	C	N	O	S	0	0	0
			10203	6543	1691	1938	31			

- Molecule 2 is a protein called AR2 monoclonal antibody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	216	Total	C	N	O	S	0	0	0
			1657	1039	279	334	5			
2	E	216	Total	C	N	O	S	0	0	0
			1657	1039	279	334	5			

- Molecule 3 is a protein called AR2 monoclonal antibody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	217	Total	C	N	O	S	0	0	0
			1632	1025	271	329	7			
3	F	217	Total	C	N	O	S	0	0	0
			1632	1025	271	329	7			

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		
4	A	1	Total	Zn	0	0
			1	1		


- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total 1	Ca 1	0	0
5	A	1	Total 1	Ca 1	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Botulinum neurotoxin type A







• Molecule 1: Botulinum neurotoxin type A




- Molecule 2: AR2 monoclonal antibody




- Molecule 2: AR2 monoclonal antibody



- Molecule 3: AR2 monoclonal antibody



- Molecule 3: AR2 monoclonal antibody





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.97Å 197.60Å 146.10Å 90.00° 91.18° 90.00°	Depositor
Resolution (Å)	40.00 – 3.79 48.97 – 3.79	Depositor EDS
% Data completeness (in resolution range)	95.5 (40.00-3.79) 95.5 (48.97-3.79)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.90 (at 3.77Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.224 , 0.278 0.203 , 0.242	Depositor DCC
$R_{free}$ test set	2686 reflections (5.38%)	DCC
Wilson B-factor (Å <sup>2</sup> )	95.2	Xtriage
Anisotropy	0.122	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 112.2	EDS
Estimated twinning fraction	0.085 for h,-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 52593 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	26985	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	84.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.64	0/10416	0.67	0/14098
1	B	0.64	0/10420	0.66	0/14104
2	C	0.64	0/1695	0.60	0/2303
2	E	0.65	1/1695 (0.1%)	0.62	0/2303
3	D	0.56	0/1673	0.66	1/2281 (0.0%)
3	F	0.52	0/1673	0.64	0/2281
All	All	0.63	1/27572 (0.0%)	0.65	1/37370 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	26
1	B	0	27
2	C	0	3
2	E	0	3
3	D	0	6
3	F	0	5
All	All	0	70

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	127	GLU	CD-OE1	5.77	1.31	1.25

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	153	GLU	C-N-CD	-8.33	102.27	120.60

There are no chirality outliers.

5 of 70 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	118	TRP	Peptide
1	A	119	GLY	Peptide
1	A	484	ASP	Peptide
1	A	507	ASN	Peptide
1	A	547	LYS	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10200	0	10003	357	0
1	B	10203	0	9999	372	0
2	C	1657	0	1595	64	0
2	E	1657	0	1595	70	0
3	D	1632	0	1552	64	0
3	F	1632	0	1552	73	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
All	All	26985	0	26296	992	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 992 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:269:HIS:CD2	1:B:365:THR:HG21	1.40	1.56
1:A:596:LYS:CB	1:A:597:ALA:HB3	1.39	1.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:132:SER:HA	3:F:133:SER:CB	1.42	1.46
1:B:540:GLY:CA	1:B:541:LYS:HB2	1.40	1.43
1:B:578:ASN:HB3	1:B:579:PRO:CA	1.41	1.43

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1259/1295 (97%)	1084 (86%)	130 (10%)	45 (4%)	4	41
1	B	1259/1295 (97%)	1081 (86%)	139 (11%)	39 (3%)	5	45
2	C	214/218 (98%)	182 (85%)	30 (14%)	2 (1%)	21	68
2	E	214/218 (98%)	193 (90%)	14 (6%)	7 (3%)	5	43
3	D	215/224 (96%)	182 (85%)	24 (11%)	9 (4%)	3	36
3	F	215/224 (96%)	192 (89%)	17 (8%)	6 (3%)	6	47
All	All	3376/3474 (97%)	2914 (86%)	354 (10%)	108 (3%)	5	44

5 of 108 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	121	SER
1	A	400	ASN
1	A	560	GLU
1	A	564	SER
1	A	566	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1124/1178 (95%)	1020 (91%)	104 (9%)	11	47
1	B	1124/1178 (95%)	1037 (92%)	87 (8%)	16	56
2	C	186/190 (98%)	165 (89%)	21 (11%)	7	38
2	E	186/190 (98%)	165 (89%)	21 (11%)	7	38
3	D	181/190 (95%)	162 (90%)	19 (10%)	8	41
3	F	181/190 (95%)	165 (91%)	16 (9%)	12	50
All	All	2982/3116 (96%)	2714 (91%)	268 (9%)	12	49

5 of 268 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	254	SER
1	B	764	PHE
2	E	169	GLU
1	B	297	LEU
1	B	568	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 48 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	53	ASN
1	B	246	ASN
3	D	82	GLN
1	B	67	GLN
1	B	205	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1267/1295 (97%)	-0.14	2 (0%) 95 91	29, 84, 98, 110	0
1	B	1267/1295 (97%)	-0.12	9 (0%) 89 80	33, 85, 97, 111	0
2	C	216/218 (99%)	-0.12	0 100 100	62, 84, 92, 95	0
2	E	216/218 (99%)	-0.23	0 100 100	67, 85, 92, 95	0
3	D	217/224 (96%)	-0.13	0 100 100	75, 85, 95, 103	0
3	F	217/224 (96%)	-0.15	0 100 100	72, 84, 92, 99	0
All	All	3400/3474 (97%)	-0.14	11 (0%) 94 89	29, 84, 97, 111	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	366	TYR	3.2
1	B	1198	SER	2.9
1	B	1237	MET	2.4
1	B	1160	PHE	2.3
1	B	1256	ASN	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	CA	B	1298	1/1	0.91	0.14	-	110,110,110,110	0
4	ZN	B	1297	1/1	0.99	0.13	-	72,72,72,72	0
4	ZN	A	1	1/1	0.96	0.15	-	81,81,81,81	0
5	CA	A	1297	1/1	0.85	0.23	-	101,101,101,101	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.