



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:08 AM GMT

PDB ID : 3NZX  
Title : Crystal structure of the yeast 20S proteasome in complex with ligand 2c  
Authors : Groll, M.; Gallastegui, N.; Marechal, X.; Le Ravalec, V.; Basse, N.; Richey, N.;  
Genin, E.; Huber, R.; Moroder, M.; Vidal, V.; Reboud-Ravaux, M.  
Deposited on : 2010-07-17  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	<div> <div>2%</div> <div>73% 24% •</div> </div>
1	O	250	<div> <div>2%</div> <div>73% 24% •</div> </div>
2	B	258	<div> <div>2%</div> <div>58% 32% • 5%</div> </div>
2	P	258	<div> <div>2%</div> <div>58% 32% 5% 5%</div> </div>
3	C	254	<div> <div>2%</div> <div>57% 35% • 5%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Q	254	
4	D	260	
4	R	260	
5	E	234	
5	S	234	
6	F	288	
6	T	288	
7	G	252	
7	U	252	
8	H	261	
8	V	261	
9	I	205	
9	W	205	
10	J	198	
10	X	198	
11	K	287	
11	Y	287	
12	L	241	
12	Z	241	
13	1	266	
13	M	266	
14	2	215	
14	N	215	
15	3	5	
15	4	5	

## 2 Entry composition

There are 16 unique types of molecules in this entry. The entry contains 50982 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome component Y7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			
1	O	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			

- Molecule 2 is a protein called Proteasome component Y13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	244	Total	C	N	O	S	0	0	0
			1904	1201	321	379	3			
2	P	244	Total	C	N	O	S	0	0	0
			1904	1201	321	379	3			

- Molecule 3 is a protein called Proteasome component PRE6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	241	Total	C	N	O	S	0	0	0
			1890	1181	331	374	4			
3	Q	241	Total	C	N	O	S	0	0	0
			1890	1181	331	374	4			

- Molecule 4 is a protein called Proteasome component PUP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	242	Total	C	N	O	S	0	0	0
			1861	1162	314	378	7			
4	R	242	Total	C	N	O	S	0	0	0
			1861	1162	314	378	7			

- Molecule 5 is a protein called Proteasome component PRE5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	233	Total	C	N	O	S	0	0	0
			1795	1129	312	350	4			
5	S	233	Total	C	N	O	S	0	0	0
			1795	1129	312	350	4			

- Molecule 6 is a protein called Proteasome component C1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	244	Total	C	N	O	S	0	0	0
			1896	1205	330	357	4			
6	T	244	Total	C	N	O	S	0	0	0
			1896	1205	330	357	4			

- Molecule 7 is a protein called Proteasome component C7-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	243	Total	C	N	O	S	0	0	0
			1921	1221	322	370	8			
7	U	243	Total	C	N	O	S	0	0	0
			1921	1221	322	370	8			

- Molecule 8 is a protein called Proteasome component PUP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	222	Total	C	N	O	S	0	0	0
			1684	1061	293	323	7			
8	V	222	Total	C	N	O	S	0	0	0
			1684	1061	293	323	7			

- Molecule 9 is a protein called Proteasome component PUP3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			
9	W	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			

- Molecule 10 is a protein called Proteasome component C11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	198	Total	C	N	O	S	0	0	0
			1585	1005	269	305	6			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	X	198	Total	C	N	O	S	0	0	0
			1585	1005	269	305	6			

- Molecule 11 is a protein called Proteasome component PRE2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			
11	Y	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			

- Molecule 12 is a protein called Proteasome component C5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			
12	Z	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			

- Molecule 13 is a protein called Proteasome component PRE4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			
13	1	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			

- Molecule 14 is a protein called Proteasome component PRE3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			
14	2	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			

- Molecule 15 is a protein called TMC-95A mimic ligand 2c.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
15	3	5	Total	C	N	O	0	0	0
			55	42	5	8			
15	4	5	Total	C	N	O	0	0	0
			55	42	5	8			

- Molecule 16 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
16	A	57	Total O 57 57	0	0
16	B	36	Total O 36 36	0	0
16	C	44	Total O 44 44	0	0
16	D	41	Total O 41 41	0	0
16	E	23	Total O 23 23	0	0
16	F	45	Total O 45 45	0	0
16	G	64	Total O 64 64	0	0
16	H	51	Total O 51 51	0	0
16	I	69	Total O 69 69	0	0
16	J	51	Total O 51 51	0	0
16	K	40	Total O 40 40	0	0
16	L	56	Total O 56 56	0	0
16	M	73	Total O 73 73	0	0
16	N	54	Total O 54 54	0	0
16	O	34	Total O 34 34	0	0
16	P	29	Total O 29 29	0	0
16	Q	27	Total O 27 27	0	0
16	R	31	Total O 31 31	0	0
16	S	20	Total O 20 20	0	0
16	T	38	Total O 38 38	0	0
16	U	64	Total O 64 64	0	0

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	V	47	Total 47	O 47	0	0
16	W	61	Total 61	O 61	0	0
16	X	48	Total 48	O 48	0	0
16	Y	47	Total 47	O 47	0	0
16	Z	48	Total 48	O 48	0	0
16	1	78	Total 78	O 78	0	0
16	2	58	Total 58	O 58	0	0




### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome component Y7



- Molecule 1: Proteasome component Y7




- Molecule 2: Proteasome component Y13




- Molecule 2: Proteasome component Y13







• Molecule 3: Proteasome component PRE6




• Molecule 3: Proteasome component PRE6




• Molecule 4: Proteasome component PUP2




- Molecule 4: Proteasome component PUP2




- Molecule 5: Proteasome component PRE5




- Molecule 5: Proteasome component PRE5




- Molecule 6: Proteasome component C1









• Molecule 8: Proteasome component PUP1




• Molecule 9: Proteasome component PUP3



• Molecule 9: Proteasome component PUP3




• Molecule 10: Proteasome component C11






• Molecule 10: Proteasome component C11



• Molecule 11: Proteasome component PRE2




• Molecule 11: Proteasome component PRE2




• Molecule 12: Proteasome component C5







- Molecule 12: Proteasome component C5



- Molecule 13: Proteasome component PRE4



- Molecule 13: Proteasome component PRE4



- Molecule 14: Proteasome component PRE3

■ D92 ■ N94 ■ I100 ■ I101 ■ A102 ■ D105 ■ D106 ■ K10B ■ I106 ■ G108 ■ T112 ■ I113 ■ V119 ■ H120 ■ A127 ■ I134 ■ I135 ■ G136 ■ K140 ■ E144 ■ E145 ■ M146 ■ E149 ■ E150 ■ H157 ■ S158 ■ Q161 ■ A162 ■ I163 ■ G170 ■ R174 ■ M175 ■ V176 ■ W177 ■ L178 ■ L187 ■ Q181 ■ I181

E86	E89	I100	I101	A102	K106	K107	G108	T112	I113	P114	L115	G116	V119	H120	A127	I134	I135	G136	K140	E144	I145	M146	E149	E150	K156	H157	S158	G161	A162	I163	G170	R174	M175	V176	<b>E177</b>	L178	L187	Q181	L191
-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-------------	------	------	------	------



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	134.20Å 300.11Å 143.63Å 90.00° 112.60° 90.00°	Depositor
Resolution (Å)	15.00 – 2.70 49.68 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (15.00-2.70) 99.7 (49.68-2.50)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.42 (at 2.51Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.226 , 0.248 0.225 , 0.226	Depositor DCC
$R_{free}$ test set	14176 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	39.7	Xtriage
Anisotropy	0.902	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 52.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 359106 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	50982	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TY5, BOC, ABN, RE3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/1952	0.64	0/2642
1	O	0.38	0/1952	0.64	0/2642
2	B	0.37	0/1934	0.64	0/2618
2	P	0.39	0/1934	0.64	0/2618
3	C	0.36	0/1919	0.62	0/2598
3	Q	0.38	0/1919	0.62	0/2598
4	D	0.37	0/1886	0.63	0/2541
4	R	0.38	0/1886	0.63	0/2541
5	E	0.36	0/1823	0.62	0/2463
5	S	0.38	0/1823	0.62	0/2463
6	F	0.38	0/1936	0.62	0/2614
6	T	0.39	0/1936	0.63	0/2614
7	G	0.41	0/1959	0.64	0/2652
7	U	0.40	0/1959	0.64	0/2652
8	H	0.37	0/1715	0.66	0/2326
8	V	0.36	0/1715	0.66	0/2326
9	I	0.40	0/1611	0.68	0/2174
9	W	0.41	0/1611	0.68	0/2174
10	J	0.40	0/1613	0.66	0/2173
10	X	0.40	0/1613	0.67	0/2173
11	K	0.46	0/1681	0.67	1/2274 (0.0%)
11	Y	0.46	0/1681	0.68	1/2274 (0.0%)
12	L	0.41	0/1795	0.68	0/2420
12	Z	0.41	0/1795	0.68	0/2420
13	1	0.40	0/1855	0.68	1/2514 (0.0%)
13	M	0.39	0/1855	0.67	1/2514 (0.0%)
14	2	0.40	0/1541	0.64	1/2087 (0.0%)
14	N	0.40	0/1541	0.64	0/2087
15	3	0.78	0/4	2.03	0/4
15	4	0.90	0/4	2.10	0/4
All	All	0.39	0/50448	0.65	5/68200 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
12	L	0	1
12	Z	0	1
All	All	0	2

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	Y	4	LEU	CA-CB-CG	5.65	128.30	115.30
11	K	4	LEU	CA-CB-CG	5.59	128.15	115.30
13	1	95	LEU	N-CA-C	-5.45	96.29	111.00
13	M	95	LEU	N-CA-C	-5.24	96.86	111.00
14	2	22	THR	N-CA-C	-5.18	97.01	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
12	L	145	TYR	Sidechain
12	Z	145	TYR	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1915	0	1926	54	0
1	O	1915	0	1926	58	0
2	B	1904	0	1901	92	0
2	P	1904	0	1901	89	0
3	C	1890	0	1900	97	0
3	Q	1890	0	1900	98	0
4	D	1861	0	1836	52	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	R	1861	0	1836	55	0
5	E	1795	0	1797	95	0
5	S	1795	0	1797	108	0
6	F	1896	0	1886	73	0
6	T	1896	0	1886	84	0
7	G	1921	0	1910	72	0
7	U	1921	0	1910	77	0
8	H	1684	0	1688	57	0
8	V	1684	0	1688	60	0
9	I	1581	0	1574	63	0
9	W	1581	0	1574	59	0
10	J	1585	0	1590	89	0
10	X	1585	0	1590	77	0
11	K	1644	0	1595	73	0
11	Y	1644	0	1595	79	0
12	L	1757	0	1711	70	0
12	Z	1757	0	1711	74	0
13	1	1824	0	1832	65	0
13	M	1824	0	1832	59	0
14	2	1512	0	1481	53	0
14	N	1512	0	1481	50	0
15	3	55	0	44	4	0
15	4	55	0	44	4	0
16	1	78	0	0	4	0
16	2	58	0	0	4	0
16	A	57	0	0	2	0
16	B	36	0	0	1	0
16	C	44	0	0	2	0
16	D	41	0	0	2	0
16	E	23	0	0	3	0
16	F	45	0	0	2	0
16	G	64	0	0	0	0
16	H	51	0	0	5	0
16	I	69	0	0	3	0
16	J	51	0	0	7	0
16	K	40	0	0	4	0
16	L	56	0	0	7	0
16	M	73	0	0	4	0
16	N	54	0	0	4	0
16	O	34	0	0	1	0
16	P	29	0	0	3	0
16	Q	27	0	0	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
16	R	31	0	0	2	0
16	S	20	0	0	1	0
16	T	38	0	0	3	0
16	U	64	0	0	8	0
16	V	47	0	0	4	0
16	W	61	0	0	1	0
16	X	48	0	0	3	0
16	Y	47	0	0	8	0
16	Z	48	0	0	4	0
All	All	50982	0	49342	1804	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 1804 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:K:208:ASN:ND2	9:W:29:ASN:HD21	1.22	1.32
9:I:29:ASN:HD21	11:Y:208:ASN:ND2	1.23	1.30
11:K:10(B):LYS:H	11:K:10(B):LYS:HD2	1.05	1.18
2:P:202:THR:HG22	2:P:204:SER:H	1.10	1.14
11:Y:10(B):LYS:H	11:Y:10(B):LYS:HD2	1.05	1.13

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/250 (99%)	235 (95%)	10 (4%)	3 (1%)	16	39
1	O	248/250 (99%)	234 (94%)	11 (4%)	3 (1%)	16	39

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	242/258 (94%)	220 (91%)	18 (7%)	4 (2%)	11	29
2	P	242/258 (94%)	221 (91%)	17 (7%)	4 (2%)	11	29
3	C	239/254 (94%)	222 (93%)	13 (5%)	4 (2%)	11	29
3	Q	239/254 (94%)	221 (92%)	14 (6%)	4 (2%)	11	29
4	D	240/260 (92%)	217 (90%)	20 (8%)	3 (1%)	15	37
4	R	240/260 (92%)	216 (90%)	20 (8%)	4 (2%)	11	29
5	E	231/234 (99%)	207 (90%)	20 (9%)	4 (2%)	11	29
5	S	231/234 (99%)	208 (90%)	19 (8%)	4 (2%)	11	29
6	F	242/288 (84%)	228 (94%)	14 (6%)	0	100	100
6	T	242/288 (84%)	228 (94%)	13 (5%)	1 (0%)	39	69
7	G	241/252 (96%)	229 (95%)	11 (5%)	1 (0%)	39	69
7	U	241/252 (96%)	228 (95%)	12 (5%)	1 (0%)	39	69
8	H	220/261 (84%)	207 (94%)	9 (4%)	4 (2%)	11	27
8	V	220/261 (84%)	208 (94%)	8 (4%)	4 (2%)	11	27
9	I	202/205 (98%)	192 (95%)	8 (4%)	2 (1%)	19	45
9	W	202/205 (98%)	193 (96%)	7 (4%)	2 (1%)	19	45
10	J	196/198 (99%)	185 (94%)	8 (4%)	3 (2%)	13	32
10	X	196/198 (99%)	185 (94%)	8 (4%)	3 (2%)	13	32
11	K	210/287 (73%)	198 (94%)	9 (4%)	3 (1%)	14	35
11	Y	210/287 (73%)	199 (95%)	8 (4%)	3 (1%)	14	35
12	L	220/241 (91%)	202 (92%)	17 (8%)	1 (0%)	34	63
12	Z	220/241 (91%)	200 (91%)	19 (9%)	1 (0%)	34	63
13	1	231/266 (87%)	216 (94%)	14 (6%)	1 (0%)	39	69
13	M	231/266 (87%)	217 (94%)	13 (6%)	1 (0%)	39	69
14	2	194/215 (90%)	184 (95%)	10 (5%)	0	100	100
14	N	194/215 (90%)	185 (95%)	9 (5%)	0	100	100
15	3	1/5 (20%)	1 (100%)	0	0	100	100
15	4	1/5 (20%)	1 (100%)	0	0	100	100
All	All	6314/6948 (91%)	5887 (93%)	359 (6%)	68 (1%)	17	42

5 of 68 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	58	LEU
3	C	203	THR
4	D	12(G)	GLU
5	E	5	ARG
5	E	202	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	198 (95%)	11 (5%)	28	57
1	O	209/209 (100%)	200 (96%)	9 (4%)	35	66
2	B	203/216 (94%)	191 (94%)	12 (6%)	24	51
2	P	203/216 (94%)	189 (93%)	14 (7%)	19	43
3	C	213/226 (94%)	202 (95%)	11 (5%)	29	58
3	Q	213/226 (94%)	202 (95%)	11 (5%)	29	58
4	D	198/215 (92%)	185 (93%)	13 (7%)	21	45
4	R	198/215 (92%)	185 (93%)	13 (7%)	21	45
5	E	192/193 (100%)	173 (90%)	19 (10%)	10	22
5	S	192/193 (100%)	174 (91%)	18 (9%)	11	25
6	F	201/239 (84%)	179 (89%)	22 (11%)	8	18
6	T	201/239 (84%)	179 (89%)	22 (11%)	8	18
7	G	207/210 (99%)	194 (94%)	13 (6%)	22	48
7	U	207/210 (99%)	195 (94%)	12 (6%)	25	52
8	H	181/214 (85%)	175 (97%)	6 (3%)	45	76
8	V	181/214 (85%)	175 (97%)	6 (3%)	45	76
9	I	172/173 (99%)	165 (96%)	7 (4%)	37	69
9	W	172/173 (99%)	167 (97%)	5 (3%)	50	80
10	J	175/175 (100%)	163 (93%)	12 (7%)	19	43
10	X	175/175 (100%)	164 (94%)	11 (6%)	22	48

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
11	K	169/235 (72%)	157 (93%)	12 (7%)	18	41
11	Y	169/235 (72%)	156 (92%)	13 (8%)	16	36
12	L	185/201 (92%)	169 (91%)	16 (9%)	13	29
12	Z	185/201 (92%)	169 (91%)	16 (9%)	13	29
13	1	199/224 (89%)	187 (94%)	12 (6%)	24	50
13	M	199/224 (89%)	189 (95%)	10 (5%)	30	60
14	2	162/178 (91%)	152 (94%)	10 (6%)	23	49
14	N	162/178 (91%)	153 (94%)	9 (6%)	26	54
All	All	5332/5816 (92%)	4987 (94%)	345 (6%)	21	46

5 of 345 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
13	M	62	LEU
2	P	212	PHE
12	Z	99	THR
13	M	129	PHE
1	O	10(A)	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 193 such sidechains are listed below:

Mol	Chain	Res	Type
13	M	89	GLN
3	Q	121	GLN
12	Z	85	HIS
13	M	149	GLN
2	P	95	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
15	TY5	3	2	15	19,20,21	1.80	7 (36%)	22,25,27	0.96	2 (9%)
15	RE3	3	4	15	15,17,18	1.77	6 (40%)	21,25,27	2.18	6 (28%)
15	TY5	4	2	15	19,20,21	1.78	6 (31%)	22,25,27	0.97	1 (4%)
15	RE3	4	4	15	15,17,18	1.83	5 (33%)	21,25,27	2.18	7 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
15	TY5	3	2	15	-	0/9/11/13	0/2/2/2
15	RE3	3	4	15	-	0/5/23/25	0/2/2/2
15	TY5	4	2	15	-	0/9/11/13	0/2/2/2
15	RE3	4	4	15	-	0/5/23/25	0/2/2/2

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
15	3	4	RE3	CB-CA	-2.63	1.50	1.54
15	3	2	TY5	C53-C52	2.03	1.43	1.38
15	4	2	TY5	C53-C52	2.04	1.43	1.38
15	3	4	RE3	CZ3-CE3	2.04	1.43	1.38
15	3	2	TY5	C54-C55	2.05	1.43	1.38

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	3	4	RE3	CE2-NE1-CD1	-5.09	109.14	111.88
15	4	4	RE3	CE2-NE1-CD1	-5.02	109.17	111.88
15	3	4	RE3	CG-CD2-CE2	-4.14	106.97	108.80
15	4	4	RE3	CG-CD2-CE2	-3.98	107.04	108.80
15	3	4	RE3	O-C-CA	-2.64	118.62	125.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
15	3	2	TY5	2	0
15	4	2	TY5	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	250/250 (100%)	-0.46	3 (1%) 81 81	40, 54, 84, 107	0
1	O	250/250 (100%)	-0.41	4 (1%) 74 75	40, 56, 85, 107	0
2	B	244/258 (94%)	-0.29	4 (1%) 74 75	41, 58, 95, 120	0
2	P	244/258 (94%)	-0.24	6 (2%) 61 61	41, 59, 96, 120	0
3	C	241/254 (94%)	-0.11	5 (2%) 67 68	42, 63, 112, 129	0
3	Q	241/254 (94%)	0.18	21 (8%) 13 10	45, 65, 112, 129	0
4	D	242/260 (93%)	-0.22	8 (3%) 50 50	43, 61, 96, 127	0
4	R	242/260 (93%)	-0.17	8 (3%) 50 50	44, 62, 96, 127	0
5	E	233/234 (99%)	-0.20	4 (1%) 73 74	45, 65, 91, 113	0
5	S	233/234 (99%)	0.13	12 (5%) 31 30	45, 67, 93, 113	0
6	F	244/288 (84%)	-0.42	2 (0%) 87 88	42, 58, 91, 108	0
6	T	244/288 (84%)	-0.15	6 (2%) 61 61	43, 58, 92, 107	0
7	G	243/252 (96%)	-0.44	3 (1%) 81 81	38, 53, 79, 118	0
7	U	243/252 (96%)	-0.36	2 (0%) 87 88	39, 54, 79, 118	0
8	H	222/261 (85%)	-0.43	1 (0%) 91 93	37, 52, 70, 103	0
8	V	222/261 (85%)	-0.55	1 (0%) 91 93	38, 52, 70, 104	0
9	I	204/205 (99%)	-0.49	1 (0%) 91 93	37, 50, 68, 87	0
9	W	204/205 (99%)	-0.39	1 (0%) 91 93	37, 50, 68, 86	0
10	J	198/198 (100%)	-0.45	4 (2%) 68 69	37, 51, 69, 127	0
10	X	198/198 (100%)	-0.47	5 (2%) 61 61	39, 51, 69, 127	0
11	K	212/287 (73%)	-0.39	4 (1%) 70 70	36, 49, 76, 85	0
11	Y	212/287 (73%)	-0.43	3 (1%) 78 77	35, 51, 77, 86	0
12	L	222/241 (92%)	-0.47	3 (1%) 78 77	35, 51, 72, 100	0
12	Z	222/241 (92%)	-0.50	4 (1%) 71 72	35, 51, 72, 100	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
13	1	233/266 (87%)	-0.59	0 100 100	37, 51, 64, 70	0
13	M	233/266 (87%)	-0.56	1 (0%) 93 94	37, 51, 64, 70	0
14	2	196/215 (91%)	-0.53	0 100 100	35, 48, 67, 81	0
14	N	196/215 (91%)	-0.58	0 100 100	37, 47, 67, 82	0
15	3	1/5 (20%)	-0.24	0 100 100	46, 46, 46, 46	0
15	4	1/5 (20%)	0.12	0 100 100	48, 48, 48, 48	0
All	All	6370/6948 (91%)	-0.35	116 (1%) 71 72	35, 54, 87, 129	0

The worst 5 of 116 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	R	12(F)	GLY	11.8
10	J	192	ALA	10.0
3	C	55	THR	8.9
7	U	6	ALA	8.7
4	D	12(E)	SER	8.3

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
15	TY5	4	2	19/20	0.94	0.16	-	41,46,55,55	0
15	RE3	4	4	16/17	0.94	0.14	-	50,52,54,57	0
15	RE3	3	4	16/17	0.94	0.16	-	49,51,52,54	0
15	TY5	3	2	19/20	0.96	0.15	-	42,45,54,55	0

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.