



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:45 AM GMT

PDB ID : 2O0R
Title : The three-dimensional structure of N-Succinyldiaminopimelate aminotransferase from Mycobacterium tuberculosis
Authors : Weyand, S.; Kefala, G.; Weiss, M.S.; TB Structural Genomics Consortium (TBSGC)
Deposited on : 2006-11-28
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	411	 4% 86% 7% • 6%
1	B	411	 3% 80% 12% • 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CL	B	602	-	-	-	X
3	NA	B	603	-	-	-	X
4	GOL	B	702	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6319 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rv0858c (N-Succinyldiaminopimelate aminotransferase).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	385	Total	C	N	O	P	S	0	5	0
			2931	1856	517	545	1	12			
1	B	384	Total	C	N	O	P	S	0	5	0
			2916	1848	507	548	1	12			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	INITIATING METHIONINE	UNP O53870
A	1	ALA	-	CLONING ARTIFACT	UNP O53870
A	232	LLP	LYS	MODIFIED RESIDUE	UNP O53870
A	398	GLY	-	CLONING ARTIFACT	UNP O53870
A	399	VAL	-	CLONING ARTIFACT	UNP O53870
A	400	PRO	-	CLONING ARTIFACT	UNP O53870
A	401	ARG	-	CLONING ARTIFACT	UNP O53870
A	402	GLY	-	CLONING ARTIFACT	UNP O53870
A	403	LEU	-	CLONING ARTIFACT	UNP O53870
A	404	GLU	-	CLONING ARTIFACT	UNP O53870
A	405	HIS	-	EXPRESSION TAG	UNP O53870
A	406	HIS	-	EXPRESSION TAG	UNP O53870
A	407	HIS	-	EXPRESSION TAG	UNP O53870
A	408	HIS	-	EXPRESSION TAG	UNP O53870
A	409	HIS	-	EXPRESSION TAG	UNP O53870
A	410	HIS	-	EXPRESSION TAG	UNP O53870
B	0	MET	-	INITIATING METHIONINE	UNP O53870
B	1	ALA	-	CLONING ARTIFACT	UNP O53870
B	232	LLP	LYS	MODIFIED RESIDUE	UNP O53870
B	398	GLY	-	CLONING ARTIFACT	UNP O53870
B	399	VAL	-	CLONING ARTIFACT	UNP O53870
B	400	PRO	-	CLONING ARTIFACT	UNP O53870
B	401	ARG	-	CLONING ARTIFACT	UNP O53870
B	402	GLY	-	CLONING ARTIFACT	UNP O53870
B	403	LEU	-	CLONING ARTIFACT	UNP O53870

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	404	GLU	-	CLONING ARTIFACT	UNP O53870
B	405	HIS	-	EXPRESSION TAG	UNP O53870
B	406	HIS	-	EXPRESSION TAG	UNP O53870
B	407	HIS	-	EXPRESSION TAG	UNP O53870
B	408	HIS	-	EXPRESSION TAG	UNP O53870
B	409	HIS	-	EXPRESSION TAG	UNP O53870
B	410	HIS	-	EXPRESSION TAG	UNP O53870


- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Cl 1 1	0	0
2	A	1	Total Cl 1 1	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Na 1 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	225	Total	O	0	0
			225	225		
5	B	232	Total	O	0	0
			232	232		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Rv0858c (N-Succinyldiaminopimelate aminotransferase)



- Molecule 1: Rv0858c (N-Succinyldiaminopimelate aminotransferase)



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	54.34Å 56.13Å 247.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 17.92 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (20.00-2.00) 99.9 (17.92-2.00)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.37 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.164 , 0.213 0.167 , 0.211	Depositor DCC
R_{free} test set	1080 reflections (2.11%)	DCC
Wilson B-factor (Å ²)	23.1	Xtriage
Anisotropy	0.450	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.3	EDS
Estimated twinning fraction	0.026 for k,h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 52200 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6319	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, LLP, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	0/2992	0.80	2/4084 (0.0%)
1	B	0.85	0/2977	0.84	1/4066 (0.0%)
All	All	0.82	0/5969	0.82	3/8150 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	7	ARG	NE-CZ-NH2	-5.50	117.55	120.30
1	A	133	ARG	NE-CZ-NH2	-5.42	117.59	120.30
1	B	75	ARG	NE-CZ-NH2	-5.15	117.72	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	141	ASP	Peptide
1	B	23	ARG	Peptide
1	B	364	ASP	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2931	0	2898	27	0
1	B	2916	0	2873	38	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	B	1	0	0	0	0
4	A	6	0	8	0	0
4	B	6	0	8	1	0
5	A	225	0	0	5	0
5	B	232	0	0	1	0
All	All	6319	0	5787	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23:ARG:HH11	1:B:23:ARG:HG2	1.34	0.92
1:A:292:ARG:HD3	5:A:906:HOH:O	1.79	0.82
1:B:120:ASP:OD2	4:B:702:GOL:H31	1.82	0.79
1:A:379:ASP:HB2	5:A:842:HOH:O	1.85	0.77
1:B:197[B]:THR:HG21	5:B:733:HOH:O	1.84	0.76
1:A:23:ARG:HE	1:A:23:ARG:HA	1.53	0.74
1:A:1:ALA:HB2	5:A:826:HOH:O	1.94	0.68
1:B:23:ARG:HG2	1:B:23:ARG:NH1	2.03	0.65
1:A:318:LEU:C	1:A:318:LEU:HD12	2.18	0.64
1:A:12:THR:HG21	1:A:120:ASP:OD2	1.97	0.63
1:B:23:ARG:HH11	1:B:23:ARG:CG	2.11	0.60
1:B:364:ASP:O	1:B:365:VAL:HB	2.02	0.60
1:B:36:GLU:HG3	1:B:235:ASN:OD1	2.01	0.59
1:B:12:THR:HG21	1:B:120:ASP:OD2	2.02	0.59
1:A:1:ALA:C	1:A:2:THR:HG1	2.05	0.59
1:A:68:ARG:NH2	5:A:790:HOH:O	2.38	0.56
1:A:367:ASN:ND2	1:A:367:ASN:H	2.04	0.56
1:A:204:LEU:HD21	1:A:291:LEU:HD12	1.85	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:166:ASN:HD21	1:B:170:ASN:H	1.55	0.54
1:A:2:THR:N	5:A:839:HOH:O	2.42	0.52
1:B:323:ARG:HD3	1:B:368:HIS:ND1	2.24	0.51
1:A:237:THR:HB	1:B:57:TYR:CE1	2.45	0.51
1:B:247:PRO:HB2	1:B:249[A]:GLU:OE2	2.11	0.51
1:B:376:LYS:HB2	1:B:381:LEU:HD13	1.92	0.51
1:A:27:VAL:HG11	1:A:384:ALA:HB2	1.92	0.51
1:B:64:ALA:HB3	1:B:65:PRO:HD3	1.92	0.51
1:A:204:LEU:HD21	1:A:291:LEU:CD1	2.40	0.50
1:B:111:GLU:OE1	1:B:159:ARG:NH1	2.45	0.49
1:B:305:ILE:HG12	1:B:389:SER:HA	1.94	0.48
1:A:167:SER:HA	1:A:168:PRO:C	2.32	0.48
1:A:1:ALA:C	1:A:2:THR:OG1	2.52	0.47
1:B:198:ASP:OD1	1:B:232:LLP:N1	2.48	0.47
1:B:12:THR:HB	1:B:14:PHE:HD1	1.80	0.46
1:B:387:ARG:O	1:B:390:VAL:HG22	2.16	0.46
1:A:111:GLU:HG3	1:A:159[B]:ARG:HG3	1.97	0.46
1:B:204:LEU:HD11	1:B:291:LEU:HD13	1.97	0.46
1:A:23:ARG:NE	1:A:23:ARG:HA	2.27	0.45
1:A:1:ALA:O	1:B:253:GLY:CA	2.64	0.45
1:B:23:ARG:HD3	1:B:23:ARG:O	2.16	0.45
1:B:225:ILE:HD11	1:B:250:LEU:HB3	1.99	0.45
1:B:229:SER:HA	1:B:242:GLY:HA2	1.98	0.45
1:B:166:ASN:ND2	1:B:170:ASN:H	2.13	0.45
1:B:85[B]:GLU:HG2	1:B:86:THR:HG23	1.98	0.45
1:B:323:ARG:HH22	1:B:365:VAL:HG22	1.81	0.44
1:B:20:LEU:HD13	1:B:338:PRO:HG2	1.98	0.44
1:A:16:GLU:H	1:A:16:GLU:HG2	1.50	0.44
1:A:318:LEU:O	1:A:318:LEU:HD12	2.17	0.44
1:A:367:ASN:HD22	1:A:367:ASN:H	1.66	0.44
1:B:204:LEU:HD21	1:B:291:LEU:HD12	1.99	0.43
1:A:1:ALA:O	1:B:253:GLY:HA3	2.19	0.43
1:B:167:SER:HA	1:B:168:PRO:C	2.39	0.43
1:A:157:THR:HB	1:A:158:PRO:HD2	2.01	0.42
1:A:249:GLU:CD	1:A:249:GLU:H	2.21	0.42
1:A:12:THR:CG2	1:A:120:ASP:OD2	2.64	0.42
1:B:113:LEU:HD11	1:B:156:VAL:HG22	2.01	0.42
1:B:101:ALA:HA	1:B:254:VAL:HG22	2.02	0.41
1:B:162:ALA:HA	1:B:194:VAL:O	2.20	0.41
1:B:364:ASP:O	1:B:365:VAL:CB	2.67	0.41
1:A:198:ASP:OD1	1:A:232:LLP:N1	2.54	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:66:LEU:HA	1:B:274:ALA:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/411 (94%)	373 (97%)	10 (3%)	2 (0%)	34	26
1	B	384/411 (93%)	372 (97%)	10 (3%)	2 (0%)	34	26
All	All	769/822 (94%)	745 (97%)	20 (3%)	4 (0%)	34	26

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	365	VAL
1	A	263	TYR
1	B	263	TYR
1	A	354	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	295/309 (96%)	288 (98%)	7 (2%)	57	58

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	294/309 (95%)	284 (97%)	10 (3%)	44	41
All	All	589/618 (95%)	572 (97%)	17 (3%)	57	49

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	GLU
1	A	23	ARG
1	A	118	PHE
1	A	288[A]	ARG
1	A	288[B]	ARG
1	A	316	TYR
1	A	367	ASN
1	B	20	LEU
1	B	23	ARG
1	B	222[A]	GLU
1	B	222[B]	GLU
1	B	249[A]	GLU
1	B	249[B]	GLU
1	B	292	ARG
1	B	316	TYR
1	B	341	VAL
1	B	381	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	289	ASN
1	A	367	ASN
1	B	47	GLN
1	B	132	HIS
1	B	166	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	A	232	1	23,24,25	1.73	3 (13%)	28,32,34	1.86	7 (25%)
1	LLP	B	232	1	23,24,25	1.59	4 (17%)	28,32,34	2.29	9 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	232	1	-	0/15/17/19	0/1/1/1
1	LLP	B	232	1	-	0/15/17/19	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	232	LLP	O3-C3	-5.49	1.24	1.37
1	B	232	LLP	O3-C3	-4.34	1.26	1.37
1	A	232	LLP	C3-C2	-2.69	1.38	1.40
1	A	232	LLP	C6-N1	2.14	1.39	1.34
1	B	232	LLP	C6-N1	2.53	1.39	1.34
1	B	232	LLP	C4-C4'	2.65	1.51	1.46
1	B	232	LLP	C2-N1	3.04	1.40	1.34

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	232	LLP	C4-C4'-NZ	-3.83	103.77	125.06
1	B	232	LLP	OP4-P-OP1	-3.68	97.78	107.14
1	B	232	LLP	C4-C4'-NZ	-3.24	107.02	125.06
1	B	232	LLP	CD-CE-NZ	-3.23	105.69	110.98

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	232	LLP	C5-C6-N1	-3.19	118.32	123.86
1	B	232	LLP	C3-C4-C4'	-2.88	116.43	120.16
1	A	232	LLP	OP4-P-OP1	-2.82	99.95	107.14
1	B	232	LLP	C5-C6-N1	-2.78	119.04	123.86
1	A	232	LLP	CD-CE-NZ	-2.39	107.07	110.98
1	A	232	LLP	O-C-CA	-2.13	119.93	125.49
1	B	232	LLP	OP2-P-OP1	2.05	117.17	110.58
1	B	232	LLP	O3-C3-C2	3.03	122.93	117.66
1	A	232	LLP	OP4-C5'-C5	4.00	115.60	108.99
1	A	232	LLP	C3-C4-C5	4.12	121.19	118.11
1	B	232	LLP	C3-C4-C5	4.98	121.84	118.11
1	B	232	LLP	OP4-C5'-C5	6.92	120.43	108.99

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	232	LLP	1	0
1	B	232	LLP	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	701	-	5,5,5	0.41	0	5,5,5	0.66	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	B	702	-	5,5,5	0.35	0	5,5,5	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	701	-	-	0/4/4/4	0/0/0/0
4	GOL	B	702	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	702	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	384/411 (93%)	-0.10	16 (4%) 40 41	7, 14, 31, 60	1 (0%)
1	B	383/411 (93%)	-0.10	13 (3%) 49 50	7, 14, 32, 51	0
All	All	767/822 (93%)	-0.10	29 (3%) 44 45	7, 14, 33, 60	1 (0%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	23	ARG	6.9
1	A	23	ARG	6.0
1	B	355	ALA	5.8
1	B	364	ASP	5.0
1	A	365	VAL	4.3
1	B	22	THR	4.2
1	A	354	PRO	3.9
1	A	355	ALA	3.9
1	B	365	VAL	3.9
1	B	339	GLU	3.8
1	A	1	ALA	3.7
1	A	393	GLU	3.7
1	A	19	ALA	3.7
1	A	22	THR	3.1
1	A	24	ILE	3.0
1	B	354	PRO	3.0
1	A	328	ASP	2.6
1	A	364	ASP	2.6
1	B	332	GLU	2.5
1	B	326	GLY	2.4
1	A	12	THR	2.3
1	A	15	ALA	2.2
1	A	339	GLU	2.2
1	A	17	MET	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	328	ASP	2.2
1	B	19	ALA	2.2
1	B	24	ILE	2.1
1	B	329	ASP	2.1
1	A	20	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	LLP	B	232	24/25	0.97	0.09	-	11,14,19,24	0
1	LLP	A	232	24/25	0.98	0.08	-	9,14,23,28	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	CL	B	602	1/1	0.99	0.17	3.87	12,12,12,12	0
4	GOL	B	702	6/6	0.86	0.18	3.57	33,37,39,46	0
3	NA	B	603	1/1	0.91	0.14	3.03	32,32,32,32	0
4	GOL	A	701	6/6	0.86	0.20	1.60	30,36,41,44	0
2	CL	A	601	1/1	0.98	0.17	-	11,11,11,11	0

6.5 Other polymers

There are no such residues in this entry.