



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:46 AM GMT

PDB ID : 2O54  
Title : Structure of E. coli topoisomerase III in complex with an 8-base single stranded oligonucleotide. Frozen in glycerol at pH 7.0  
Authors : Changela, A.; Digate, R.J.; Mondragon, A.  
Deposited on : 2006-12-05  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	8	<div> <div>13%</div> <div>13% 50% 25% 13%</div> </div>
1	D	8	<div> <div>25% 63% 13%</div> </div>
2	A	659	<div> <div>11%</div> <div>83% 11% . .</div> </div>
2	B	659	<div> <div>4%</div> <div>84% 11% . .</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CL	A	800	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10538 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(\*CP\*GP\*CP\*AP\*AP\*CP\*TP\*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	7	Total	C	N	O	P	0	0	0
			138	67	26	39	6			
1	D	8	Total	C	N	O	P	0	0	0
			158	77	28	46	7			

- Molecule 2 is a protein called DNA topoisomerase 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	630	Total	C	N	O	S	0	0	0
			5013	3165	912	916	20			
2	B	631	Total	C	N	O	S	0	0	0
			5016	3168	913	915	20			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	654	HIS	-	EXPRESSION TAG	UNP P14294
A	655	HIS	-	EXPRESSION TAG	UNP P14294
A	656	HIS	-	EXPRESSION TAG	UNP P14294
A	657	HIS	-	EXPRESSION TAG	UNP P14294
A	658	HIS	-	EXPRESSION TAG	UNP P14294
A	659	HIS	-	EXPRESSION TAG	UNP P14294
B	654	HIS	-	EXPRESSION TAG	UNP P14294
B	655	HIS	-	EXPRESSION TAG	UNP P14294
B	656	HIS	-	EXPRESSION TAG	UNP P14294
B	657	HIS	-	EXPRESSION TAG	UNP P14294
B	658	HIS	-	EXPRESSION TAG	UNP P14294
B	659	HIS	-	EXPRESSION TAG	UNP P14294

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cl	0	0
			2	2		

- Molecule 4 is ACETIC ACID (three-letter code: ACY) (formula: C<sub>2</sub>H<sub>4</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		



- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	79	Total	O	0	0
			79	79		
5	B	117	Total	O	0	0
			117	117		
5	C	1	Total	O	0	0
			1	1		
5	D	10	Total	O	0	0
			10	10		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: 5'-D(\*CP\*GP\*CP\*AP\*AP\*CP\*TP\*T)-3'




- Molecule 1: 5'-D(\*CP\*GP\*CP\*AP\*AP\*CP\*TP\*T)-3'




- Molecule 2: DNA topoisomerase 3



- Molecule 2: DNA topoisomerase 3





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	101.83Å 101.83Å 452.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.10 – 2.50 29.10 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.7 (29.10-2.50) 97.7 (29.10-2.50)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.88 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.222 , 0.269 0.222 , 0.264	Depositor DCC
$R_{free}$ test set	4104 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	45.2	Xtriage
Anisotropy	0.089	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 45.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.41$ , $\langle L^2 \rangle = 0.23$	Xtriage
Outliers	0 of 81830 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	10538	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACY, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	C	1.10	0/154	1.99	5/235 (2.1%)
1	D	1.07	0/176	1.84	7/269 (2.6%)
2	A	0.48	0/5123	0.61	2/6945 (0.0%)
2	B	0.53	0/5127	0.63	1/6952 (0.0%)
All	All	0.54	0/10580	0.70	15/14401 (0.1%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	1	DC	O4'-C1'-N1	9.54	114.68	108.00
1	C	5	DA	O4'-C1'-N9	8.85	114.19	108.00
1	D	5	DA	O4'-C1'-N9	8.57	114.00	108.00
1	D	8	DT	O4'-C4'-C3'	-6.87	101.75	104.50
1	D	2	DG	P-O3'-C3'	6.67	127.70	119.70
1	C	6	DC	P-O3'-C3'	6.64	127.67	119.70
1	C	7	DT	C6-C5-C7	-6.18	119.19	122.90
1	D	3	DC	P-O3'-C3'	6.14	127.07	119.70
1	C	4	DA	O4'-C1'-N9	-6.05	103.76	108.00
1	D	1	DC	P-O3'-C3'	-5.39	113.23	119.70
2	A	178	ARG	NE-CZ-NH1	5.39	123.00	120.30
2	B	455	ARG	NE-CZ-NH1	5.38	122.99	120.30
2	A	178	ARG	NE-CZ-NH2	-5.25	117.68	120.30
1	D	8	DT	C6-C5-C7	-5.17	119.80	122.90
1	D	1	DC	O4'-C1'-N1	5.15	111.60	108.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	138	0	80	2	0
1	D	158	0	92	2	0
2	A	5013	0	5014	49	0
2	B	5016	0	5020	39	0
3	A	2	0	0	0	0
4	B	4	0	3	0	0
5	A	79	0	0	2	0
5	B	117	0	0	0	0
5	C	1	0	0	0	0
5	D	10	0	0	0	0
All	All	10538	0	10209	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:396:THR:OG1	2:A:399:GLU:HG3	1.78	0.83
2:A:588:TYR:HB3	2:A:656:HIS:HE1	1.47	0.80
2:A:197:ARG:HD2	2:A:574:TRP:CZ3	2.17	0.79
2:A:588:TYR:HB3	2:A:656:HIS:CE1	2.23	0.74
2:A:169:ASP:OD2	2:A:197:ARG:HG2	1.89	0.72
2:A:520:ASP:HB3	2:A:525:GLU:OE2	1.89	0.72
2:B:525:GLU:OE2	2:B:525:GLU:HA	1.91	0.70
2:B:219:LYS:NZ	2:B:251:ASP:OD1	2.25	0.69
2:A:646:GLU:HB2	2:A:649:SER:HB3	1.77	0.67
2:A:187:ALA:HB2	2:A:611:VAL:HG13	1.76	0.66
2:B:87:VAL:HG22	2:B:90:ARG:HH21	1.61	0.66
2:B:259:ARG:NH1	2:B:263:GLU:OE2	2.30	0.65
2:B:92:LEU:HG	2:B:121:LEU:HD13	1.77	0.64
2:A:197:ARG:CD	2:A:574:TRP:CZ3	2.81	0.64
2:A:21:LYS:HG3	2:A:22:PRO:HA	1.81	0.62
2:A:126:ARG:HH12	2:A:152:ASN:HD22	1.50	0.59
2:A:515:ILE:HD13	2:A:576:SER:HB2	1.83	0.59

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:98:ILE:HB	2:B:129:VAL:HG22	1.86	0.58
2:A:169:ASP:OD2	2:A:197:ARG:CG	2.51	0.58
1:D:7:DT:H5'	2:B:8:LYS:HE2	1.86	0.58
2:A:503:ILE:HG23	2:A:516:LEU:HD13	1.87	0.57
2:A:577:VAL:O	2:A:581:ILE:HG13	2.04	0.57
2:A:169:ASP:OD2	2:A:197:ARG:HD3	2.04	0.56
2:B:221:PHE:CD1	2:B:255:ARG:HB3	2.41	0.56
2:B:106:ARG:HG3	2:B:162:ALA:HB2	1.87	0.56
2:B:150:ARG:HH11	2:B:150:ARG:HG2	1.70	0.55
2:A:58:TYR:O	2:A:178:ARG:NH2	2.26	0.55
2:B:427:LEU:HD21	2:B:473:LEU:HD12	1.89	0.55
2:B:583:GLU:O	2:B:584:LYS:HB2	2.08	0.54
2:B:72:LYS:NZ	2:B:649:SER:HB2	2.23	0.54
2:B:7:GLU:HB2	2:B:102:GLY:HA2	1.90	0.54
2:A:1:MET:HG2	2:A:35:GLY:O	2.08	0.54
2:A:419:VAL:HB	2:A:443:GLU:HB2	1.89	0.53
2:A:453:LYS:O	2:A:457:GLU:HG2	2.09	0.52
1:C:6:DC:H3'	1:C:7:DT:H5'	1.90	0.52
2:B:277:THR:O	2:B:469:LYS:HD2	2.10	0.52
2:A:520:ASP:CB	2:A:525:GLU:OE2	2.58	0.52
2:A:106:ARG:HB2	2:A:162:ALA:HB2	1.92	0.51
2:B:177:THR:HG23	2:B:194:SER:HA	1.92	0.51
2:B:455:ARG:HD2	2:B:456:ASP:OD2	2.11	0.51
1:C:2:DG:N7	2:A:178:ARG:HD2	2.26	0.51
2:B:100:HIS:CE1	2:B:112:VAL:HB	2.45	0.50
2:A:361:GLN:HG2	2:A:449:LEU:HD21	1.94	0.50
2:B:46:LEU:HD12	2:B:114:GLU:HG3	1.93	0.50
2:B:574:TRP:CZ2	2:B:595:LEU:CD1	2.96	0.49
2:A:197:ARG:HD2	2:A:574:TRP:CE3	2.49	0.48
2:A:83:LYS:O	2:A:87:VAL:HG23	2.13	0.48
2:B:187:ALA:HB2	2:B:611:VAL:HB	1.94	0.48
2:B:88:ILE:O	2:B:92:LEU:HB2	2.13	0.48
2:A:276:VAL:HG21	2:A:467:VAL:HG13	1.96	0.48
2:B:100:HIS:NE2	2:B:112:VAL:HB	2.29	0.47
2:A:615:ARG:O	2:A:617:ILE:HG13	2.15	0.47
2:A:509:ASP:HB3	2:A:512:LEU:HB2	1.95	0.47
2:B:507:VAL:CG2	2:B:568:PRO:HG2	2.45	0.46
2:B:208:ARG:NH2	2:B:566:THR:O	2.48	0.46
2:B:232:ALA:HB3	2:B:234:GLU:OE1	2.15	0.46
1:D:8:DT:O3'	2:B:321:GLU:OE1	2.30	0.46
2:A:240:TRP:O	2:A:242:PRO:HD3	2.16	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:229:VAL:HG12	2:B:474:LEU:HB3	1.98	0.45
2:A:169:ASP:OD2	2:A:197:ARG:CD	2.65	0.45
2:B:525:GLU:CA	2:B:525:GLU:OE2	2.62	0.45
2:A:277:THR:O	2:A:469:LYS:HE2	2.17	0.45
2:A:98:ILE:HB	2:A:129:VAL:HG22	1.99	0.45
2:B:7:GLU:HG3	2:B:111:LEU:HD11	1.98	0.44
2:B:444:ALA:O	2:B:447:ARG:HB2	2.17	0.44
2:B:247:GLU:N	2:B:248:PRO:HD2	2.32	0.44
2:B:574:TRP:CZ2	2:B:595:LEU:HD11	2.53	0.44
2:A:197:ARG:HH21	2:A:571:THR:HG23	1.83	0.44
2:A:133:LEU:HD11	2:A:155:PHE:CZ	2.52	0.44
2:A:268:ARG:HD3	5:A:861:HOH:O	2.17	0.44
2:B:89:LYS:HD3	2:B:119:LEU:HD22	1.98	0.44
2:A:450:LEU:HB3	2:A:454:GLU:HG3	2.00	0.43
2:A:588:TYR:CB	2:A:656:HIS:HE1	2.24	0.43
2:A:584:LYS:NZ	5:A:811:HOH:O	2.52	0.43
2:A:1:MET:HE1	2:A:3:LEU:HB2	2.00	0.42
2:B:327:THR:O	2:B:328:TYR:C	2.57	0.42
2:A:151:SER:O	2:A:154:GLU:HG2	2.20	0.42
2:B:579:THR:O	2:B:583:GLU:HG2	2.20	0.42
2:A:133:LEU:HD11	2:A:155:PHE:HZ	1.85	0.41
2:A:202:VAL:O	2:A:203:LEU:C	2.58	0.41
2:A:117:ASP:HB3	2:A:648:GLY:HA2	2.00	0.41
2:A:69:VAL:HB	2:B:593:GLN:OE1	2.20	0.41
2:A:265:VAL:O	2:A:269:ILE:HG12	2.21	0.41
2:A:88:ILE:O	2:A:92:LEU:HB2	2.20	0.41
2:B:6:ALA:HA	2:B:101:ALA:O	2.20	0.41
2:B:422:LYS:HA	2:B:438:ALA:O	2.21	0.41
2:A:276:VAL:HG21	2:A:467:VAL:CG1	2.52	0.40
2:A:428:ASP:HA	2:A:432:GLY:O	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	626/659 (95%)	591 (94%)	28 (4%)	7 (1%)	17	31
2	B	627/659 (95%)	608 (97%)	19 (3%)	0	100	100
All	All	1253/1318 (95%)	1199 (96%)	47 (4%)	7 (1%)	30	50

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	455	ARG
2	A	615	ARG
2	A	616	GLY
2	A	463	PRO
2	A	525	GLU
2	A	453	LYS
2	A	647	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	535/555 (96%)	520 (97%)	15 (3%)	51	78
2	B	535/555 (96%)	518 (97%)	17 (3%)	46	74
All	All	1070/1110 (96%)	1038 (97%)	32 (3%)	48	76

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	1	MET
2	A	126	ARG
2	A	140	GLN
2	A	153	SER
2	A	158	LEU
2	A	251	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	A	280	ASN
2	A	458	GLU
2	A	462	THR
2	A	467	VAL
2	A	520	ASP
2	A	527	THR
2	A	586	CYS
2	A	611	VAL
2	A	652	ILE
2	B	18	VAL
2	B	48	GLN
2	B	57	ARG
2	B	80	SER
2	B	98	ILE
2	B	127	GLN
2	B	246	CYS
2	B	263	GLU
2	B	270	SER
2	B	277	THR
2	B	285	SER
2	B	318	LYS
2	B	372	CYS
2	B	481	VAL
2	B	497	LEU
2	B	508	GLN
2	B	569	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	93	HIS
2	A	317	GLN
2	A	656	HIS
2	B	50	GLN
2	B	76	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	ACY	B	803	-	1,3,3	3.60	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ACY	B	803	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	803	ACY	CH3-C	3.60	1.53	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	C	7/8 (87%)	-0.04	1 (14%) 4 3	24, 28, 62, 71	0
1	D	8/8 (100%)	-0.57	0 100 100	17, 24, 36, 44	0
2	A	630/659 (95%)	0.52	70 (11%) 7 7	14, 51, 104, 145	0
2	B	631/659 (95%)	0.10	26 (4%) 41 46	15, 35, 79, 138	0
All	All	1276/1334 (95%)	0.30	97 (7%) 17 18	14, 42, 95, 145	0

All (97) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	657	HIS	8.1
2	A	249	TYR	7.6
2	A	658	HIS	6.5
2	B	658	HIS	6.2
2	A	654	HIS	5.9
2	A	647	VAL	5.8
2	A	218	ALA	5.6
2	A	508	GLN	5.5
2	A	544	LYS	5.4
2	A	547	TYR	5.2
2	A	456	ASP	5.2
2	B	656	HIS	5.2
2	A	657	HIS	4.8
2	A	22	PRO	4.6
2	A	612	ARG	4.6
2	A	252	GLU	4.6
2	A	458	GLU	4.4
2	B	654	HIS	4.3
2	A	248	PRO	4.2
2	B	653	ALA	4.1
2	A	459	ASN	4.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	A	260	PRO	4.0
2	A	440	PHE	4.0
2	A	510	LYS	3.9
2	A	258	HIS	3.9
2	A	507	VAL	3.9
2	A	123	PRO	3.6
2	A	512	LEU	3.6
2	B	124	GLU	3.6
2	A	511	ASP	3.5
2	B	655	HIS	3.5
2	A	487	PRO	3.5
2	A	484	GLN	3.5
2	A	306	GLY	3.5
2	A	486	GLN	3.5
2	A	124	GLU	3.4
2	A	489	ARG	3.3
2	A	267	ASN	3.3
2	B	24	ARG	3.3
2	B	253	GLU	3.3
2	A	127	GLN	3.3
2	A	460	ASP	3.2
2	A	253	GLU	3.1
2	A	505	ARG	3.0
2	A	216	PHE	3.0
2	A	616	GLY	3.0
2	A	655	HIS	3.0
2	A	217	VAL	2.9
2	B	618	VAL	2.9
2	A	5	ILE	2.9
2	A	648	GLY	2.9
2	A	615	ARG	2.9
2	A	514	LYS	2.8
2	A	244	GLU	2.8
2	A	112	VAL	2.7
2	B	652	ILE	2.7
2	A	649	SER	2.7
2	A	656	HIS	2.6
2	B	392	ALA	2.6
2	A	215	ASN	2.6
2	A	34	ASN	2.6
2	B	612	ARG	2.6
2	B	394	ASN	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	A	455	ARG	2.5
2	A	546	ARG	2.5
2	B	21	LYS	2.5
2	A	264	HIS	2.4
2	A	504	ALA	2.4
2	B	123	PRO	2.4
2	B	619	ALA	2.4
2	B	252	GLU	2.4
2	A	221	PHE	2.4
2	A	517	ARG	2.4
2	B	81	VAL	2.4
2	A	617	ILE	2.4
2	A	650	GLY	2.3
2	A	453	LYS	2.3
2	B	283	ARG	2.3
2	B	649	SER	2.2
2	A	245	ALA	2.2
2	B	648	GLY	2.2
2	A	462	THR	2.2
2	A	543	LYS	2.2
2	A	116	LEU	2.2
2	B	112	VAL	2.1
2	A	515	ILE	2.1
2	A	257	LEU	2.1
2	A	488	PRO	2.1
2	B	620	PRO	2.1
1	C	7	DT	2.1
2	A	282	LYS	2.1
2	B	469	LYS	2.1
2	A	431	LYS	2.1
2	A	482	GLU	2.1
2	A	585	GLN	2.0
2	A	509	ASP	2.0
2	B	486	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CL	A	800	1/1	0.99	0.15	2.31	23,23,23,23	0
4	ACY	B	803	4/4	0.92	0.17	1.84	23,23,24,25	0
3	CL	A	801	1/1	1.00	0.13	0.18	20,20,20,20	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.