



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:08 AM GMT

PDB ID : 3O5N
Title : Tetrahydroquinoline carboxylates are potent inhibitors of the Shank PDZ domain, a putative target in autism disorders
Authors : Saupe, J.; Roske, Y.; Schillinger, C.; Kamdem, N.; Radetzki, S.; Diehl, A.; Oschkinat, H.; Krause, G.; Heinemann, U.; Rademann, J.
Deposited on : 2010-07-28
Resolution : 1.83 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	2634 (1.86-1.82)
Clashscore	102246	2862 (1.86-1.82)
Ramachandran outliers	100387	2831 (1.86-1.82)
Sidechain outliers	100360	2832 (1.86-1.82)
RSRZ outliers	91569	2639 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	112	<div><div>3%</div><div>47%32%9%•11%</div></div>
1	B	112	<div><div>4%</div><div>46%33%6%•13%</div></div>
1	C	112	<div><div>4%</div><div>47%34%••13%</div></div>
1	D	112	<div><div>21%</div><div>41%36%11%•11%</div></div>
1	E	112	<div><div>24%</div><div>40%33%8%•18%</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	112	
1	G	112	
1	H	112	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BR0	E	1	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6346 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SH3 and multiple ankyrin repeat domains protein 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	100	Total	C	N	O	S	0	0	0
			780	500	139	140	1			
1	B	98	Total	C	N	O	S	0	0	0
			764	489	137	137	1			
1	C	97	Total	C	N	O	S	0	0	0
			747	479	132	135	1			
1	D	100	Total	C	N	O	S	0	2	0
			792	507	145	139	1			
1	E	92	Total	C	N	O	S	0	1	0
			720	462	129	128	1			
1	F	97	Total	C	N	O	S	0	1	0
			760	486	138	135	1			
1	G	95	Total	C	N	O	S	0	1	0
			745	478	133	133	1			
1	H	93	Total	C	N	O	S	0	0	0
			726	467	128	130	1			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	633	GLY	-	expression tag	UNP Q4ACU6
A	634	ALA	-	expression tag	UNP Q4ACU6
A	635	ALA	-	expression tag	UNP Q4ACU6
A	636	SER	-	expression tag	UNP Q4ACU6
B	633	GLY	-	expression tag	UNP Q4ACU6
B	634	ALA	-	expression tag	UNP Q4ACU6
B	635	ALA	-	expression tag	UNP Q4ACU6
B	636	SER	-	expression tag	UNP Q4ACU6
C	633	GLY	-	expression tag	UNP Q4ACU6
C	634	ALA	-	expression tag	UNP Q4ACU6
C	635	ALA	-	expression tag	UNP Q4ACU6
C	636	SER	-	expression tag	UNP Q4ACU6
D	633	GLY	-	expression tag	UNP Q4ACU6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	634	ALA	-	expression tag	UNP Q4ACU6
D	635	ALA	-	expression tag	UNP Q4ACU6
D	636	SER	-	expression tag	UNP Q4ACU6
E	633	GLY	-	expression tag	UNP Q4ACU6
E	634	ALA	-	expression tag	UNP Q4ACU6
E	635	ALA	-	expression tag	UNP Q4ACU6
E	636	SER	-	expression tag	UNP Q4ACU6
F	633	GLY	-	expression tag	UNP Q4ACU6
F	634	ALA	-	expression tag	UNP Q4ACU6
F	635	ALA	-	expression tag	UNP Q4ACU6
F	636	SER	-	expression tag	UNP Q4ACU6
G	633	GLY	-	expression tag	UNP Q4ACU6
G	634	ALA	-	expression tag	UNP Q4ACU6
G	635	ALA	-	expression tag	UNP Q4ACU6
G	636	SER	-	expression tag	UNP Q4ACU6
H	633	GLY	-	expression tag	UNP Q4ACU6
H	634	ALA	-	expression tag	UNP Q4ACU6
H	635	ALA	-	expression tag	UNP Q4ACU6
H	636	SER	-	expression tag	UNP Q4ACU6

- Molecule 2 is (3AS,4R,9BR)-9-NITRO-3A,4,5,9B-TETRAHYDRO-3H-CYCLOPENTA[C]QUINOLINE-4,6-DICARBOXYLIC ACID (three-letter code: BR0) (formula: C₁₄H₁₂N₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	E	1	Total	C	N	O	0
			22	14	2	6	


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	53	Total 53	O 53	0	0
3	B	41	Total 41	O 41	0	0
3	C	34	Total 34	O 34	0	0
3	D	39	Total 39	O 39	0	0
3	E	23	Total 23	O 23	0	0
3	F	46	Total 46	O 46	0	0
3	G	14	Total 14	O 14	0	0
3	H	40	Total 40	O 40	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SH3 and multiple ankyrin repeat domains protein 3




- Molecule 1: SH3 and multiple ankyrin repeat domains protein 3





- Molecule 1: SH3 and multiple ankyrin repeat domains protein 3




- Molecule 1: SH3 and multiple ankyrin repeat domains protein 3




- Molecule 1: SH3 and multiple ankyrin repeat domains protein 3




- Molecule 1: SH3 and multiple ankyrin repeat domains protein 3



- Molecule 1: SH3 and multiple ankyrin repeat domains protein 3



- Molecule 1: SH3 and multiple ankyrin repeat domains protein 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.95Å 64.06Å 101.92Å 90.00° 90.09° 90.00°	Depositor
Resolution (Å)	33.97 – 1.83 33.97 – 1.83	Depositor EDS
% Data completeness (in resolution range)	88.7 (33.97-1.83) 96.1 (33.97-1.83)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.87 (at 1.83Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.233 , 0.283 0.273 , 0.317	Depositor DCC
R_{free} test set	3064 reflections (5.25%)	DCC
Wilson B-factor (Å ²)	30.2	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.8	EDS
Estimated twinning fraction	0.514 for H, K, L 0.486 for H, -K, -L 0.118 for h,-k,-l	Xtriage
Reported twinning fraction	0.514 for H, K, L 0.486 for H, -K, -L	Depositor
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 61804 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	6346	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.30 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 8.1796e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BR0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/793	0.82	0/1071
1	B	0.66	0/776	0.84	2/1047 (0.2%)
1	C	0.70	0/759	0.82	1/1027 (0.1%)
1	D	0.58	0/811	0.81	1/1094 (0.1%)
1	E	0.58	0/734	0.79	0/990
1	F	0.56	0/775	0.82	0/1046
1	G	0.72	0/760	0.85	0/1027
1	H	0.56	0/738	0.75	0/998
All	All	0.64	0/6146	0.81	4/8300 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	2
1	F	0	1
All	All	0	3

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	696	ALA	N-CA-C	5.95	127.06	111.00
1	B	648	LEU	CA-CB-CG	5.63	128.25	115.30
1	C	702	ASP	CB-CG-OD1	5.09	122.88	118.30
1	D	638	ASP	N-CA-C	5.08	124.70	111.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	686	SER	Peptide
1	D	687	VAL	Peptide
1	F	653	HIS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	780	0	798	44	0
1	B	764	0	780	59	0
1	C	747	0	751	37	0
1	D	792	0	815	68	0
1	E	720	0	745	65	1
1	F	760	0	779	65	0
1	G	745	0	760	44	0
1	H	726	0	740	56	0
2	E	22	0	10	9	0
3	A	53	0	0	7	0
3	B	41	0	0	9	0
3	C	34	0	0	5	0
3	D	39	0	0	7	1
3	E	23	0	0	5	0
3	F	46	0	0	12	0
3	G	14	0	0	3	0
3	H	40	0	0	4	0
All	All	6346	0	6178	405	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 33.

The worst 5 of 405 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:732:VAL:HG21	1:H:677:ALA:HB1	1.21	1.19
1:H:727:GLY:O	1:H:730:ARG:NE	1.74	1.18

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:730:ARG:HD2	1:E:678:PHE:CD2	1.80	1.16
1:D:730:ARG:HD2	1:E:678:PHE:HD2	1.01	1.16
1:D:687:VAL:HG12	1:D:688:ASP:H	1.09	1.15

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:718:LYS:NZ	3:D:29:HOH:O[2_657]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	96/112 (86%)	81 (84%)	7 (7%)	8 (8%)	1	0
1	B	94/112 (84%)	82 (87%)	9 (10%)	3 (3%)	5	0
1	C	93/112 (83%)	89 (96%)	1 (1%)	3 (3%)	5	0
1	D	98/112 (88%)	78 (80%)	13 (13%)	7 (7%)	1	0
1	E	89/112 (80%)	78 (88%)	6 (7%)	5 (6%)	2	0
1	F	94/112 (84%)	84 (89%)	8 (8%)	2 (2%)	9	1
1	G	92/112 (82%)	88 (96%)	3 (3%)	1 (1%)	17	5
1	H	89/112 (80%)	79 (89%)	8 (9%)	2 (2%)	8	1
All	All	745/896 (83%)	659 (88%)	55 (7%)	31 (4%)	3	0

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	692	VAL
1	A	693	ALA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	694	TRP
1	B	696	ALA
1	C	672	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	84/91 (92%)	75 (89%)	9 (11%)	8	1
1	B	81/91 (89%)	68 (84%)	13 (16%)	3	0
1	C	79/91 (87%)	71 (90%)	8 (10%)	9	1
1	D	84/91 (92%)	70 (83%)	14 (17%)	3	0
1	E	78/91 (86%)	64 (82%)	14 (18%)	2	0
1	F	81/91 (89%)	67 (83%)	14 (17%)	2	0
1	G	80/91 (88%)	72 (90%)	8 (10%)	9	1
1	H	78/91 (86%)	70 (90%)	8 (10%)	9	1
All	All	645/728 (89%)	557 (86%)	88 (14%)	5	0

5 of 88 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	704	LEU
1	E	659	VAL
1	H	661	ARG
1	D	723	LEU
1	E	638	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	708	ASN
1	D	729	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	682	GLN
1	D	719	GLN
1	E	649	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BR0	E	1	-	15,24,24	3.15	4 (26%)	13,36,36	2.87	4 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BR0	E	1	-	-	0/3/34/34	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1	BR0	CA-N	2.10	1.48	1.46
2	E	1	BR0	CB-CA	2.67	1.56	1.53
2	E	1	BR0	CD1-CAI	4.23	1.53	1.33
2	E	1	BR0	OAC-NAT	10.30	1.43	1.22

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1	BR0	CG-CD1-CAI	-6.86	107.13	112.46
2	E	1	BR0	CAK-CAI-CD1	-6.29	104.73	112.75
2	E	1	BR0	CAG-CAH-CAP	2.03	122.21	118.96
2	E	1	BR0	CB-CA-N	3.46	112.07	108.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	1	BR0	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	100/112 (89%)	0.74	3 (3%) 54 50	18, 28, 40, 44	0
1	B	98/112 (87%)	0.74	5 (5%) 32 28	22, 30, 37, 41	0
1	C	97/112 (86%)	0.71	4 (4%) 41 37	18, 28, 36, 38	0
1	D	100/112 (89%)	1.37	23 (23%) 1 1	28, 37, 44, 46	0
1	E	92/112 (82%)	1.48	27 (29%) 1 0	27, 38, 45, 48	0
1	F	97/112 (86%)	1.35	18 (18%) 2 1	30, 39, 45, 49	0
1	G	95/112 (84%)	0.68	2 (2%) 67 64	20, 28, 36, 49	0
1	H	93/112 (83%)	1.32	23 (24%) 1 0	27, 37, 46, 53	0
All	All	772/896 (86%)	1.05	105 (13%) 4 3	18, 33, 44, 53	0

The worst 5 of 105 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	678	PHE	7.2
1	D	738	VAL	5.6
1	D	672	PHE	5.4
1	F	691	GLY	4.8
1	D	689	VAL	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	BR0	E	1	22/22	0.84	0.21	0.35	51,52,52,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.