



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 04:48 AM GMT

PDB ID : 2O8C
Title : human MutSalpha (MSH2/MSH6) bound to ADP and an O6-methyl-guanine
T mispair
Authors : Warren, J.J.; Pohlhaus, T.J.; Changela, A.; Modrich, P.L.; Beese, L.S.
Deposited on : 2006-12-12
Resolution : 3.37 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1084 (3.46-3.30)
Clashscore	102246	1158 (3.46-3.30)
Ramachandran outliers	100387	1139 (3.46-3.30)
Sidechain outliers	100360	1138 (3.46-3.30)
RSRZ outliers	91569	1089 (3.46-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	15	<div> <div>7%</div> <div> <div></div> <div>73%</div> <div>27%</div> </div> </div>
2	F	15	<div> <div></div> <div> <div>93%</div> <div>7%</div> </div> </div>
3	A	934	<div> <div>9%</div> <div> <div></div> <div>81%</div> <div>8%</div> <div>11%</div> </div> </div>
4	B	1022	<div> <div>5%</div> <div> <div></div> <div>81%</div> <div>10%</div> <div>9%</div> </div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 14571 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*GP*AP*AP*CP*CP*GP*CP*(6OG)P*CP*GP*C P*TP*AP*GP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	15	Total	C	N	O	P	0	0	0
			308	146	62	86	14			

- Molecule 2 is a DNA chain called 5'-D(*CP*CP*TP*AP*GP*CP*GP*TP*GP*CP*GP*G P*TP*TP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	15	Total	C	N	O	P	0	0	0
			303	145	53	91	14			

- Molecule 3 is a protein called DNA mismatch repair protein Msh2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	830	Total	C	N	O	S	0	0	0
			6439	4085	1092	1228	34			

- Molecule 4 is a protein called DNA mismatch repair protein MSH6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	932	Total	C	N	O	S	0	0	0
			7443	4721	1277	1394	51			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	339	MET	-	INITIATING METHIONINE	UNP P52701
B	340	GLY	-	CLONING ARTIFACT	UNP P52701

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Mg 1 1	0	0
5	A	1	Total Mg 1 1	0	0

- Molecule 6 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C N O P 27 10 5 10 2	0	0
6	B	1	Total C N O P 27 10 5 10 2	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total O 1 1	0	0
7	B	20	Total O 20 20	0	0
7	F	1	Total O 1 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(*GP*AP*AP*CP*CP*GP*CP*(6OG)P*CP*GP*CP*TP*AP*GP*G)-3',



- Molecule 2: 5'-D(*CP*CP*TP*AP*GP*CP*GP*TP*GP*CP*GP*GP*TP*TP*C)-3'



- Molecule 3: DNA mismatch repair protein Msh2





4 Data and refinement statistics

Property	Value	Source
Space group	P 43 3 2	Depositor
Cell constants a, b, c, α , β , γ	259.81Å 259.81Å 259.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 3.37 48.25 – 3.37	Depositor EDS
% Data completeness (in resolution range)	99.1 (20.00-3.37) 99.1 (48.25-3.37)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	0.20	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 3.40Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.256 , 0.290 0.241 , 0.272	Depositor DCC
R_{free} test set	2126 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	92.3	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 145.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtriage
Outliers	0 of 42382 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	14571	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 6OG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	0.55	0/319	1.18	4/488 (0.8%)
2	F	0.59	0/338	1.23	1/520 (0.2%)
3	A	0.31	0/6539	0.54	0/8828
4	B	0.33	0/7587	0.57	0/10226
All	All	0.33	0/14783	0.61	5/20062 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	4	DC	O4'-C4'-C3'	-8.04	101.18	106.00
2	F	21	DC	P-O3'-C3'	6.36	127.33	119.70
1	E	4	DC	P-O3'-C3'	5.61	126.44	119.70
1	E	14	DG	O4'-C1'-N9	5.58	111.91	108.00
1	E	4	DC	O4'-C1'-N1	5.57	111.90	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	308	0	170	1	0
2	F	303	0	171	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	6439	0	6410	37	0
4	B	7443	0	7415	68	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	27	0	12	0	0
6	B	27	0	12	0	0
7	A	1	0	0	0	0
7	B	20	0	0	0	0
7	F	1	0	0	0	0
All	All	14571	0	14190	104	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 104 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:385:ARG:HD2	4:B:386:PRO:HD2	1.66	0.76
4:B:380:ASP:HB2	4:B:384:ARG:H	1.60	0.66
3:A:359:ARG:NH2	3:A:691:ILE:O	2.28	0.65
3:A:5:PRO:HB3	3:A:81:SER:HB3	1.77	0.65
4:B:897:ASN:HB3	4:B:901:ARG:HE	1.62	0.64

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	816/934 (87%)	736 (90%)	73 (9%)	7 (1%)	21	63
4	B	912/1022 (89%)	836 (92%)	66 (7%)	10 (1%)	17	58
All	All	1728/1956 (88%)	1572 (91%)	139 (8%)	17 (1%)	19	60

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	146	GLY
3	A	28	GLU
4	B	667	ASP
4	B	898	PRO
3	A	672	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	693/808 (86%)	693 (100%)	0	100	100
4	B	819/899 (91%)	819 (100%)	0	100	100
All	All	1512/1707 (89%)	1512 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	377	GLN
4	B	534	ASN
4	B	989	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	6OG	E	8	1,2	16,25,26	1.43	1 (6%)	21,36,39	2.93	5 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	6OG	E	8	1,2	-	2/5/23/24	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	8	6OG	C6-N1	4.58	1.39	1.31

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	8	6OG	C5-C6-N1	-7.26	110.99	123.81
1	E	8	6OG	C1'-N9-C4	-2.49	122.94	127.16
1	E	8	6OG	N3-C2-N1	-2.45	123.71	127.44
1	E	8	6OG	C2-N1-C6	7.22	125.96	116.03
1	E	8	6OG	C-O6-C6	7.22	124.75	117.26

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	E	8	6OG	C-O6-C6-N1
1	E	8	6OG	C-O6-C6-C5

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	ADP	A	936	5	22,29,29	1.06	1 (4%)	27,45,45	1.79	4 (14%)
6	ADP	B	202	5	22,29,29	1.03	1 (4%)	27,45,45	1.90	3 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ADP	A	936	5	-	0/12/32/32	0/3/3/3
6	ADP	B	202	5	-	0/12/32/32	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	202	ADP	C5-C4	3.23	1.47	1.40
6	A	936	ADP	C5-C4	3.26	1.47	1.40

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	202	ADP	N3-C2-N1	-7.05	123.50	128.89
6	A	936	ADP	N3-C2-N1	-6.54	123.89	128.89
6	B	202	ADP	PA-O3A-PB	-3.78	119.98	132.67
6	B	202	ADP	C4-C5-N7	-3.31	106.43	109.48
6	A	936	ADP	C4-C5-N7	-3.12	106.61	109.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	14/15 (93%)	0.68	1 (7%) 19 20	68, 78, 91, 91	0
2	F	15/15 (100%)	0.41	0 100 100	65, 72, 89, 91	0
3	A	830/934 (88%)	0.67	84 (10%) 9 10	20, 73, 76, 83	0
4	B	932/1022 (91%)	0.42	49 (5%) 30 31	30, 73, 82, 89	0
All	All	1791/1986 (90%)	0.54	134 (7%) 17 19	20, 73, 80, 91	0

The worst 5 of 134 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	206	THR	7.8
4	B	968	VAL	6.7
1	E	15	DG	6.4
4	B	954	TYR	5.8
3	A	541	THR	5.4

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	6OG	E	8	23/24	0.91	0.27	-	68,68,69,69	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	ADP	A	936	27/27	0.89	0.23	-0.39	70,70,75,75	0
6	ADP	B	202	27/27	0.90	0.19	-0.86	82,82,83,83	0
5	MG	B	102	1/1	0.96	0.09	-	58,58,58,58	0
5	MG	A	935	1/1	0.97	0.30	-	73,73,73,73	0

6.5 Other polymers [i](#)

There are no such residues in this entry.