



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:13 PM GMT

PDB ID : 4O8G  
Title : Structure of Infrared Fluorescent Protein 1.4  
Authors : Bhattacharya, S.; Forest, K.T.  
Deposited on : 2013-12-27  
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1226 (1.66-1.66)
Clashscore	102246	1323 (1.66-1.66)
Ramachandran outliers	100387	1295 (1.66-1.66)
Sidechain outliers	100360	1295 (1.66-1.66)
RSRZ outliers	91569	1227 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	<div> <div>9%</div> <div>89%</div> <div>5% 5%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2745 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bacteriophytochrome.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	311	2432	1553	429	439	11	0	10	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	SER	CONFLICT	UNP Q9RZA4
A	54	VAL	MET	ENGINEERED MUTATION	UNP Q9RZA4
A	119	ALA	GLY	ENGINEERED MUTATION	UNP Q9RZA4
A	135	ILE	THR	CONFLICT	UNP Q9RZA4
A	186	MET	VAL	ENGINEERED MUTATION	UNP Q9RZA4
A	195	MET	LEU	ENGINEERED MUTATION	UNP Q9RZA4
A	196	GLN	HIS	ENGINEERED MUTATION	UNP Q9RZA4
A	207	HIS	ASP	ENGINEERED MUTATION	UNP Q9RZA4
A	208	THR	ILE	ENGINEERED MUTATION	UNP Q9RZA4
A	288	VAL	ALA	ENGINEERED MUTATION	UNP Q9RZA4
A	307	GLU	TYR	ENGINEERED MUTATION	UNP Q9RZA4
A	311	LYS	LEU	ENGINEERED MUTATION	UNP Q9RZA4
A	314	GLY	LEU	ENGINEERED MUTATION	UNP Q9RZA4
A	318	ARG	VAL	ENGINEERED MUTATION	UNP Q9RZA4
A	322	LEU	ALA	CONFLICT	UNP Q9RZA4
A	323	GLU	ASP	CONFLICT	UNP Q9RZA4
A	324	HIS	-	EXPRESSION TAG	UNP Q9RZA4
A	325	HIS	-	EXPRESSION TAG	UNP Q9RZA4
A	326	HIS	-	EXPRESSION TAG	UNP Q9RZA4
A	327	HIS	-	EXPRESSION TAG	UNP Q9RZA4
A	328	HIS	-	EXPRESSION TAG	UNP Q9RZA4
A	329	HIS	-	EXPRESSION TAG	UNP Q9RZA4


- Molecule 2 is 3-[2-[(Z)-[3-(2-CARBOXYETHYL)-5-[(Z)-(4-ETHENYL-3-METHYL-5-OXIDANYLIDENE-PYRROL-2-YLIDENE)METHYL]-4-METHYL-PYRROL-1-IUM-2-YLIDENE]METHYL]-5-[(Z)-[(3E)-3-ETHYLIDENE-4-METHYL-5-OXIDANYLIDENE-PYRROLIDIN-2-YLIDENE]METHYL]-4-METHYL-1H-PYRROL-3-YL]PROPANOIC ACID

(three-letter code: LBV) (formula:  $C_{33}H_{37}N_4O_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	1
			43	33	4	6		

- Molecule 3 is 3-[2-[(Z)-[5-[(Z)-(4-ETHENYL-3-METHYL-5-OXIDANYLIDENE-PYRROL-2-YLIDENE)METHYL]-3-(3-HYDROXY-3-OXOPROPYL)-4-METHYL-PYRROL-1-IUM-2-YLIDENE]METHYL]-5-[(Z)-[(3E,4R)-3-ETHYLIDENE-4-METHYL-5-OXIDANYLIDENE-PYRROLIDIN-2-YLIDENE]METHYL]-4-METHYL-1H-PYRROL-3-YL]PROPANOIC ACID (three-letter code: LBW) (formula:  $C_{33}H_{37}N_4O_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	1
			43	33	4	6		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	227	Total	O	0	0
			227	227		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Bacteriophytochrome



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.16Å 53.23Å 66.80Å 90.00° 90.60° 90.00°	Depositor
Resolution (Å)	24.04 – 1.65 24.04 – 1.65	Depositor EDS
% Data completeness (in resolution range)	93.4 (24.04-1.65) 93.4 (24.04-1.65)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.03 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.185 , 0.211 0.195 , 0.220	Depositor DCC
$R_{free}$ test set	1876 reflections (5.20%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.5	Xtriage
Anisotropy	0.050	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 46.1	EDS
Estimated twinning fraction	0.007 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.009 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.014 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.013 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.016 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 37943 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2745	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LBW, LBV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.47	0/2523	0.71	4/3447 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	70	ARG	NE-CZ-NH1	6.47	123.54	120.30
1	A	70	ARG	NE-CZ-NH2	-6.47	117.07	120.30
1	A	254	ARG	NE-CZ-NH2	6.10	123.35	120.30
1	A	254	ARG	NE-CZ-NH1	-5.16	117.72	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2432	0	2462	11	0
2	A	43	0	33	1	0
3	A	43	0	33	3	0
4	A	227	0	0	2	0
All	All	2745	0	2528	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 3.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:MET:HB2	1:A:288[A]:VAL:CG1	2.29	0.62
1:A:259:MET:HG2	4:A:698:HOH:O	2.06	0.55
1:A:174:MET:HB2	1:A:288[A]:VAL:HG13	1.90	0.54
1:A:288[A]:VAL:HG23	4:A:510:HOH:O	2.08	0.54
3:A:402[B]:LBW:CBA	3:A:402[B]:LBW:HMA1	2.42	0.49
1:A:290:HIS:CE1	3:A:402[B]:LBW:O_D	2.66	0.49
1:A:203:PHE:CZ	2:A:401[A]:LBV:HBD2	2.48	0.49
1:A:143:ALA:O	1:A:147[A]:LEU:HD23	2.13	0.49
1:A:98:GLN:HG2	1:A:116[B]:HIS:NE2	2.29	0.47
1:A:196:GLN:OE1	1:A:196:GLN:HA	2.17	0.45
1:A:193:GLU:HA	1:A:194:GLY:HA2	1.79	0.43
1:A:147[A]:LEU:HD22	1:A:159:VAL:HG12	2.02	0.41
3:A:402[B]:LBW:HBD1	3:A:402[B]:LBW:O_D	2.20	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	315/329 (96%)	312 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/265 (98%)	258 (99%)	3 (1%)	80	64

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	105	TRP
1	A	222	ARG
1	A	323	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	LBV	A	401[A]	1	35,46,46	4.29	16 (45%)	37,67,67	1.70	7 (18%)
3	LBW	A	402[B]	1	35,46,46	4.05	17 (48%)	37,67,67	1.63	9 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LBV	A	401[A]	1	-	2/22/74/74	0/4/4/4
3	LBW	A	402[B]	1	-	2/22/74/74	0/4/4/4

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401[A]	LBV	C3D-C2D	-6.36	1.23	1.36
3	A	402[B]	LBW	C3D-C2D	-5.90	1.24	1.36
3	A	402[B]	LBW	C4A-C3A	-4.94	1.35	1.45
2	A	401[A]	LBV	C4A-C3A	-4.53	1.36	1.45
2	A	401[A]	LBV	C1D-C2D	-4.07	1.37	1.45
3	A	402[B]	LBW	C1D-C2D	-3.65	1.38	1.45
3	A	402[B]	LBW	O_D-C4D	-2.58	1.18	1.23
3	A	402[B]	LBW	CMA-C2A	-2.35	1.45	1.53
3	A	402[B]	LBW	C4A-N_A	-2.30	1.33	1.37
2	A	401[A]	LBV	CMA-C2A	-2.12	1.46	1.53
2	A	401[A]	LBV	C3D-C4D	2.04	1.53	1.47
3	A	402[B]	LBW	CHB-C4A	2.40	1.39	1.34
2	A	401[A]	LBV	C4C-C3C	2.46	1.51	1.45
3	A	402[B]	LBW	C4C-C3C	2.64	1.51	1.45
3	A	402[B]	LBW	C1D-N_D	2.82	1.42	1.37
3	A	402[B]	LBW	C4B-CHC	2.99	1.51	1.40
2	A	401[A]	LBV	C4B-CHC	3.18	1.52	1.40
2	A	401[A]	LBV	CHB-C4A	3.40	1.41	1.34
3	A	402[B]	LBW	C1B-CHB	3.41	1.53	1.40
2	A	401[A]	LBV	C1D-N_D	3.55	1.44	1.37
2	A	401[A]	LBV	C1B-CHB	3.90	1.55	1.40
2	A	401[A]	LBV	CHD-C4C	3.97	1.50	1.40
3	A	402[B]	LBW	CHD-C4C	4.07	1.50	1.40
3	A	402[B]	LBW	CAA-C3A	4.07	1.38	1.33
2	A	401[A]	LBV	CHC-C1C	4.22	1.38	1.35
3	A	402[B]	LBW	CHC-C1C	5.13	1.39	1.35

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401[A]	LBV	CAA-C3A	5.80	1.40	1.33
3	A	402[B]	LBW	C4D-N_D	8.60	1.57	1.37
2	A	401[A]	LBV	C4D-N_D	9.22	1.58	1.37
3	A	402[B]	LBW	C2A-C1A	10.45	1.61	1.52
2	A	401[A]	LBV	C2A-C1A	10.80	1.62	1.52
3	A	402[B]	LBW	C1A-N_A	13.48	1.54	1.37
2	A	401[A]	LBV	C1A-N_A	14.67	1.56	1.37

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402[B]	LBW	C1D-N_D-C4D	-4.03	104.83	110.73
2	A	401[A]	LBV	C1D-N_D-C4D	-3.79	105.18	110.73
3	A	402[B]	LBW	CBB-CAB-C3B	-3.50	106.25	112.53
2	A	401[A]	LBV	CHB-C4A-C3A	-3.20	121.15	127.21
3	A	402[B]	LBW	CHB-C4A-N_A	-2.55	121.44	130.87
3	A	402[B]	LBW	CHD-C1D-N_D	-2.30	121.19	126.16
2	A	401[A]	LBV	C2C-C1C-N_C	-2.12	106.56	109.86
3	A	402[B]	LBW	CHB-C4A-C3A	-2.12	123.20	127.21
3	A	402[B]	LBW	C2C-C1C-N_C	-2.10	106.61	109.86
2	A	401[A]	LBV	CBB-CAB-C3B	-2.06	108.83	112.53
3	A	402[B]	LBW	C2D-C1D-N_D	2.14	110.10	107.00
2	A	401[A]	LBV	C3A-C4A-N_A	2.23	110.24	107.32
2	A	401[A]	LBV	CHD-C4C-C3C	2.26	130.40	124.88
3	A	402[B]	LBW	C3A-C4A-N_A	2.66	110.81	107.32
3	A	402[B]	LBW	O_A-C1A-N_A	3.02	128.49	124.83
2	A	401[A]	LBV	O_A-C1A-N_A	4.75	130.57	124.83

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402[B]	LBW	C1D-CHD-C4C-C3C
2	A	401[A]	LBV	C1D-CHD-C4C-C3C
3	A	402[B]	LBW	C1D-CHD-C4C-N_C
2	A	401[A]	LBV	C1D-CHD-C4C-N_C

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401[A]	LBV	1	0
3	A	402[B]	LBW	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	311/329 (94%)	0.32	30 (9%) 10 8	10, 17, 41, 77	4 (1%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	105	TRP	7.1
1	A	96	ALA	5.7
1	A	130	GLU	5.0
1	A	4	ASP	4.8
1	A	197	ALA	4.4
1	A	138	HIS	4.3
1	A	6	LEU	4.3
1	A	106	PRO	3.7
1	A	322	LEU	3.6
1	A	95	ASP	3.3
1	A	5	PRO	3.2
1	A	139	ALA	3.2
1	A	196	GLN	3.2
1	A	140	LEU	3.2
1	A	321	ALA	3.1
1	A	110	HIS	3.0
1	A	193	GLU	3.0
1	A	93	CYS	3.0
1	A	194	GLY	2.9
1	A	109	GLY	2.8
1	A	141	ARG	2.8
1	A	288[A]	VAL	2.6
1	A	92	GLY	2.6
1	A	195	MET	2.5
1	A	94	PRO	2.5
1	A	80	GLU	2.5
1	A	104	ASP	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	90[A]	PRO	2.2
1	A	192	ARG	2.2
1	A	124	LEU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	LBV	A	401[A]	43/43	0.95	0.11	0.39	10,11,14,17	43
3	LBW	A	402[B]	43/43	0.95	0.11	0.39	9,10,13,17	43

## 6.5 Other polymers [i](#)

There are no such residues in this entry.