



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 26, 2016 – 04:09 PM BST

PDB ID : 1ONV
Title : NMR Structure of a Complex Containing the TFIIF Subunit RAP74 and the RNAP II CTD Phosphatase FCP1
Authors : Nguyen, B.D.; Abbott, K.L.; Potempa, K.; Kobor, M.S.; Archambault, J.; Greenblatt, J.; Legault, P.; Omichinski, J.G.
Deposited on : 2003-03-02

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	82	 55% 24% • 18%
2	B	83	 14% 6% 5% 75%

2 Ensemble composition and analysis ⓘ

This entry contains 20 models. Model 12 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average, minimized average structure*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:453-A:517, B:945-B:961 (82)	0.54	12

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters. No single-model clusters were found.

Cluster number	Models
1	2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 15, 16, 17, 20
2	13, 19
3	14, 18
4	1, 9

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1437 atoms, of which 733 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Transcription initiation factor IIF, alpha subunit.

Mol	Chain	Residues	Atoms						Trace
1	A	67	Total	C	H	N	O	S	0
			1145	346	593	101	102	3	

- Molecule 2 is a protein called serine phosphatase FCP1a.


Mol	Chain	Residues	Atoms						Trace
2	B	21	Total	C	H	N	O	S	0
			292	89	140	23	38	2	

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Transcription initiation factor IIF, alpha subunit




- Molecule 2: serine phosphatase FCP1a



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 12. Colouring as in section 4.1 above.

- Molecule 1: Transcription initiation factor IIF, alpha subunit



- Molecule 2: serine phosphatase FCP1a



ASP	PRO
GLU	GLU
8941	GLN
8942	GLU
8943	GLU
8944	GLU
8949	PRO
	GLN
8953	PRO
	ARG
8957	LYS
8958	PRO
8959	GLY
8960	THR
8961	ARG
	ARG
	GLU
	ARG
	THR
	LEU
	GLY
	ALA
	PRO
	ALA
	ALA
	SER
	SER
	GLU
	ARG
	SER
	ALA
	ALA
	SER
	GLU
	SER
	SER
	ARG
	GLU
	SER
	SER
	ASN
	GLU
	ASP
	GLU
	ALA
	ALA
	SER
	GLU
	SER
	SER
	ARG
	GLU
	SER
	SER
	ASN
	GLU

5 Refinement protocol and experimental data overview

The models were refined using the following method: *Torsion Angle Dynamics*.

Of the 70 calculated structures, 20 were deposited, based on the following criterion: *structures with acceptable covalent geometry, structures with the least restraint violations, structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure solution	1.0
CNS	refinement	modified CNS with conformational database potential

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	537	580	579	27±5
2	B	130	122	122	15±5
All	All	13340	14040	14020	588

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

5 of 260 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:474:LEU:HD21	1:A:490:VAL:HG22	1.05	1.27	19	2
2:B:957:LEU:HD23	2:B:960:LEU:HD21	1.03	1.22	1	1
1:A:500:LEU:HD21	1:A:515:LEU:HD11	0.95	1.32	2	3
2:B:957:LEU:HD11	2:B:960:LEU:HD13	0.95	1.38	16	2
1:A:497:LEU:HD13	2:B:953:LEU:HD21	0.92	1.42	7	12

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	64/82 (78%)	60±2 (94±2%)	3±1 (5±2%)	1±1 (1±1%)	23	69
2	B	16/83 (19%)	16±1 (99±3%)	0±1 (1±3%)	0±0 (0±0%)	100	100
All	All	1600/3300 (48%)	1521 (95%)	65 (4%)	14 (1%)	26	73

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	479	THR	7
1	A	481	LYS	4
1	A	478	GLN	2
1	A	508	ASN	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	62/77 (81%)	58±2 (93±3%)	4±2 (7±3%)	23	68
2	B	13/66 (20%)	12±1 (92±5%)	1±1 (8±5%)	18	63
All	All	1500/2860 (52%)	1389 (93%)	111 (7%)	22	67

5 of 39 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	466	LYS	11
1	A	515	LEU	9
2	B	960	LEU	8
1	A	498	LYS	8
1	A	496	ILE	7

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided