



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 04:54 AM GMT

PDB ID : 2ONC  
Title : Crystal structure of human DPP-4  
Authors : Feng, J.; Zhang, Z.; Wallace, M.B.; Stafford, J.A.; Kaldor, S.W.; Kassel, D.B.; Navre, M.; Shi, L.; Skene, R.J.; Asakawa, T.; Takeuchi, K.; Xu, R.; Webb, D.R.; Gwaltney, S.L.  
Deposited on : 2007-01-23  
Resolution : 2.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4549 (2.58-2.50)
Clashscore	102246	5292 (2.58-2.50)
Ramachandran outliers	100387	5194 (2.58-2.50)
Sidechain outliers	100360	5196 (2.58-2.50)
RSRZ outliers	91569	4561 (2.58-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	731	<div> <div>5%</div> <div>85%</div> <div>12%</div> <div>..</div> </div>
1	B	731	<div> <div>3%</div> <div>84%</div> <div>15%</div> <div>.</div> </div>
1	C	731	<div> <div>7%</div> <div>86%</div> <div>11%</div> <div>..</div> </div>
1	D	731	<div> <div>10%</div> <div>85%</div> <div>13%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	807	-	-	-	X
2	NAG	C	807	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 25016 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl peptidase 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	724	Total	C	N	O	S	0	1	0
			5925	3807	973	1119	26			
1	B	728	Total	C	N	O	S	0	1	0
			5958	3826	981	1125	26			
1	C	723	Total	C	N	O	S	0	1	0
			5918	3802	972	1118	26			
1	D	722	Total	C	N	O	S	0	0	0
			5907	3795	969	1117	26			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	36	HIS	-	EXPRESSION TAG	UNP P27487
A	37	HIS	-	EXPRESSION TAG	UNP P27487
A	38	ALA	-	EXPRESSION TAG	UNP P27487
A	39	SER	-	EXPRESSION TAG	UNP P27487
A	40	ALA	-	EXPRESSION TAG	UNP P27487
B	36	HIS	-	EXPRESSION TAG	UNP P27487
B	37	HIS	-	EXPRESSION TAG	UNP P27487
B	38	ALA	-	EXPRESSION TAG	UNP P27487
B	39	SER	-	EXPRESSION TAG	UNP P27487
B	40	ALA	-	EXPRESSION TAG	UNP P27487
C	36	HIS	-	EXPRESSION TAG	UNP P27487
C	37	HIS	-	EXPRESSION TAG	UNP P27487
C	38	ALA	-	EXPRESSION TAG	UNP P27487
C	39	SER	-	EXPRESSION TAG	UNP P27487
C	40	ALA	-	EXPRESSION TAG	UNP P27487
D	36	HIS	-	EXPRESSION TAG	UNP P27487
D	37	HIS	-	EXPRESSION TAG	UNP P27487
D	38	ALA	-	EXPRESSION TAG	UNP P27487
D	39	SER	-	EXPRESSION TAG	UNP P27487
D	40	ALA	-	EXPRESSION TAG	UNP P27487

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	D	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	B	2	Total	C	N	O	0	0
			28	16	2	10		
3	C	2	Total	C	N	O	0	0
			28	16	2	10		
3	D	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 4 is 2-({2-[(3R)-3-AMINOPIPERIDIN-1-YL]-4-OXOQUINAZOLIN-3(4H)-YL}METHYL)BENZONITRILE (three-letter code: SY1) (formula: C<sub>21</sub>H<sub>21</sub>N<sub>5</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			27	21	5	1		
4	A	1	Total	C	N	O	0	0
			27	21	5	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			27	21	5	1		
4	B	1	Total	C	N	O	0	0
			27	21	5	1		
4	C	1	Total	C	N	O	0	0
			27	21	5	1		
4	C	1	Total	C	N	O	0	0
			27	21	5	1		
4	D	1	Total	C	N	O	0	0
			27	21	5	1		
4	D	1	Total	C	N	O	0	0
			27	21	5	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	227	Total	O	0	0
			227	227		
5	B	213	Total	O	0	0
			213	213		
5	C	219	Total	O	0	0
			219	219		
5	D	97	Total	O	0	0
			97	97		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Dipeptidyl peptidase 4




#### • Molecule 1: Dipeptidyl peptidase 4




#### • Molecule 1: Dipeptidyl peptidase 4







• Molecule 1: Dipeptidyl peptidase 4



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	122.36Å 123.70Å 145.36Å 90.00° 114.89° 90.00°	Depositor
Resolution (Å)	50.00 – 2.55 38.65 – 2.55	Depositor EDS
% Data completeness (in resolution range)	92.8 (50.00-2.55) 92.8 (38.65-2.55)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.61 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.195 , 0.254 0.195 , 0.254	Depositor DCC
$R_{free}$ test set	5942 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	46.8	Xtriage
Anisotropy	0.526	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 49.5	EDS
Estimated twinning fraction	0.012 for h,-k,-h-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 118368 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	25016	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SY1, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.44	0/6096	0.59	2/8291 (0.0%)
1	B	0.42	0/6135	0.58	0/8344
1	C	0.43	0/6092	0.59	2/8285 (0.0%)
1	D	0.40	0/6077	0.55	0/8266
All	All	0.42	0/24400	0.58	4/33186 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	316	LEU	CA-CB-CG	5.45	127.84	115.30
1	C	57	LEU	CA-CB-CG	5.37	127.66	115.30
1	A	316	LEU	CA-CB-CG	5.09	127.00	115.30
1	A	142	LEU	CA-CB-CG	5.03	126.88	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5925	0	5639	53	0
1	B	5958	0	5670	58	0
1	C	5918	0	5635	43	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	5907	0	5621	46	0
2	A	56	0	52	0	0
2	B	56	0	52	0	0
2	C	56	0	52	0	0
2	D	28	0	26	0	0
3	A	56	0	50	1	0
3	B	28	0	25	0	0
3	C	28	0	25	0	0
3	D	28	0	25	0	0
4	A	54	0	42	6	0
4	B	54	0	42	6	0
4	C	54	0	42	6	0
4	D	54	0	42	5	0
5	A	227	0	0	5	0
5	B	213	0	0	5	0
5	C	219	0	0	0	0
5	D	97	0	0	2	0
All	All	25016	0	23040	220	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 220 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:153:GLN:HE22	1:B:170:ASN:H	1.07	1.00
1:C:153:GLN:HE22	1:C:170:ASN:H	1.13	0.95
1:A:153:GLN:HE22	1:A:170:ASN:H	1.20	0.89
4:B:800:SY1:H52	4:B:800:SY1:H191	1.57	0.87
1:D:153:GLN:HE22	1:D:170:ASN:H	1.25	0.81

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	720/731 (98%)	681 (95%)	37 (5%)	2 (0%)	46	66
1	B	725/731 (99%)	686 (95%)	37 (5%)	2 (0%)	46	66
1	C	718/731 (98%)	678 (94%)	39 (5%)	1 (0%)	56	76
1	D	716/731 (98%)	671 (94%)	44 (6%)	1 (0%)	56	76
All	All	2879/2924 (98%)	2716 (94%)	157 (6%)	6 (0%)	52	73

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	85	ASN
1	A	87	SER
1	B	463	LYS
1	B	83	TYR
1	C	536	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	648/654 (99%)	611 (94%)	37 (6%)	25	44
1	B	652/654 (100%)	616 (94%)	36 (6%)	27	46
1	C	648/654 (99%)	614 (95%)	34 (5%)	29	49
1	D	646/654 (99%)	618 (96%)	28 (4%)	35	59
All	All	2594/2616 (99%)	2459 (95%)	135 (5%)	29	49

5 of 135 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	498	SER
1	C	94	THR
1	D	489	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	504	LEU
1	B	566	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 58 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	506	ASN
1	C	169	ASN
1	D	506	ASN
1	B	572	ASN
1	B	710	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

10 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	805	3	14,14,15	0.62	0	15,19,21	1.41	2 (13%)
3	NAG	A	806	3	14,14,15	0.49	0	15,19,21	0.90	0
3	NAG	A	808	1,3	14,14,15	0.63	0	15,19,21	0.65	0
3	NAG	A	809	3	14,14,15	0.46	0	15,19,21	1.11	1 (6%)
3	NAG	B	805	1,3	14,14,15	0.49	0	15,19,21	0.75	1 (6%)
3	NAG	B	806	3	14,14,15	0.54	0	15,19,21	1.00	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	C	804	1,3	14,14,15	0.54	0	15,19,21	1.53	2 (13%)
3	NAG	C	805	3	14,14,15	0.47	0	15,19,21	0.84	1 (6%)
3	NAG	D	803	1,3	14,14,15	0.44	0	15,19,21	0.92	1 (6%)
3	NAG	D	804	3	14,14,15	0.49	0	15,19,21	1.14	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	805	3	-	0/6/23/26	0/1/1/1
3	NAG	A	806	3	-	0/6/23/26	0/1/1/1
3	NAG	A	808	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	809	3	-	0/6/23/26	0/1/1/1
3	NAG	B	805	1,3	-	0/6/23/26	0/1/1/1
3	NAG	B	806	3	-	0/6/23/26	0/1/1/1
3	NAG	C	804	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	805	3	-	0/6/23/26	0/1/1/1
3	NAG	D	803	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	804	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	804	NAG	C2-N2-C7	-3.49	118.56	123.04
3	B	805	NAG	C1-O5-C5	2.03	114.83	112.25
3	C	805	NAG	C1-O5-C5	2.19	115.03	112.25
3	B	806	NAG	C4-C3-C2	2.22	114.67	111.23
3	D	803	NAG	C1-O5-C5	2.60	115.54	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	805	NAG	1	0

## 5.6 Ligand geometry ⓘ

22 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	SY1	A	800	-	28,30,30	2.28	3 (10%)	29,42,42	2.02	6 (20%)
4	SY1	A	801	-	28,30,30	2.32	4 (14%)	29,42,42	1.98	6 (20%)
2	NAG	A	802	1	14,14,15	0.59	0	15,19,21	1.09	2 (13%)
2	NAG	A	803	1	14,14,15	0.59	0	15,19,21	1.07	1 (6%)
2	NAG	A	804	1	14,14,15	0.52	0	15,19,21	0.95	1 (6%)
2	NAG	A	807	1	14,14,15	0.58	0	15,19,21	1.19	1 (6%)
4	SY1	B	800	-	28,30,30	2.30	2 (7%)	29,42,42	1.80	6 (20%)
4	SY1	B	801	-	28,30,30	2.34	3 (10%)	29,42,42	1.40	4 (13%)
2	NAG	B	802	1	14,14,15	0.60	0	15,19,21	0.98	1 (6%)
2	NAG	B	803	1	14,14,15	0.57	0	15,19,21	1.21	1 (6%)
2	NAG	B	804	1	14,14,15	0.51	0	15,19,21	0.94	1 (6%)
2	NAG	B	807	1	14,14,15	0.49	0	15,19,21	1.15	1 (6%)
4	SY1	C	800	-	28,30,30	2.33	2 (7%)	29,42,42	1.84	6 (20%)
4	SY1	C	801	-	28,30,30	2.47	4 (14%)	29,42,42	2.53	6 (20%)
2	NAG	C	802	1	14,14,15	0.59	0	15,19,21	0.96	0
2	NAG	C	803	1	14,14,15	0.49	0	15,19,21	1.08	1 (6%)
2	NAG	C	806	1	14,14,15	0.59	0	15,19,21	0.99	1 (6%)
2	NAG	C	807	1	14,14,15	0.59	0	15,19,21	1.25	1 (6%)
4	SY1	D	800	-	28,30,30	2.32	2 (7%)	29,42,42	1.61	6 (20%)
4	SY1	D	801	-	28,30,30	2.31	4 (14%)	29,42,42	2.20	7 (24%)
2	NAG	D	802	1	14,14,15	0.61	0	15,19,21	1.12	1 (6%)
2	NAG	D	805	1	14,14,15	0.49	0	15,19,21	1.02	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SY1	A	800	-	-	0/10/20/20	0/4/4/4
4	SY1	A	801	-	-	0/10/20/20	0/4/4/4
2	NAG	A	802	1	-	0/6/23/26	0/1/1/1
2	NAG	A	803	1	-	0/6/23/26	0/1/1/1
2	NAG	A	804	1	-	0/6/23/26	0/1/1/1
2	NAG	A	807	1	-	0/6/23/26	0/1/1/1
4	SY1	B	800	-	-	0/10/20/20	0/4/4/4
4	SY1	B	801	-	-	0/10/20/20	0/4/4/4
2	NAG	B	802	1	-	0/6/23/26	0/1/1/1
2	NAG	B	803	1	-	0/6/23/26	0/1/1/1
2	NAG	B	804	1	-	0/6/23/26	0/1/1/1
2	NAG	B	807	1	-	0/6/23/26	0/1/1/1
4	SY1	C	800	-	-	0/10/20/20	0/4/4/4
4	SY1	C	801	-	-	0/10/20/20	0/4/4/4
2	NAG	C	802	1	-	0/6/23/26	0/1/1/1
2	NAG	C	803	1	-	0/6/23/26	0/1/1/1
2	NAG	C	806	1	-	0/6/23/26	0/1/1/1
2	NAG	C	807	1	-	0/6/23/26	0/1/1/1
4	SY1	D	800	-	-	0/10/20/20	0/4/4/4
4	SY1	D	801	-	-	0/10/20/20	0/4/4/4
2	NAG	D	802	1	-	0/6/23/26	0/1/1/1
2	NAG	D	805	1	-	0/6/23/26	0/1/1/1

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	800	SY1	C25-C26	-10.55	1.28	1.44
4	B	801	SY1	C25-C26	-10.34	1.28	1.44
4	D	800	SY1	C25-C26	-10.31	1.29	1.44
4	A	800	SY1	C25-C26	-10.25	1.29	1.44
4	B	800	SY1	C25-C26	-10.07	1.29	1.44

The worst 5 of 61 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	801	SY1	C25-C26-N27	-10.84	159.03	177.82
4	D	801	SY1	C25-C26-N27	-7.98	164.00	177.82
4	A	801	SY1	C25-C26-N27	-5.49	168.30	177.82
4	A	800	SY1	C3-C2-N1	-5.12	100.45	110.71
4	A	800	SY1	C7-C2-N1	-5.10	99.27	111.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	800	SY1	4	0
4	A	801	SY1	2	0
4	B	800	SY1	6	0
4	C	800	SY1	6	0
4	D	800	SY1	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	724/731 (99%)	0.28	39 (5%)	29 34	28, 45, 76, 113	1 (0%)
1	B	728/731 (99%)	0.16	21 (2%)	55 61	32, 49, 74, 102	0
1	C	723/731 (98%)	0.26	48 (6%)	22 25	31, 49, 77, 94	0
1	D	722/731 (98%)	0.54	70 (9%)	10 11	35, 62, 91, 133	0
All	All	2897/2924 (99%)	0.31	178 (6%)	25 28	28, 51, 83, 133	1 (0%)

The worst 5 of 178 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	392	LYS	6.5
1	D	83	TYR	6.4
1	A	88	VAL	6.3
1	D	77	LEU	6.0
1	C	88	VAL	5.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	A	808	14/15	0.95	0.16	1.69	57,61,63,65	0
3	NAG	B	805	14/15	0.92	0.16	0.76	68,70,73,77	0
3	NAG	C	804	14/15	0.94	0.14	0.16	62,65,68,72	0
3	NAG	D	803	14/15	0.93	0.14	-0.77	67,69,70,72	0
3	NAG	C	805	14/15	0.84	0.31	-	74,77,80,80	0
3	NAG	B	806	14/15	0.86	0.26	-	80,82,83,83	0
3	NAG	A	806	14/15	0.79	0.22	-	101,102,103,103	0
3	NAG	A	805	14/15	0.52	0.19	-	97,99,101,101	0
3	NAG	A	809	14/15	0.91	0.15	-	67,68,69,69	0
3	NAG	D	804	14/15	0.86	0.21	-	74,75,76,76	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	A	807	14/15	0.90	0.39	6.47	60,64,66,66	0
2	NAG	C	807	14/15	0.88	0.36	5.74	70,74,75,76	0
4	SY1	C	801	27/27	0.90	0.17	1.25	62,63,64,64	0
4	SY1	D	801	27/27	0.94	0.20	0.75	56,58,62,65	0
4	SY1	A	800	27/27	0.97	0.23	0.65	36,41,42,43	0
4	SY1	A	801	27/27	0.94	0.15	0.14	47,50,53,53	0
4	SY1	B	801	27/27	0.95	0.15	0.09	48,49,51,52	0
4	SY1	D	800	27/27	0.96	0.20	-0.37	43,46,47,47	0
4	SY1	C	800	27/27	0.97	0.17	-0.46	33,37,41,41	0
4	SY1	B	800	27/27	0.96	0.19	-1.09	38,40,44,45	0
2	NAG	C	802	14/15	0.81	0.28	-	72,73,74,75	0
2	NAG	D	802	14/15	0.83	0.23	-	78,80,81,81	0
2	NAG	B	807	14/15	0.79	0.14	-	75,77,78,79	0
2	NAG	A	803	14/15	0.79	0.31	-	72,75,77,78	0
2	NAG	B	804	14/15	0.82	0.28	-	72,74,76,77	0
2	NAG	A	802	14/15	0.68	0.19	-	105,105,106,106	0
2	NAG	B	803	14/15	0.93	0.17	-	62,64,66,67	0
2	NAG	B	802	14/15	0.78	0.13	-	98,100,100,100	0
2	NAG	C	806	14/15	0.80	0.16	-	82,84,86,86	0
2	NAG	A	804	14/15	0.88	0.30	-	70,72,76,77	0
2	NAG	D	805	14/15	0.88	0.17	-	80,81,82,83	0
2	NAG	C	803	14/15	0.83	0.29	-	69,71,74,75	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.