



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:26 PM GMT

PDB ID : 4OY4
Title : calcium-free CaMPARI v0.2
Authors : Fosque, B.F.; Schreiter, E.R.
Deposited on : 2014-02-10
Resolution : 2.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	7858 (2.04-2.00)
Clashscore	102246	9060 (2.04-2.00)
Ramachandran outliers	100387	8952 (2.04-2.00)
Sidechain outliers	100360	8951 (2.04-2.00)
RSRZ outliers	91569	7873 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	413	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	504	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2870 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chimera protein of Calmodulin, GPF-like protein EosFP, and Myosin light chain kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	335	Total	C	N	O	S	2	3	0
			2744	1742	461	519	22			

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	MET	-	expression tag	UNP P62161
A	-7	HIS	-	expression tag	UNP P62161
A	-6	HIS	-	expression tag	UNP P62161
A	-5	HIS	-	expression tag	UNP P62161
A	-4	HIS	-	expression tag	UNP P62161
A	-3	HIS	-	expression tag	UNP P62161
A	-2	HIS	-	expression tag	UNP P62161
A	-1	GLY	-	expression tag	UNP P62161
A	0	SER	-	expression tag	UNP P62161
A	59	ASP	ASN	engineered mutation	UNP P62161
A	146	ALA	-	linker	UNP P62161
A	147	LYS	-	linker	UNP P62161
A	148	LEU	-	linker	UNP P62161
A	149	GLU	-	linker	UNP P62161
A	150	CYS	-	linker	UNP P62161
A	165	HIS	THR	engineered mutation	UNP Q5S6Z9
A	232	ARG	-	linker	UNP Q5S6Z9
A	233	ARG	-	linker	UNP Q5S6Z9
A	234	GLY	-	linker	UNP Q5S6Z9
A	235	GLY	-	linker	UNP Q5S6Z9
A	236	THR	-	linker	UNP Q5S6Z9
A	237	GLY	-	linker	UNP Q5S6Z9
A	238	GLY	-	linker	UNP Q5S6Z9
A	239	SER	-	linker	UNP Q5S6Z9
A	240	MET	-	linker	UNP Q5S6Z9
A	241	VAL	-	linker	UNP Q5S6Z9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	251	LYS	ASN	engineered mutation	UNP Q5S6Z9
A	274	TYR	PHE	engineered mutation	UNP Q5S6Z9
A	279	THR	SER	engineered mutation	UNP Q5S6Z9
A	303	CR8	HIS	chromophore	UNP Q5S6Z9
A	303	CR8	TYR	chromophore	UNP Q5S6Z9
A	303	CR8	GLY	chromophore	UNP Q5S6Z9
A	309	VAL	ALA	engineered mutation	UNP Q5S6Z9
A	310	LYS	GLU	engineered mutation	UNP Q5S6Z9
A	314	ASN	HIS	engineered mutation	UNP Q5S6Z9
A	333	MET	LEU	engineered mutation	UNP Q5S6Z9
A	342	TYR	ILE	engineered mutation	UNP Q5S6Z9
A	361	TYR	HIS	engineered mutation	UNP Q5S6Z9
A	363	THR	VAL	engineered mutation	UNP Q5S6Z9
A	380	MET	GLU	engineered mutation	UNP Q5S6Z9
A	383	TRP	-	linker	UNP Q5S6Z9
A	384	THR	-	linker	UNP Q5S6Z9
A	385	ARG	-	linker	UNP Q5S6Z9
A	386	SER	-	linker	UNP Q5S6Z9
A	387	SER	-	linker	UNP Q5S6Z9
A	392	ASN	GLN	engineered mutation	UNP Q6LDG3

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	104	Total	O	0	0
			104	104		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chimera protein of Calmodulin, GPF-like protein EosFP, and Myosin light chain kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	68.67Å 68.67Å 172.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	53.75 – 2.03 53.75 – 2.03	Depositor EDS
% Data completeness (in resolution range)	99.9 (53.75-2.03) 99.9 (53.75-2.03)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.33 (at 2.03Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.196 , 0.235 0.205 , 0.242	Depositor DCC
R_{free} test set	1388 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	26.5	Xtriage
Anisotropy	0.224	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 44.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	2 of 27630 reflections (0.007%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2870	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CR8, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.11	3/2788 (0.1%)	1.04	9/3744 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	214	LYS	CD-CE	-27.25	0.83	1.51
1	A	383	TRP	CB-CG	-5.23	1.40	1.50
1	A	184	TYR	CD2-CE2	5.18	1.47	1.39

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	214	LYS	CG-CD-CE	8.46	137.28	111.90
1	A	214	LYS	CD-CE-NZ	-7.31	94.88	111.70
1	A	306	ARG	NE-CZ-NH2	6.88	123.74	120.30
1	A	306	ARG	NE-CZ-NH1	-6.47	117.06	120.30
1	A	380	MET	CG-SD-CE	-6.14	90.37	100.20
1	A	313	ASP	CB-CG-OD2	-5.69	113.18	118.30
1	A	248	MET	CG-SD-CE	5.59	109.14	100.20
1	A	49	ASP	CB-CG-OD1	5.58	123.32	118.30
1	A	23	ASP	N-CA-CB	-5.45	100.80	110.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2744	0	2650	20	0
2	A	10	0	0	0	0
3	A	12	0	16	0	0
4	A	104	0	0	2	0
All	All	2870	0	2666	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:ARG:NH2	1:A:225:SER:O	2.28	0.66
1:A:253[A]:ARG:NH1	1:A:255:GLU:OE2	2.26	0.63
1:A:240:MET:HG3	1:A:274:TYR:CZ	2.38	0.58
1:A:81:GLU:O	1:A:85:ARG:HG3	2.08	0.53
1:A:84:ILE:HD12	1:A:145:THR:HG22	1.91	0.51
1:A:85:ARG:NH2	1:A:138:GLU:OE1	2.44	0.50
1:A:76:LYS:HA	1:A:76:LYS:HE2	1.94	0.50
1:A:327:TYR:CZ	1:A:347:ILE:HD12	2.50	0.46
1:A:27:THR:OG1	1:A:30:GLU:HG3	2.16	0.46
1:A:390:LYS:HD3	4:A:619:HOH:O	2.16	0.45
1:A:11:PHE:CZ	1:A:71:MET:HE3	2.51	0.45
1:A:250:ILE:HD13	4:A:637:HOH:O	2.15	0.45
1:A:240:MET:HA	1:A:274:TYR:CE1	2.52	0.45
1:A:375:LYS:HD2	1:A:375:LYS:N	2.30	0.44
1:A:79:ASP:HA	1:A:84:ILE:HD11	2.00	0.43
1:A:18:PHE:CD1	1:A:34:VAL:HG22	2.54	0.43
1:A:138:GLU:O	1:A:142:GLN:HG2	2.20	0.42
1:A:35[A]:MET:HE2	1:A:35[A]:MET:HB3	1.78	0.41
1:A:50:MET:HA	1:A:53:GLU:HG3	2.02	0.41
1:A:81:GLU:OE2	1:A:85:ARG:NH1	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	331/413 (80%)	325 (98%)	5 (2%)	1 (0%)	46	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	90	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/351 (84%)	284 (97%)	10 (3%)	44	41

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	54	VAL
1	A	189	LYS
1	A	210	LYS
1	A	214	LYS
1	A	306	ARG
1	A	375	LYS
1	A	385	ARG
1	A	387	SER
1	A	388	ARG
1	A	389	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CR8	A	303	1	20,27,28	2.26	6 (30%)	16,37,39	1.66	4 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CR8	A	303	1	-	0/8/25/26	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	303	CR8	C17-N13	-5.13	1.39	1.49
1	A	303	CR8	C4-C1	-4.54	1.35	1.45
1	A	303	CR8	C2-C1	-2.27	1.40	1.45
1	A	303	CR8	C8-C7	2.36	1.42	1.36
1	A	303	CR8	O19-C	2.96	1.39	1.19
1	A	303	CR8	O3-C1	4.06	1.37	1.24

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	303	CR8	C4-C5-C7	-3.30	119.07	122.00
1	A	303	CR8	C16-C14-N15	2.30	128.86	123.77
1	A	303	CR8	C-C17-N13	2.74	116.81	112.37
1	A	303	CR8	C23-N11-C10	2.99	110.42	105.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	501	-	4,4,4	0.51	0	6,6,6	0.41	0
2	SO4	A	502	-	4,4,4	0.45	0	6,6,6	0.57	0
3	GOL	A	503	-	5,5,5	0.82	0	5,5,5	2.16	2 (40%)
3	GOL	A	504	-	5,5,5	0.28	0	5,5,5	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	501	-	-	0/0/0/0	0/0/0/0
2	SO4	A	502	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	503	-	-	0/4/4/4	0/0/0/0
3	GOL	A	504	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	503	GOL	C3-C2-C1	-2.50	101.32	111.12
3	A	503	GOL	O1-C1-C2	3.71	128.17	110.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	334/413 (80%)	0.06	7 (2%) 67 68	16, 27, 60, 84	1 (0%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	56	ALA	4.1
1	A	387	SER	3.8
1	A	386	SER	3.2
1	A	90	VAL	2.4
1	A	385	ARG	2.3
1	A	57	ASP	2.3
1	A	388	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CR8	A	303	25/26	0.98	0.10	-	15,17,20,20	0

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GOL	A	504	6/6	0.92	0.14	3.30	43,44,44,44	0
2	SO4	A	501	5/5	0.90	0.14	-0.12	56,67,72,73	0
3	GOL	A	503	6/6	0.84	0.22	-	29,43,49,50	0
2	SO4	A	502	5/5	0.92	0.12	-	67,70,74,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.