



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:26 AM GMT

PDB ID : 3P03
Title : Crystal structure of BetP-G153D with choline bound
Authors : Perez, C.; Ressler, S.; Ziegler, Z.
Deposited on : 2010-09-27
Resolution : 3.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1005 (3.42-3.30)
Clashscore	102246	1076 (3.42-3.30)
Ramachandran outliers	100387	1059 (3.42-3.30)
Sidechain outliers	100360	1058 (3.42-3.30)
RSRZ outliers	91569	1010 (3.42-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	566	 8% 30% 55% 11%
1	B	566	 8% 28% 52% 16%
1	C	566	 4% 27% 57% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CHT	C	2486	-	-	X	X

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 11349 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glycine betaine transporter BetP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	504	Total	C	N	O	S	0	0	0
			3868	2542	641	669	16			
1	B	476	Total	C	N	O	S	0	0	0
			3612	2391	577	628	16			
1	C	508	Total	C	N	O	S	0	0	0
			3862	2545	627	674	16			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	44	ALA	GLU	ENGINEERED MUTATION	UNP P54582
A	45	ALA	GLU	ENGINEERED MUTATION	UNP P54582
A	46	ALA	GLU	ENGINEERED MUTATION	UNP P54582
A	153	ASP	GLY	ENGINEERED MUTATION	UNP P54582
B	44	ALA	GLU	ENGINEERED MUTATION	UNP P54582
B	45	ALA	GLU	ENGINEERED MUTATION	UNP P54582
B	46	ALA	GLU	ENGINEERED MUTATION	UNP P54582
B	153	ASP	GLY	ENGINEERED MUTATION	UNP P54582
C	44	ALA	GLU	ENGINEERED MUTATION	UNP P54582
C	45	ALA	GLU	ENGINEERED MUTATION	UNP P54582
C	46	ALA	GLU	ENGINEERED MUTATION	UNP P54582
C	153	ASP	GLY	ENGINEERED MUTATION	UNP P54582


- Molecule 2 is CHOLINE ION (three-letter code: CHT) (formula: C₅H₁₄NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	C	1	7	5	1	1	0	0



• Molecule 1: Glycine betaine transporter BetP



Y553	Q481	K353	G418
R554	H482	S354	I421
E555	G483	A355	V422
Q556	Q484	D356	F423
Q557	L485	G357	E424
R558	E486	T358	Q425
F559	L487	A359	N426
R560	N488		G427
A561		W362	E428
R562	V491	L363	S429
L563	T492	G364	I430
A564	A493	S365	W431
A565	A494	W366	G432
E566	W495	T367	D433
R567	G496	I368	G434
R568	V497	F369	A435
VAL	A498	Y370	A436
HIS		W371	E437
ASN	T502	A372	W373
GLU	G503	E438	Q439
HIS	L504	W374	L440
ARG	T505	I375	F441
LYS	L506	S376	G442
ARG	L507	W377	L443
GLU		S378	L444
LEU	L515	P379	
ALA		F380	L447
ALA	L518	V381	P448
LYS	Q519	W383	
ARG	N520	F384	Q451
ARG	V521	L385	I452
ARG	T522	A386	M453
GLU	I523	R387	G454
ARG	V524	I388	I455
LYS	A525	S389	I456
ALA	L526	R390	A457
SER	T527	G391	W458
GLY	P528	R392	I459
ALA	F529	S393	L460
GLY	L530	I394	L461
LYS		R395	G462
ARG	L534	E396	T463
ARG	G535	F397	F464
	L536	I398	F465
	H537	L399	I466
	F538	G400	T467
	A539	V401	S468
	L540		A469
	V541		D470
	W542		S471
	D543		A472
	L544		S473
	S545		T474
	N546		W475
	D547		W476
	W548		G477
	I549		T478
	T550		M479
	L551		S480
	E552		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	117.56Å 129.31Å 183.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.20 – 3.35 48.18 – 3.30	Depositor EDS
% Data completeness (in resolution range)	91.2 (46.20-3.35) 87.6 (48.18-3.30)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	15.91 (at 3.33Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.2_432)	Depositor
R, R_{free}	0.245 , 0.300 0.281 , 0.319	Depositor DCC
R_{free} test set	3705 reflections (11.05%)	DCC
Wilson B-factor (Å ²)	70.2	Xtriage
Anisotropy	0.012	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 105.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 37324 reflections	Xtriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	11349	wwPDB-VP
Average B, all atoms (Å ²)	107.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CHT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/3967	0.56	1/5397 (0.0%)
1	B	0.37	0/3706	0.59	0/5051
1	C	0.44	0/3960	0.64	0/5396
All	All	0.38	0/11633	0.59	1/15844 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	367	THR	C-N-CA	-5.06	109.06	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3868	0	3897	495	0
1	B	3612	0	3647	542	0
1	C	3862	0	3899	579	0
2	C	7	0	14	6	0
All	All	11349	0	11457	1576	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 69.

The worst 5 of 1576 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:170:VAL:CG1	1:C:171:PRO:HD2	1.50	1.41
1:C:226:GLY:HA2	1:C:227:GLU:CB	1.47	1.40
1:C:247:VAL:HG12	1:C:502:ILE:CD1	1.54	1.35
1:B:69:ILE:O	1:B:73:THR:HG23	1.33	1.29
1:B:208:THR:HG22	1:B:213:ARG:O	1.32	1.27

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	498/566 (88%)	426 (86%)	62 (12%)	10 (2%)	9	45
1	B	470/566 (83%)	403 (86%)	61 (13%)	6 (1%)	15	54
1	C	504/566 (89%)	435 (86%)	60 (12%)	9 (2%)	11	47
All	All	1472/1698 (87%)	1264 (86%)	183 (12%)	25 (2%)	11	48

5 of 25 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	377	TRP
1	A	403	LEU
1	B	231	GLU
1	B	432	GLY
1	C	231	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	396/440 (90%)	378 (96%)	18 (4%)	34	72
1	B	371/440 (84%)	356 (96%)	15 (4%)	38	74
1	C	396/440 (90%)	369 (93%)	27 (7%)	20	58
All	All	1163/1320 (88%)	1103 (95%)	60 (5%)	29	67

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	371	TRP
1	C	76	TRP
1	C	465	PHE
1	B	397	PHE
1	C	121	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	445	HIS
1	C	173	HIS
1	C	517	ASN
1	B	343	ASN
1	B	426	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CHT	C	2486	-	6,6,6	0.77	0	8,8,8	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CHT	C	2486	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2486	CHT	6	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	504/566 (89%)	0.40	45 (8%) 12 12	28, 107, 309, 534	0
1	B	476/566 (84%)	0.44	48 (10%) 9 9	14, 97, 292, 557	0
1	C	508/566 (89%)	0.00	21 (4%) 41 40	7, 62, 194, 353	0
All	All	1488/1698 (87%)	0.28	114 (7%) 16 17	7, 86, 284, 557	0

The worst 5 of 114 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	296	SER	15.6
1	B	296	SER	9.1
1	A	297	GLY	8.5
1	A	299	GLY	7.3
1	C	299	GLY	7.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	CHT	C	2486	7/7	0.69	0.46	6.19	85,89,97,97	0

6.5 Other polymers [i](#)

There are no such residues in this entry.