



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:30 PM GMT

PDB ID : 4P4F  
Title : X-ray structure of human glutamate carboxypeptidase II (GCP II) in complex with a phosphoramidate inhibitor NC-2-40  
Authors : Barinka, C.  
Deposited on : 2014-03-12  
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*


The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	707	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	802	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	803	-	-	-	X
2	NAG	A	809	-	-	-	X
2	NAG	A	812	-	-	-	X

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 6648 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate carboxypeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	694	Total	C	N	O	S	0	62	0
			5834	3741	975	1095	23			

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is BETA-D-MANNOSE (three-letter code: BMA) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Zn	0	0
			2	2		


- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Cl	0	0
			1	1		

- Molecule 8 is N-[(S)-(biphenyl-4-ylmethoxy)(hydroxy)phosphoryl]-L-glutamic acid (three-letter code: 2GB) (formula: C<sub>18</sub>H<sub>20</sub>NO<sub>7</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	A	1	Total	C	N	O	P	0	0
			27	18	1	7	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	582	Total O 582 582	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Glutamate carboxypeptidase 2





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	101.77Å 130.61Å 159.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.86 20.01 – 1.86	Depositor EDS
% Data completeness (in resolution range)	98.2 (20.00-1.86) 98.2 (20.01-1.86)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.67 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.152 , 0.176 0.152 , 0.174	Depositor DCC
$R_{free}$ test set	1744 reflections (2.05%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.9	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 56.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 86819 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6648	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BMA, NAG, CL, CA, 2GB, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.92	1/6156 (0.0%)	0.75	2/8332 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	433	GLU	CB-CG	5.16	1.61	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	174[A]	LEU	CA-CB-CG	5.01	126.83	115.30
1	A	174[B]	LEU	CA-CB-CG	5.01	126.83	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5834	0	5685	62	0
2	A	168	0	149	1	0
3	A	22	0	19	0	0
4	A	11	0	10	0	0
5	A	2	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	1	0	0	0	0
7	A	1	0	0	0	0
8	A	27	0	17	1	0
9	A	582	0	0	13	1
All	All	6648	0	5880	63	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:505[B]:GLU:OE1	1:A:606[B]:LYS:HE2	1.41	1.21
1:A:505[B]:GLU:CD	1:A:606[B]:LYS:HE2	1.62	1.20
1:A:660[A]:VAL:O	1:A:664[A]:MET:HG2	1.38	1.19
1:A:597[B]:CYS:SG	1:A:671:LEU:HD22	1.99	1.02
1:A:597[B]:CYS:SG	1:A:671:LEU:CD2	2.49	1.00
1:A:505[B]:GLU:OE1	1:A:606[B]:LYS:CE	2.12	0.97
1:A:412:PRO:HA	1:A:589[B]:ASN:HD21	1.29	0.97
1:A:688[B]:ARG:NH1	9:A:1062:HOH:O	2.15	0.80
1:A:656[B]:SER:O	9:A:901:HOH:O	2.03	0.76
1:A:516:GLY:O	1:A:526[B]:GLN:NE2	2.21	0.73
1:A:505[B]:GLU:OE1	1:A:606[B]:LYS:CD	2.37	0.72
1:A:688[B]:ARG:HG3	1:A:688[B]:ARG:NH2	2.05	0.69
1:A:80:ILE:HD12	1:A:88[B]:GLN:HG2	1.76	0.67
1:A:656[B]:SER:OG	9:A:901:HOH:O	2.14	0.66
1:A:597[B]:CYS:SG	1:A:671:LEU:HD23	2.37	0.65
1:A:505[B]:GLU:OE2	1:A:606[B]:LYS:HE2	1.96	0.65
1:A:437[A]:GLU:OE1	9:A:903:HOH:O	2.14	0.64
2:A:802:NAG:H81	9:A:1215:HOH:O	1.97	0.64
1:A:539:LYS:HE3	9:A:1476:HOH:O	1.97	0.64
1:A:641:GLU:HG3	9:A:1318:HOH:O	1.97	0.63
1:A:688[B]:ARG:HG3	1:A:688[B]:ARG:HH21	1.62	0.63
1:A:688[B]:ARG:HH21	1:A:688[B]:ARG:CG	2.13	0.61
1:A:505[B]:GLU:CD	1:A:606[B]:LYS:CE	2.55	0.60
1:A:400:ARG:O	1:A:404[B]:THR:HG23	2.02	0.59
1:A:506:PHE:HB2	1:A:509:MET:HG3	1.88	0.55
1:A:174[B]:LEU:HD11	1:A:342:VAL:HG21	1.90	0.54
1:A:539:LYS:HG3	1:A:540:ASN:N	2.24	0.53
1:A:526[B]:GLN:HG2	1:A:694:PRO:HD3	1.89	0.53
1:A:685:PRO:O	1:A:688[B]:ARG:HD3	2.08	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:480:GLU:HG3	9:A:1456:HOH:O	2.09	0.53
1:A:198:GLY:O	1:A:223:LYS:HE2	2.08	0.53
1:A:505[B]:GLU:OE1	1:A:606[B]:LYS:HD2	2.09	0.52
1:A:659[B]:ILE:O	1:A:663[B]:MET:HG3	2.11	0.49
1:A:188:LEU:HD21	1:A:329[A]:VAL:HG11	1.94	0.49
1:A:412:PRO:HA	1:A:589[B]:ASN:ND2	2.12	0.48
1:A:169:MET:HA	1:A:344:MET:O	2.15	0.47
1:A:662[B]:ARG:NE	9:A:905:HOH:O	2.30	0.46
1:A:188:LEU:CD2	1:A:329[A]:VAL:HG11	2.46	0.46
1:A:413:ARG:H	1:A:589[B]:ASN:ND2	2.13	0.46
1:A:540:ASN:HD21	1:A:542:GLU:HB2	1.80	0.46
1:A:362:LEU:CD1	1:A:406:LYS:HD2	2.46	0.46
1:A:188:LEU:HD21	1:A:329[A]:VAL:CG1	2.47	0.45
1:A:548:GLY:HA2	8:A:820:2GB:CAQ	2.46	0.45
1:A:66[B]:LYS:HB3	1:A:68[B]:GLU:HG2	1.97	0.45
1:A:58[B]:MET:SD	1:A:586:GLU:HG2	2.57	0.45
1:A:239[A]:VAL:HG22	1:A:247:ASN:ND2	2.32	0.45
1:A:540:ASN:C	1:A:540:ASN:HD22	2.20	0.45
1:A:263:GLY:HA2	1:A:522:GLU:OE1	2.17	0.44
1:A:514[B]:LYS:HG2	9:A:1110:HOH:O	2.17	0.44
1:A:88[B]:GLN:CD	9:A:1295:HOH:O	2.56	0.44
1:A:469:LEU:O	1:A:595:PHE:HA	2.19	0.42
1:A:278:ALA:HB3	1:A:280[A]:ARG:NH1	2.34	0.42
1:A:367:GLU:OE1	1:A:662[A]:ARG:NH1	2.52	0.42
1:A:656[B]:SER:CB	9:A:901:HOH:O	2.68	0.41
1:A:506:PHE:CB	1:A:509:MET:HG3	2.50	0.41
1:A:225:VAL:O	1:A:294:VAL:HA	2.21	0.41
1:A:660[A]:VAL:O	1:A:664[A]:MET:CG	2.33	0.41
1:A:387:ASP:HA	1:A:388:PRO:HA	1.93	0.41
1:A:469:LEU:HD21	1:A:671:LEU:HD23	2.02	0.40
1:A:645[A]:LYS:HD3	9:A:1346:HOH:O	2.21	0.40
1:A:659[B]:ILE:HA	1:A:659[B]:ILE:HD13	1.79	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:908:HOH:O	9:A:952:HOH:O[2_565]	1.80	0.40

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	749/707 (106%)	730 (98%)	16 (2%)	3 (0%)	39 22

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	335	GLY
1	A	540	ASN
1	A	382	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	651/603 (108%)	640 (98%)	11 (2%)	68 54

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58[A]	MET
1	A	58[B]	MET
1	A	102[A]	GLU
1	A	102[B]	GLU
1	A	519	ASN
1	A	536	ARG
1	A	537	TYR
1	A	539	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	540	ASN
1	A	600	TYR
1	A	673	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	303	GLN
1	A	540	ASN
1	A	618	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

13 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	801	1,2	14,14,15	0.50	0	15,19,21	1.23	1 (6%)
2	NAG	A	802	2	14,14,15	0.58	0	15,19,21	1.18	2 (13%)
2	NAG	A	803	1,2	14,14,15	0.75	0	15,19,21	1.35	0
2	NAG	A	804	2	14,14,15	0.43	0	15,19,21	1.23	1 (6%)
2	NAG	A	805	1,2	14,14,15	0.66	0	15,19,21	1.27	2 (13%)
2	NAG	A	806	2	14,14,15	0.50	0	15,19,21	1.15	1 (6%)
2	NAG	A	807	1	14,14,15	0.41	0	15,19,21	2.20	2 (13%)
2	NAG	A	808	1	14,14,15	0.79	0	15,19,21	1.33	2 (13%)
2	NAG	A	809	1,2	14,14,15	0.86	1 (7%)	15,19,21	1.02	1 (6%)
2	NAG	A	810	3,2	14,14,15	0.67	0	15,19,21	1.12	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	812	1,2	14,14,15	0.65	0	15,19,21	1.49	2 (13%)
2	NAG	A	813	3,2	14,14,15	0.59	0	15,19,21	1.57	3 (20%)
4	MAN	A	815	3	11,11,12	0.57	0	14,15,17	1.16	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	801	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	802	2	-	0/6/23/26	0/1/1/1
2	NAG	A	803	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	804	2	-	0/6/23/26	0/1/1/1
2	NAG	A	805	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	806	2	-	0/6/23/26	0/1/1/1
2	NAG	A	807	1	-	0/6/23/26	0/1/1/1
2	NAG	A	808	1	-	0/6/23/26	0/1/1/1
2	NAG	A	809	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	810	3,2	-	0/6/23/26	0/1/1/1
2	NAG	A	812	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	813	3,2	-	0/6/23/26	0/1/1/1
4	MAN	A	815	3	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	809	NAG	O7-C7	2.51	1.29	1.23

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	813	NAG	C3-C4-C5	-3.43	104.22	110.20
2	A	812	NAG	C2-N2-C7	-3.14	119.01	123.04
2	A	808	NAG	C2-N2-C7	-2.61	119.69	123.04
2	A	805	NAG	O7-C7-C8	-2.08	118.24	122.06
2	A	807	NAG	C6-C5-C4	-2.08	107.89	113.02
2	A	813	NAG	C4-C3-C2	-2.06	108.03	111.23
2	A	802	NAG	C2-N2-C7	2.08	125.71	123.04
2	A	805	NAG	C1-O5-C5	2.30	115.16	112.25
2	A	801	NAG	C1-O5-C5	2.40	115.29	112.25

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	806	NAG	O5-C5-C6	2.44	112.63	107.35
2	A	810	NAG	C8-C7-N2	2.47	120.83	116.11
2	A	808	NAG	O5-C5-C6	2.47	112.70	107.35
2	A	813	NAG	C8-C7-N2	2.50	120.89	116.11
2	A	810	NAG	C2-N2-C7	2.52	126.28	123.04
2	A	802	NAG	C1-O5-C5	2.79	115.79	112.25
2	A	809	NAG	C2-N2-C7	2.83	126.68	123.04
4	A	815	MAN	O5-C5-C6	3.06	113.97	107.35
2	A	812	NAG	C1-O5-C5	3.43	116.60	112.25
2	A	804	NAG	C1-O5-C5	3.61	116.83	112.25
2	A	807	NAG	C1-O5-C5	7.74	122.07	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	802	NAG	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 4 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	801	1,2	14,14,15	0.50	0	15,19,21	1.23	1 (6%)
2	NAG	A	802	2	14,14,15	0.58	0	15,19,21	1.18	2 (13%)



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	803	1,2	14,14,15	0.75	0	15,19,21	1.35	0
2	NAG	A	804	2	14,14,15	0.43	0	15,19,21	1.23	1 (6%)
2	NAG	A	805	1,2	14,14,15	0.66	0	15,19,21	1.27	2 (13%)
2	NAG	A	806	2	14,14,15	0.50	0	15,19,21	1.15	1 (6%)
2	NAG	A	807	1	14,14,15	0.41	0	15,19,21	2.20	2 (13%)
2	NAG	A	808	1	14,14,15	0.79	0	15,19,21	1.33	2 (13%)
2	NAG	A	809	1,2	14,14,15	0.86	1 (7%)	15,19,21	1.02	1 (6%)
2	NAG	A	810	3,2	14,14,15	0.67	0	15,19,21	1.12	2 (13%)
3	BMA	A	811	2	11,11,12	0.85	0	14,15,17	1.09	0
2	NAG	A	812	1,2	14,14,15	0.65	0	15,19,21	1.49	2 (13%)
2	NAG	A	813	3,2	14,14,15	0.59	0	15,19,21	1.57	3 (20%)
3	BMA	A	814	2,4	11,11,12	0.55	0	14,15,17	0.97	1 (7%)
4	MAN	A	815	3	11,11,12	0.57	0	14,15,17	1.16	1 (7%)
8	2GB	A	820	5	21,28,28	3.22	7 (33%)	27,38,38	2.06	7 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	801	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	802	2	-	0/6/23/26	0/1/1/1
2	NAG	A	803	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	804	2	-	0/6/23/26	0/1/1/1
2	NAG	A	805	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	806	2	-	0/6/23/26	0/1/1/1
2	NAG	A	807	1	-	0/6/23/26	0/1/1/1
2	NAG	A	808	1	-	0/6/23/26	0/1/1/1
2	NAG	A	809	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	810	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	811	2	-	0/2/19/22	0/1/1/1
2	NAG	A	812	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	813	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	814	2,4	-	0/2/19/22	0/1/1/1
4	MAN	A	815	3	-	0/2/19/22	0/1/1/1
8	2GB	A	820	5	-	0/17/24/24	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	820	2GB	CAN-CAI	-2.47	1.42	1.49
2	A	809	NAG	O7-C7	2.51	1.29	1.23
8	A	820	2GB	CAL-CAK	2.85	1.43	1.38
8	A	820	2GB	OAT-CAB	4.39	1.63	1.44
8	A	820	2GB	CAB-CAH	5.22	1.62	1.50
8	A	820	2GB	PBA-OAU	5.75	1.52	1.46
8	A	820	2GB	CB-CA	6.56	1.62	1.53
8	A	820	2GB	PBA-N	8.26	1.70	1.61

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	820	2GB	OAU-PBA-N	-3.67	106.51	113.08
2	A	813	NAG	C3-C4-C5	-3.43	104.22	110.20
2	A	812	NAG	C2-N2-C7	-3.14	119.01	123.04
8	A	820	2GB	CAM-CAI-CAN	-2.74	116.48	121.39
2	A	808	NAG	C2-N2-C7	-2.61	119.69	123.04
8	A	820	2GB	CG-CB-CA	-2.21	108.50	112.99
2	A	805	NAG	O7-C7-C8	-2.08	118.24	122.06
2	A	807	NAG	C6-C5-C4	-2.08	107.89	113.02
2	A	813	NAG	C4-C3-C2	-2.06	108.03	111.23
3	A	814	BMA	C1-O5-C5	2.07	114.88	112.25
2	A	802	NAG	C2-N2-C7	2.08	125.71	123.04
2	A	805	NAG	C1-O5-C5	2.30	115.16	112.25
8	A	820	2GB	CAB-CAH-CAK	2.35	126.31	120.66
2	A	801	NAG	C1-O5-C5	2.40	115.29	112.25
2	A	806	NAG	O5-C5-C6	2.44	112.63	107.35
2	A	810	NAG	C8-C7-N2	2.47	120.83	116.11
2	A	808	NAG	O5-C5-C6	2.47	112.70	107.35
2	A	813	NAG	C8-C7-N2	2.50	120.89	116.11
2	A	810	NAG	C2-N2-C7	2.52	126.28	123.04
2	A	802	NAG	C1-O5-C5	2.79	115.79	112.25
2	A	809	NAG	C2-N2-C7	2.83	126.68	123.04
4	A	815	MAN	O5-C5-C6	3.06	113.97	107.35
8	A	820	2GB	CAL-CAI-CAN	3.20	127.12	121.39
2	A	812	NAG	C1-O5-C5	3.43	116.60	112.25
2	A	804	NAG	C1-O5-C5	3.61	116.83	112.25
8	A	820	2GB	OAV-PBA-OAT	4.09	118.53	106.60
8	A	820	2GB	OAT-CAB-CAH	5.82	119.03	109.17
2	A	807	NAG	C1-O5-C5	7.74	122.07	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	802	NAG	1	0
8	A	820	2GB	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	694/707 (98%)	-0.33	21 (3%) 54 51	14, 27, 49, 71	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	153	ASN	5.0
1	A	541	TRP	4.9
1	A	55	LYS	3.8
1	A	152	GLU	3.6
1	A	656[A]	SER	3.5
1	A	237	PRO	3.2
1	A	335	GLY	3.1
1	A	336	ASN	3.1
1	A	719	VAL	3.0
1	A	542	GLU	3.0
1	A	155[A]	SER	3.0
1	A	134	ASP	2.9
1	A	506	PHE	2.8
1	A	652	ASP	2.7
1	A	174[A]	LEU	2.7
1	A	507	SER	2.6
1	A	408	GLU	2.6
1	A	487	GLY	2.4
1	A	56	HIS	2.4
1	A	133	GLU	2.3
1	A	543	THR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	A	802	14/15	0.71	0.31	8.81	46,54,60,61	0
2	NAG	A	809	14/15	0.94	0.12	4.16	31,34,40,40	0
2	NAG	A	812	14/15	0.93	0.10	3.41	21,29,41,46	0
2	NAG	A	803	14/15	0.66	0.25	2.67	49,55,65,68	0
4	MAN	A	815	11/12	0.92	0.12	0.29	52,55,58,59	0
2	NAG	A	808	14/15	0.92	0.13	0.20	35,45,54,57	0
2	NAG	A	801	14/15	0.96	0.12	-	38,44,49,50	0
2	NAG	A	804	14/15	0.85	0.45	-	73,76,80,80	0
2	NAG	A	813	14/15	0.92	0.27	-	35,42,48,52	0
2	NAG	A	810	14/15	0.87	0.17	-	43,50,54,60	0
2	NAG	A	806	14/15	0.77	0.32	-	57,67,72,72	0
2	NAG	A	805	14/15	0.92	0.15	-	45,50,53,59	0
2	NAG	A	807	14/15	0.85	0.23	-	70,76,78,80	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	A	802	14/15	0.71	0.31	8.81	46,54,60,61	0
2	NAG	A	809	14/15	0.94	0.12	4.16	31,34,40,40	0
2	NAG	A	812	14/15	0.93	0.10	3.41	21,29,41,46	0
2	NAG	A	803	14/15	0.66	0.25	2.67	49,55,65,68	0
4	MAN	A	815	11/12	0.92	0.12	0.29	52,55,58,59	0
2	NAG	A	808	14/15	0.92	0.13	0.20	35,45,54,57	0
8	2GB	A	820	27/27	0.98	0.08	0.00	15,21,37,38	0
6	CA	A	818	1/1	1.00	0.06	-0.12	17,17,17,17	0
7	CL	A	819	1/1	0.99	0.04	-1.90	22,22,22,22	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	ZN	A	817	1/1	1.00	0.03	-2.15	19,19,19,19	0
5	ZN	A	816	1/1	1.00	0.03	-2.84	20,20,20,20	0
2	NAG	A	810	14/15	0.87	0.17	-	43,50,54,60	0
2	NAG	A	804	14/15	0.85	0.45	-	73,76,80,80	0
3	BMA	A	811	11/12	0.74	0.31	-	66,70,76,79	0
2	NAG	A	813	14/15	0.92	0.27	-	35,42,48,52	0
2	NAG	A	807	14/15	0.85	0.23	-	70,76,78,80	0
3	BMA	A	814	11/12	0.86	0.15	-	45,48,51,52	0
2	NAG	A	805	14/15	0.92	0.15	-	45,50,53,59	0
2	NAG	A	801	14/15	0.96	0.12	-	38,44,49,50	0
2	NAG	A	806	14/15	0.77	0.32	-	57,67,72,72	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.