



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:32 PM GMT

PDB ID : 1PHN
Title : STRUCTURE OF PHYCOCYANIN FROM CYANIDIUM CALDARIUM AT
1.65Å RESOLUTION
Authors : Stec, B.; Troxler, R.F.; Teeter, M.M.
Deposited on : 1995-06-21
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1323 (1.66-1.66)
Ramachandran outliers	100387	1295 (1.66-1.66)
Sidechain outliers	100360	1295 (1.66-1.66)
RSRZ outliers	91569	1227 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	162	
2	B	172	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PEB	B	182	X	-	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3208 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHYCOCYANIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	0	0	0
			1231	771	212	242	6			


- Molecule 2 is a protein called PHYCOCYANIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	172	Total	C	N	O	S	0	0	0
			1276	787	226	254	9			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	72	MEN	THR	CONFLICT	UNP P00311

- Molecule 3 is PHYCOCYANOBILIN (three-letter code: CYC) (formula: $C_{33}H_{40}N_4O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			43	33	4	6		
3	B	1	Total	C	N	O	0	0
			43	33	4	6		

- Molecule 4 is PHYCOERYTHROBILIN (three-letter code: PEB) (formula: C₃₃H₄₀N₄O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			43	33	4	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	280	Total O 280 280	0	0
5	B	292	Total O 292 292	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PHYCOCYANIN

Chain A:  90% 10%



• Molecule 2: PHYCOCYANIN

Chain B:  3% 88% 10%



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	106.42Å 106.42Å 176.18Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 1.65 27.22 – 1.65	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-1.65) 97.3 (27.22-1.65)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 1.65Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.183 , 0.272 0.189 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	7.8	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.24 , 73.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Outliers	0 of 78064 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3208	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CYC, MEN, PEB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/1254	0.65	0/1702
2	B	0.46	0/1278	0.71	1/1726 (0.1%)
All	All	0.48	0/2532	0.68	1/3428 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	79	ARG	NE-CZ-NH2	7.40	124.00	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1231	0	1200	11	0
2	B	1276	0	1280	23	0
3	A	43	0	36	2	0
3	B	43	0	37	1	0
4	B	43	0	35	9	0
5	A	280	0	0	3	0
5	B	292	0	0	2	0
All	All	3208	0	2588	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:182:PEB:CMC	4:B:182:PEB:HMD2	1.73	1.19
4:B:182:PEB:HMC1	4:B:182:PEB:HMD2	1.28	1.08
2:B:90:ILE:HG21	4:B:182:PEB:HBD1	1.56	0.88
2:B:90:ILE:CG2	4:B:182:PEB:HBD1	2.09	0.83
4:B:182:PEB:HMC3	4:B:182:PEB:HMD2	1.63	0.80
2:B:106:ILE:HG12	2:B:110:ARG:NH1	1.98	0.79
2:B:57:ARG:HD3	5:B:328:HOH:O	1.86	0.75
2:B:97:TYR:HB2	2:B:106:ILE:HD12	1.69	0.73
4:B:182:PEB:HMC1	4:B:182:PEB:CMD	2.13	0.69
4:B:182:PEB:CMC	4:B:182:PEB:CMD	2.64	0.65
1:A:128:TRP:CD2	3:A:184:CYC:HMC3	2.38	0.59
2:B:110:ARG:O	2:B:113:ASN:HB2	2.02	0.59
2:B:106:ILE:HG12	2:B:110:ARG:HH12	1.67	0.58
2:B:108:ASP:O	2:B:112:LEU:HB2	2.02	0.58
1:A:59:VAL:HG11	3:A:184:CYC:HMC1	1.88	0.55
1:A:73:GLN:HB3	5:A:277:HOH:O	2.05	0.54
2:B:97:TYR:HD2	2:B:106:ILE:HD11	1.71	0.54
2:B:94:TYR:HE2	2:B:110:ARG:NH1	2.05	0.54
2:B:7:LYS:O	2:B:11:GLN:HG3	2.09	0.52
1:A:1:MET:HA	1:A:109:GLU:OE2	2.11	0.51
2:B:115:LEU:O	2:B:115:LEU:HD12	2.12	0.49
4:B:182:PEB:HNA	4:B:182:PEB:HMB2	1.79	0.48
1:A:9:ILE:HB	2:B:1:MET:HE3	1.95	0.47
2:B:111:CYS:SG	2:B:172:ALA:CB	3.03	0.47
2:B:106:ILE:O	2:B:110:ARG:HD3	2.15	0.47
2:B:122:LEU:HD11	4:B:182:PEB:HAC2	1.98	0.44
2:B:94:TYR:HD2	2:B:106:ILE:HD13	1.84	0.43
2:B:7:LYS:HD3	5:B:248:HOH:O	2.19	0.43
2:B:106:ILE:HD12	2:B:106:ILE:HG21	1.71	0.42
1:A:3:THR:O	1:A:7:GLU:HG3	2.20	0.42
2:B:97:TYR:CB	2:B:106:ILE:HD12	2.44	0.42
1:A:43:SER:CB	1:A:144:LEU:HD21	2.49	0.42
1:A:38:LEU:HD21	5:A:421:HOH:O	2.20	0.42
1:A:43:SER:HB3	1:A:144:LEU:HD21	2.02	0.41
1:A:116:GLU:HG3	5:A:271:HOH:O	2.20	0.41
1:A:27:VAL:HG21	2:B:100:ILE:HD11	2.01	0.41
2:B:2:LEU:HD13	2:B:7:LYS:HA	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:36:LYS:HE2	3:B:183:CYC:HMD3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/162 (99%)	158 (99%)	2 (1%)	0	100	100
2	B	169/172 (98%)	165 (98%)	4 (2%)	0	100	100
All	All	329/334 (98%)	323 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/124 (100%)	121 (98%)	3 (2%)	57	28
2	B	132/132 (100%)	128 (97%)	4 (3%)	48	18
All	All	256/256 (100%)	249 (97%)	7 (3%)	52	23

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	GLN
1	A	140	HIS
1	A	150	ASN
2	B	30	MET
2	B	106	ILE
2	B	113	ASN
2	B	115	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	GLN
1	A	150	ASN
2	B	113	ASN
2	B	145	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MEN	B	72	2	7,8,9	0.83	0	5,9,11	1.15	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MEN	B	72	2	-	0/6/8/10	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	72	MEN	O-C-CA	-2.49	119.01	125.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CYC	A	184	1	35,46,46	1.53	5 (14%)	47,67,67	3.22	17 (36%)
4	PEB	B	182	2	36,46,46	3.03	13 (36%)	38,67,67	3.19	16 (42%)
3	CYC	B	183	2	35,46,46	2.66	6 (17%)	47,67,67	2.27	10 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CYC	A	184	1	-	2/21/74/74	0/4/4/4
4	PEB	B	182	2	1/1/14/19	3/19/74/74	0/4/4/4
3	CYC	B	183	2	-	2/21/74/74	0/4/4/4

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	183	CYC	C2C-C1C	-11.75	1.41	1.52
4	B	182	PEB	C2A-C1A	-8.60	1.44	1.52
4	B	182	PEB	CMA-C2A	-7.41	1.36	1.53
4	B	182	PEB	CHC-C1D	-6.19	1.38	1.54
4	B	182	PEB	C1D-ND	-4.49	1.38	1.46
3	B	183	CYC	CMC-C2C	-4.14	1.43	1.53
4	B	182	PEB	C4B-C3B	-3.18	1.40	1.45
3	B	183	CYC	C1C-NC	-3.08	1.33	1.37
3	A	184	CYC	C4B-C3B	-2.84	1.42	1.48
3	A	184	CYC	C1C-NC	-2.53	1.34	1.37
3	B	183	CYC	C2C-C3C	-2.42	1.47	1.54
3	B	183	CYC	C4B-C3B	-2.41	1.43	1.48
4	B	182	PEB	C2A-C3A	-2.17	1.48	1.54
3	A	184	CYC	CAD-C3D	2.26	1.55	1.52
3	A	184	CYC	OC-C1C	2.29	1.27	1.23
4	B	182	PEB	CAD-C3D	2.34	1.54	1.47
4	B	182	PEB	C4A-NA	2.40	1.42	1.37
4	B	182	PEB	C3C-C4C	2.93	1.46	1.42
4	B	182	PEB	CAA-C3A	3.69	1.61	1.54
4	B	182	PEB	C1A-NA	4.00	1.42	1.37
4	B	182	PEB	CBD-CAD	4.53	1.52	1.30
3	A	184	CYC	CHA-C1A	5.72	1.40	1.35
4	B	182	PEB	CHB-C4B	6.27	1.40	1.35
3	B	183	CYC	CHA-C1A	7.33	1.41	1.35

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	184	CYC	OC-C1C-C2C	-8.06	119.74	126.25
4	B	182	PEB	OA-C1A-C2A	-7.67	120.05	126.25
3	A	184	CYC	C4B-C3B-C2B	-6.51	104.33	108.05
3	B	183	CYC	OB-C4B-C3B	-5.33	121.70	128.09
3	A	184	CYC	C3C-C2C-C1C	-4.38	99.74	103.41
3	A	184	CYC	OB-C4B-C3B	-4.33	122.89	128.09
3	B	183	CYC	C1B-NB-C4B	-4.18	104.62	110.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	184	CYC	CHA-C1A-NA	-4.11	121.32	128.67
4	B	182	PEB	C2A-C1A-NA	-4.09	104.39	108.30
3	A	184	CYC	C1B-NB-C4B	-4.05	104.80	110.73
4	B	182	PEB	OD-C4D-C3D	-3.45	121.21	129.82
3	A	184	CYC	CAD-C3D-C4D	-2.72	124.05	127.01
3	B	183	CYC	C4B-C3B-C2B	-2.69	106.51	108.05
4	B	182	PEB	CAC-C2C-C3C	-2.51	121.85	129.00
4	B	182	PEB	CBD-CAD-C3D	-2.35	115.06	127.01
4	B	182	PEB	CHA-C1B-NB	-2.27	120.60	124.91
3	A	184	CYC	CHB-C4A-NA	-2.06	120.98	124.91
3	B	183	CYC	CAC-C3C-C4C	2.03	117.89	112.67
3	A	184	CYC	CHB-C4A-C3A	2.03	129.84	124.88
4	B	182	PEB	CHA-C1B-C2B	2.06	129.91	124.88
3	B	183	CYC	C2B-C1B-NB	2.14	110.11	107.00
3	B	183	CYC	CAC-C3C-C2C	2.15	119.55	114.13
4	B	182	PEB	CAA-C3A-C4A	2.24	118.43	112.67
4	B	182	PEB	CAA-C3A-C2A	2.28	119.85	114.13
3	B	183	CYC	CAB-C3B-C4B	2.31	123.53	121.51
4	B	182	PEB	CMA-C2A-C1A	2.80	118.32	112.43
4	B	182	PEB	CBC-CAC-C2C	2.91	117.74	112.53
3	A	184	CYC	CMA-C3A-C4A	3.03	130.00	125.06
4	B	182	PEB	CMB-C2B-C1B	3.05	130.03	125.06
3	A	184	CYC	CBD-CAD-C3D	3.20	118.27	112.53
3	A	184	CYC	C2C-C3C-C4C	3.31	107.11	101.50
4	B	182	PEB	CAC-C2C-C1C	3.61	130.92	127.01
3	A	184	CYC	CAC-C3C-C2C	3.91	123.95	114.13
3	B	183	CYC	CMA-C3A-C4A	4.06	131.67	125.06
4	B	182	PEB	CMC-C3C-C4C	4.26	131.44	127.14
3	A	184	CYC	CMC-C2C-C1C	4.92	122.76	112.43
3	B	183	CYC	C3B-C4B-NB	5.91	112.11	106.74
3	A	184	CYC	C3B-C4B-NB	7.49	113.55	106.74
3	A	184	CYC	C2C-C1C-NC	7.55	115.51	108.30
4	B	182	PEB	OA-C1A-NA	7.60	134.02	124.83
3	A	184	CYC	CAB-C3B-C4B	9.14	129.50	121.51
3	B	183	CYC	C3C-C2C-C1C	9.70	111.54	103.41
4	B	182	PEB	C3A-C2A-C1A	11.58	113.12	103.41

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	B	182	PEB	C1D

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	182	PEB	C4C-CHC-C1D-ND
3	B	183	CYC	C1B-CHB-C4A-C3A
4	B	182	PEB	C4A-CHA-C1B-C2B
3	A	184	CYC	C1B-CHB-C4A-C3A
3	B	183	CYC	C1B-CHB-C4A-NA
3	A	184	CYC	C1B-CHB-C4A-NA
4	B	182	PEB	C4A-CHA-C1B-NB

There are no ring outliers.

3 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	184	CYC	2	0
4	B	182	PEB	9	0
3	B	183	CYC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	162/162 (100%)	-0.70	0 100 100	2, 7, 18, 31	0
2	B	171/172 (99%)	-0.28	6 (3%) 48 48	3, 12, 43, 60	0
All	All	333/334 (99%)	-0.48	6 (1%) 71 75	2, 9, 29, 60	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	112	LEU	5.2
2	B	113	ASN	4.8
2	B	111	CYS	3.6
2	B	110	ARG	3.5
2	B	114	GLY	2.6
2	B	109	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	MEN	B	72	9/10	0.94	0.08	-	6,8,11,16	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	PEB	B	182	43/43	0.91	0.11	0.82	3,22,32,34	0
3	CYC	A	184	43/43	0.97	0.06	0.25	2,3,5,6	0
3	CYC	B	183	43/43	0.95	0.07	-0.13	4,8,13,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.