



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:40 PM GMT

PDB ID : 4PLP  
Title : Crystal Structure of the Homospermidine Synthase (HSS) from *Blastochloris viridis* in Complex with NAD  
Authors : Krossa, S.  
Deposited on : 2014-05-19  
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	479	 3% 93% 6%
1	B	479	 4% 93% 5%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 16118 atoms, of which 7418 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Homospermidine synthase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	475	Total	C	H	N	O	S	0	14	0
			7502	2410	3701	662	710	19			
1	B	475	Total	C	H	N	O	S	0	10	0
			7434	2390	3662	656	709	17			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP O32323
A	0	PRO	-	expression tag	UNP O32323
B	-1	GLY	-	expression tag	UNP O32323
B	0	PRO	-	expression tag	UNP O32323

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	
			70	21	26	7	14	2	
2	B	1	Total	C	H	N	O	P	
			70	21	26	7	14	2	

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	H	O		
			7	2	3	2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	527	Total	O		
			527	527	0	0
4	B	508	Total	O		
			508	508	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Homospermidine synthase



- Molecule 1: Homospermidine synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.05Å 109.79Å 192.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.49 47.71 – 1.31	Depositor EDS
% Data completeness (in resolution range)	98.6 (10.00-1.49) 67.7 (47.71-1.31)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.18	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.10 (at 1.31Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.191 , 0.225 0.203 , 0.232	Depositor DCC
$R_{free}$ test set	12039 reflections (6.36%)	DCC
Wilson B-factor (Å <sup>2</sup> )	10.8	Xtriage
Anisotropy	0.335	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 50.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 307834 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	16118	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.62	0/3918	0.71	1/5338 (0.0%)
1	B	0.59	0/3880	0.72	0/5286
All	All	0.61	0/7798	0.71	1/10624 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	191	ARG	NE-CZ-NH2	-5.16	117.72	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3801	3701	3695	13	0
1	B	3772	3662	3657	15	0
2	A	44	26	26	0	0
2	B	44	26	26	0	0
3	B	4	3	3	1	0
4	A	527	0	0	7	3
4	B	508	0	0	9	3
All	All	8700	7418	7407	29	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:466:ASP:OD1	4:A:1126:HOH:O	1.94	0.85
1:B:38:ARG:NH2	4:B:601:HOH:O	2.10	0.85
1:A:101[A]:MET:SD	4:A:902:HOH:O	2.38	0.81
1:A:204:LYS:NZ	4:A:601:HOH:O	2.19	0.74
1:B:40[A]:LYS:NZ	4:B:603:HOH:O	2.22	0.73
1:B:133:ARG:NH1	4:B:1074:HOH:O	2.26	0.69
1:A:344:ARG:NH2	4:A:995:HOH:O	2.31	0.63
1:B:24[B]:ARG:HE	1:B:53:LEU:CD2	2.13	0.61
1:B:385[B]:GLU:OE2	1:B:388:ARG:NH2	2.35	0.59
1:A:9[B]:ARG:NE	4:A:1064:HOH:O	2.37	0.58
1:B:416:ASN:OD1	4:B:936:HOH:O	2.18	0.56
1:A:37:ASP:OD1	1:A:39:SER:OG	2.13	0.56
1:B:385[B]:GLU:OE1	4:B:889:HOH:O	2.18	0.55
1:B:24[B]:ARG:NH2	4:B:604:HOH:O	2.38	0.55
1:B:251:ARG:NH2	1:B:463:ASP:OD2	2.39	0.54
1:A:196[A]:ARG:NH1	4:A:603:HOH:O	2.42	0.53
1:A:27:LEU:HD21	1:A:59[B]:VAL:HG21	1.91	0.52
3:B:502:ACT:OXT	4:B:1078:HOH:O	2.20	0.50
1:B:3:ASP:OD2	4:B:602:HOH:O	2.19	0.50
1:B:223:ASP:OD2	1:B:346[A]:SER:OG	2.25	0.49
1:A:38:ARG:NH2	1:A:58:GLY:O	2.47	0.48
1:B:191:ARG:NH2	4:B:815:HOH:O	2.47	0.47
1:B:27:LEU:CD1	1:B:41[A]:LEU:HD21	2.45	0.46
1:A:447[A]:ASP:OD1	4:A:1068:HOH:O	2.21	0.46
1:A:27:LEU:HD21	1:A:59[A]:VAL:HG11	1.98	0.45
1:B:129:LYS:HD3	1:B:131:GLU:H	1.83	0.44
1:B:129:LYS:HG2	1:B:130:PRO:HD2	1.98	0.44
1:A:127:ASP:N	1:A:127:ASP:OD1	2.49	0.44
1:A:435:THR:N	1:A:436:PRO:CD	2.82	0.43

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:645:HOH:O	4:B:663:HOH:O[1_455]	2.05	0.15
4:A:680:HOH:O	4:B:668:HOH:O[1_455]	2.15	0.05

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:697:HOH:O	4:B:671:HOH:O[1_455]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	486/479 (102%)	470 (97%)	16 (3%)	0	100	100
1	B	483/479 (101%)	470 (97%)	12 (2%)	1 (0%)	52	25
All	All	969/958 (101%)	940 (97%)	28 (3%)	1 (0%)	56	26

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	183	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	403/392 (103%)	391 (97%)	12 (3%)	48	15
1	B	399/392 (102%)	389 (98%)	10 (2%)	55	20
All	All	802/784 (102%)	780 (97%)	22 (3%)	51	18

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	HIS
1	A	69	ASP
1	A	87	PHE
1	A	128	LEU
1	A	131	GLU
1	A	249	PHE
1	A	273	PRO
1	A	293	LEU
1	A	296	HIS
1	A	323	TYR
1	A	344	ARG
1	A	476	ARG
1	B	60	ARG
1	B	87	PHE
1	B	125	ASP
1	B	128	LEU
1	B	129	LYS
1	B	191	ARG
1	B	196	ARG
1	B	244	LEU
1	B	296	HIS
1	B	323	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAD	A	501	-	38,48,48	0.67	0	47,73,73	1.87	6 (12%)
2	NAD	B	501	-	38,48,48	0.92	1 (2%)	47,73,73	1.58	9 (19%)
3	ACT	B	502	-	1,3,3	1.44	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	501	-	-	0/22/62/62	0/5/5/5
2	NAD	B	501	-	-	0/22/62/62	0/5/5/5
3	ACT	B	502	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	NAD	C5A-C4A	2.59	1.46	1.40

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	NAD	N3A-C2A-N1A	-8.22	122.60	128.89
2	B	501	NAD	N3A-C2A-N1A	-5.29	124.84	128.89
2	A	501	NAD	C4B-O4B-C1B	-4.18	105.13	109.72
2	B	501	NAD	C4B-O4B-C1B	-3.09	106.33	109.72
2	A	501	NAD	PN-O3-PA	-3.01	124.26	132.73
2	A	501	NAD	C1B-N9A-C4A	-2.82	122.69	126.94
2	A	501	NAD	O7N-C7N-C3N	-2.77	116.56	119.59
2	B	501	NAD	O7N-C7N-C3N	-2.77	116.56	119.59
2	B	501	NAD	PN-O3-PA	-2.43	125.90	132.73
2	B	501	NAD	C1B-N9A-C4A	-2.42	123.30	126.94
2	B	501	NAD	C2N-C3N-C4N	2.07	120.59	118.29
2	B	501	NAD	O4B-C1B-N9A	2.10	112.48	108.10

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	NAD	N6A-C6A-N1A	2.45	124.45	119.20
2	B	501	NAD	C5N-C6N-N1N	2.46	124.72	120.47
2	A	501	NAD	C3N-C7N-N7N	4.71	122.97	117.82

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	502	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	475/479 (99%)	-0.21	16 (3%) 49 51	14, 24, 50, 92	0
1	B	475/479 (99%)	-0.28	17 (3%) 46 49	15, 24, 52, 99	0
All	All	950/958 (99%)	-0.25	33 (3%) 48 50	14, 24, 51, 99	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	126	PRO	10.3
1	A	126	PRO	8.6
1	A	124	PHE	8.6
1	B	124	PHE	7.8
1	A	127	ASP	6.4
1	B	127	ASP	6.0
1	A	184	THR	5.8
1	B	184	THR	5.6
1	B	311	ALA	5.1
1	A	186	GLU	4.2
1	A	312	ALA	3.9
1	B	477	ASP	3.9
1	A	82	GLY	3.8
1	B	186	GLU	3.8
1	A	311	ALA	3.8
1	B	185	GLY	3.4
1	B	313	GLY	3.3
1	A	457	LEU	3.2
1	A	185	GLY	3.1
1	B	128	LEU	3.0
1	B	125	ASP	3.0
1	A	83	PRO	3.0
1	A	125	ASP	2.9
1	B	312	ALA	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	183	VAL	2.5
1	B	122	PHE	2.5
1	B	129	LYS	2.5
1	A	261	ASP	2.4
1	B	314	GLN	2.3
1	B	261	ASP	2.3
1	A	314	GLN	2.3
1	A	3[A]	ASP	2.2
1	A	129	LYS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	ACT	B	502	4/4	0.94	0.07	-0.05	29,38,47,60	0
2	NAD	A	501	44/44	0.97	0.06	-0.40	14,22,28,33	0
2	NAD	B	501	44/44	0.97	0.06	-0.72	15,22,30,31	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.