



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:41 PM GMT

PDB ID : 4PO6
Title : Crystal structure of the human TYK2 FERM and SH2 domains with an IF-NAR1 intracellular peptide
Authors : Wallweber, H.J.A.; Lupardus, P.J.
Deposited on : 2014-02-25
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	567	
2	B	34	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	602	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4154 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Non-receptor tyrosine-protein kinase TYK2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	475	Total	C	N	O	S	0	3	0
			3818	2453	690	649	26			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	21	GLY	-	EXPRESSION TAG	UNP P29597
A	22	SER	-	EXPRESSION TAG	UNP P29597
A	584	GLY	-	EXPRESSION TAG	UNP P29597
A	585	SER	-	EXPRESSION TAG	UNP P29597
A	586	GLY	-	EXPRESSION TAG	UNP P29597
A	587	SER	-	EXPRESSION TAG	UNP P29597

- Molecule 2 is a protein called Interferon alpha/beta receptor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	19	Total	C	N	O	S	0	0	0
			151	96	22	32	1			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	474	GLY	-	EXPRESSION TAG	UNP P17181
B	475	SER	-	EXPRESSION TAG	UNP P17181
B	476	GLY	-	EXPRESSION TAG	UNP P17181
B	477	SER	-	EXPRESSION TAG	UNP P17181

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	163	Total	O	0	0
			163	163		
4	B	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Non-receptor tyrosine-protein kinase TYK2



- Molecule 2: Interferon alpha/beta receptor 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	186.77Å 52.42Å 70.16Å 90.00° 108.09° 90.00°	Depositor
Resolution (Å)	38.55 – 1.99 38.55 – 1.99	Depositor EDS
% Data completeness (in resolution range)	95.6 (38.55-1.99) 96.1 (38.55-1.99)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 2.00Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.199 , 0.234 0.210 , 0.249	Depositor DCC
R_{free} test set	2137 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	35.0	Xtriage
Anisotropy	0.136	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 57.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 42642 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4154	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	1/3937 (0.0%)	0.67	0/5339
2	B	0.58	0/151	0.83	0/202
All	All	0.54	1/4088 (0.0%)	0.67	0/5541

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	121	ALA	C-N	-5.53	1.21	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3818	0	3782	21	0
2	B	151	0	149	0	0
3	A	18	0	24	2	0
4	A	163	0	0	2	0
4	B	4	0	0	0	0
All	All	4154	0	3955	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:394:SER:OG	1:A:402[A]:CYS:SG	2.46	0.72
1:A:279:HIS:HD2	4:A:784:HOH:O	1.73	0.71
1:A:71:PHE:HB3	3:A:601:GOL:H31	1.77	0.65
1:A:253:GLN:O	1:A:257:VAL:HG23	1.99	0.63
1:A:74:PHE:CE2	1:A:105:ILE:HG13	2.37	0.59
1:A:390:GLU:HG3	1:A:391:HIS:H	1.70	0.56
1:A:158:HIS:HE1	4:A:829:HOH:O	1.90	0.54
1:A:189:LEU:HD11	1:A:258:LYS:HE3	1.94	0.50
1:A:421:ASP:CG	1:A:425:ARG:HH22	2.14	0.50
1:A:529:LEU:C	1:A:529:LEU:HD23	2.33	0.49
1:A:165:ALA:HB2	1:A:267:ALA:HB2	1.95	0.49
1:A:533:LEU:HG	1:A:547:LEU:HD13	1.93	0.49
1:A:457:GLU:HG3	1:A:502:LEU:HD12	1.99	0.44
1:A:274[B]:ARG:HB2	3:A:603:GOL:H11	2.00	0.43
1:A:291:CYS:SG	1:A:392[A]:CYS:SG	3.13	0.43
1:A:523:PHE:HD2	1:A:529:LEU:HD12	1.83	0.43
1:A:155:GLN:O	1:A:159:GLU:HG3	2.19	0.43
1:A:523:PHE:CD2	1:A:529:LEU:HD12	2.54	0.42
1:A:378:CYS:HB3	1:A:397:ARG:HD2	2.02	0.41
1:A:524:PRO:HG2	1:A:528:GLU:HG3	2.01	0.41
1:A:379:ASP:OD2	1:A:565:ARG:NH2	2.43	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	468/567 (82%)	456 (97%)	10 (2%)	2 (0%)	39 33

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	17/34 (50%)	16 (94%)	1 (6%)	0	100	100
All	All	485/601 (81%)	472 (97%)	11 (2%)	2 (0%)	39	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	495	ASP
1	A	517	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	409/478 (86%)	406 (99%)	3 (1%)	88	91
2	B	18/32 (56%)	17 (94%)	1 (6%)	26	20
All	All	427/510 (84%)	423 (99%)	4 (1%)	84	88

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	287	GLU
1	A	293	ILE
1	A	327	TRP
2	B	502	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	226	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	601	-	5,5,5	0.15	0	5,5,5	0.34	0
3	GOL	A	602	-	5,5,5	0.13	0	5,5,5	0.24	0
3	GOL	A	603	-	5,5,5	0.12	0	5,5,5	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	601	-	-	0/4/4/4	0/0/0/0
3	GOL	A	602	-	-	0/4/4/4	0/0/0/0
3	GOL	A	603	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	601	GOL	1	0
3	A	603	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	475/567 (83%)	0.84	76 (16%) 3 3	24, 45, 86, 102	0
2	B	19/34 (55%)	1.20	4 (21%) 1 1	49, 78, 97, 109	0
All	All	494/601 (82%)	0.85	80 (16%) 3 3	24, 46, 87, 109	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	250	LEU	6.0
1	A	496	GLY	5.9
1	A	244	ASP	5.7
1	A	494	PRO	5.5
1	A	372	PRO	5.2
2	B	503	PHE	5.2
1	A	495	ASP	5.0
1	A	246	GLN	4.4
1	A	531	ALA	4.1
1	A	105	ILE	4.0
1	A	497	MET	3.9
1	A	232	LEU	3.8
1	A	515	VAL	3.8
1	A	129	GLY	3.7
1	A	245	PHE	3.7
1	A	248	GLY	3.6
1	A	290	PRO	3.6
1	A	247	PRO	3.6
1	A	207	LYS	3.5
1	A	107	PHE	3.4
1	A	240	ARG	3.4
1	A	272	THR	3.4
1	A	200	ILE	3.4
1	A	152	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	371	GLU	3.3
1	A	318	VAL	3.3
1	A	243	ARG	3.2
1	A	270	PHE	3.1
1	A	493	ALA	3.1
1	A	112	TRP	3.1
1	A	519	TRP	3.1
1	A	263	LEU	3.0
1	A	309	ALA	3.0
1	A	38	GLY	3.0
1	A	513	ALA	2.9
1	A	249	ARG	2.9
1	A	498	GLN	2.9
1	A	555	PRO	2.9
1	A	168	TRP	2.9
2	B	501	LYS	2.8
1	A	109	PHE	2.8
2	B	495	SER	2.8
1	A	327	TRP	2.7
1	A	373	LEU	2.7
1	A	293	ILE	2.7
1	A	544	CYS	2.7
1	A	492	GLN	2.7
1	A	233	ARG	2.7
1	A	241	PHE	2.6
1	A	271	GLY	2.6
1	A	238	PHE	2.6
1	A	520	GLY	2.6
1	A	74	PHE	2.6
1	A	156	GLY	2.5
1	A	426	LEU	2.5
1	A	267	ALA	2.5
1	A	108	TYR	2.5
1	A	499	SER	2.4
1	A	523	PHE	2.4
1	A	73	LEU	2.4
1	A	201	PRO	2.4
1	A	534	GLN	2.3
1	A	203	GLU	2.3
1	A	319	THR	2.3
1	A	84	TRP	2.2
2	B	502	CYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	418	SER	2.2
1	A	269	ARG	2.2
1	A	287	GLU	2.2
1	A	110	ARG	2.2
1	A	143	LEU	2.2
1	A	242	LEU	2.2
1	A	115	MET	2.1
1	A	155	GLN	2.1
1	A	77	PHE	2.1
1	A	524	PRO	2.1
1	A	324	ILE	2.0
1	A	202	LEU	2.0
1	A	522	SER	2.0
1	A	144	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GOL	A	602	6/6	0.92	0.17	2.10	47,54,60,63	0
3	GOL	A	601	6/6	0.85	0.20	1.70	60,62,62,64	0
3	GOL	A	603	6/6	0.80	0.22	-0.04	59,67,72,74	0

6.5 Other polymers [i](#)

There are no such residues in this entry.