



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:45 AM GMT

PDB ID : 3PP7
Title : Crystal structure of Leishmania mexicana pyruvate kinase in complex with the drug suramin, an inhibitor of glycolysis.
Authors : Morgan, H.P.; Auld, D.S.; McNae, I.W.; Nowicki, M.W.; Michels, P.A.M.; Fothergill-Gilmore, L.A.; Walkinshaw, M.D.
Deposited on : 2010-11-24
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1352 (2.38-2.34)
Clashscore	102246	1456 (2.38-2.34)
Ramachandran outliers	100387	1435 (2.38-2.34)
Sidechain outliers	100360	1436 (2.38-2.34)
RSRZ outliers	91569	1358 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	498	 5% 89% 9% ..
1	B	498	 24% 87% 11% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	500	-	-	-	X
5	SVR	B	499	-	-	-	X

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 8054 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyruvate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	492	Total	C	N	O	S	0	1	0
			3763	2345	663	729	26			
1	B	494	Total	C	N	O	S	0	0	0
			3766	2347	663	730	26			

- Molecule 2 is PYRENE-1,3,6,8-TETRASULFONIC ACID (three-letter code: PTK) (formula: $C_{16}H_{10}O_{12}S_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	S	0	0
			32	16	12	4		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	K	0	0
			2	2		
4	A	2	Total	K	0	0
			2	2		

- Molecule 5 is 8,8'-[CARBONYLBIS[IMINO-3,1-PHENYLENECARBONYLIMINO(4-METHYL-3,1-PHENYLENE)CARBONYLIMINO]]BIS-1,3,5-NAPHTHALENETRISULFONIC ACID (three-letter code: SVR) (formula: C₅₁H₄₀N₆O₂₃S₆).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	S	0	0
			41	25	2	11	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	236	Total	O	0	0
			236	236		
6	B	200	Total	O	0	0
			200	200		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pyruvate kinase



• Molecule 1: Pyruvate kinase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	123.77Å 129.48Å 165.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.73 – 2.35 44.73 – 2.35	Depositor EDS
% Data completeness (in resolution range)	98.0 (44.73-2.35) 98.0 (44.73-2.35)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.02	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.55 (at 2.34Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.191 , 0.231 0.208 , 0.247	Depositor DCC
R_{free} test set	2761 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	41.3	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 55.8	EDS
Estimated twinning fraction	0.010 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 54537 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8054	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SVR, GOL, K, PTK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/3819	0.55	0/5169
1	B	0.42	0/3821	0.56	0/5170
All	All	0.42	0/7640	0.55	0/10339

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3763	0	3758	32	0
1	B	3766	0	3768	33	0
2	A	32	0	8	0	0
3	A	6	0	8	1	0
3	B	6	0	8	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	B	41	0	18	3	0
6	A	236	0	0	2	0
6	B	200	0	0	1	0
All	All	8054	0	7568	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:26:THR:HG21	1:B:49:ARG:HH11	1.38	0.87
1:A:263:GLY:H	1:A:296:THR:HG21	1.39	0.85
1:B:26:THR:HG23	1:B:330:SER:HA	1.62	0.81
1:A:479:ILE:HD11	1:A:489:ALA:CB	2.13	0.78
1:B:26:THR:HG21	1:B:49:ARG:NH1	2.00	0.74
1:A:295:ALA:O	1:A:296:THR:HB	1.85	0.74
1:B:26:THR:CG2	1:B:330:SER:HA	2.18	0.73
1:B:29:PRO:HG3	5:B:499:SVR:H56	1.71	0.72
1:A:223:ARG:HD2	6:A:609:HOH:O	1.90	0.71
1:A:263:GLY:N	1:A:296:THR:HG21	2.06	0.70
1:B:268:GLU:HG2	6:B:624:HOH:O	1.91	0.69
1:B:26:THR:CG2	1:B:49:ARG:HH11	2.08	0.67
1:A:101:MET:HE1	1:A:121:LYS:HA	1.77	0.67
1:A:377:ASP:HB3	1:A:488:TYR:CD2	2.30	0.65
1:B:135:VAL:HG11	1:B:152:VAL:HG21	1.81	0.63
1:A:135:VAL:HG11	1:A:152:VAL:HG21	1.84	0.59
1:B:101:MET:HE1	1:B:121:LYS:HA	1.86	0.57
1:A:296:THR:HG22	1:A:297:GLN:HG3	1.86	0.57
1:B:298:MET:HE3	1:B:327:VAL:HB	1.86	0.56
1:B:26:THR:CG2	1:B:329:LEU:O	2.53	0.56
5:B:499:SVR:N63	5:B:499:SVR:S73	2.73	0.55
1:A:404:ARG:CG	1:A:404:ARG:HH21	2.19	0.55
1:A:108:TYR:O	1:A:123:LYS:HA	2.09	0.53
1:B:108:TYR:O	1:B:123:LYS:HA	2.08	0.53
1:B:268:GLU:HG3	1:B:269:ILE:HG13	1.91	0.52
1:B:374:MET:CE	1:B:378:GLU:HG3	2.40	0.52
1:A:54[B]:HIS:C	1:A:54[B]:HIS:CD2	2.84	0.51
1:A:263:GLY:CA	1:A:296:THR:HG21	2.41	0.51
1:B:237:CYS:SG	1:B:255:SER:HB3	2.51	0.50
1:B:398:VAL:HG13	1:B:479:ILE:HG12	1.92	0.50
1:A:479:ILE:HD11	1:A:489:ALA:HB1	1.90	0.49
1:B:26:THR:HG23	1:B:329:LEU:O	2.12	0.49
1:A:392:LYS:O	1:A:394:LYS:HE3	2.12	0.48
1:B:394:LYS:HB2	1:B:470:VAL:HG12	1.95	0.48
1:A:88:GLU:HA	3:A:500:GOL:H2	1.95	0.48
1:B:29:PRO:CG	5:B:499:SVR:H56	2.42	0.48
1:A:263:GLY:HA3	1:A:296:THR:HG21	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:394:LYS:HB2	1:A:470:VAL:HG12	1.97	0.47
1:A:404:ARG:HH21	1:A:404:ARG:HG3	1.79	0.46
1:B:142:TYR:HB3	1:B:146:GLY:HA2	1.98	0.46
1:B:26:THR:HG22	1:B:329:LEU:O	2.15	0.46
1:A:142:TYR:HB3	1:A:146:GLY:HA2	1.98	0.45
1:B:365:SER:HA	1:B:368:LYS:HE3	1.98	0.45
1:B:112:ASP:HA	1:B:113:PRO:HD3	1.86	0.45
1:A:404:ARG:CG	1:A:404:ARG:NH2	2.79	0.44
1:B:298:MET:CE	1:B:327:VAL:HB	2.45	0.44
1:B:134:VAL:CG1	1:B:182:CYS:HB3	2.47	0.44
1:B:17:ASN:N	1:B:17:ASN:HD22	2.15	0.44
1:B:17:ASN:H	1:B:17:ASN:HD22	1.66	0.43
1:B:385:VAL:HG21	1:B:412:TYR:HB2	1.99	0.43
1:B:374:MET:HE2	1:B:378:GLU:HG3	2.00	0.43
1:A:134:VAL:CG1	1:A:182:CYS:HB3	2.48	0.43
1:A:263:GLY:H	1:A:296:THR:CG2	2.20	0.42
1:A:490:ASN:N	1:A:490:ASN:HD22	2.18	0.42
1:A:58:GLU:HB2	6:A:529:HOH:O	2.19	0.42
1:A:38:LYS:CE	1:A:73:GLU:OE1	2.69	0.41
1:A:377:ASP:HB3	1:A:488:TYR:CE2	2.56	0.41
1:B:452:GLY:O	1:B:453:LYS:HB2	2.20	0.41
1:B:24:ILE:HG12	1:B:47:VAL:HB	2.03	0.41
1:A:451:GLU:O	1:A:453:LYS:HE2	2.21	0.41
1:A:310:ARG:NH2	1:B:315:ASP:OD1	2.55	0.40
1:B:93:GLN:HB2	1:B:117:ASP:HA	2.02	0.40
1:A:238:LYS:HG2	1:A:259:MET:SD	2.62	0.40
1:A:63:THR:O	1:A:67:VAL:HG23	2.21	0.40
1:A:40:LEU:HD12	1:A:40:LEU:HA	1.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	489/498 (98%)	479 (98%)	9 (2%)	1 (0%)	52	63
1	B	490/498 (98%)	477 (97%)	12 (2%)	1 (0%)	52	63
All	All	979/996 (98%)	956 (98%)	21 (2%)	2 (0%)	52	63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	487	GLY
1	A	296	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	412/416 (99%)	393 (95%)	19 (5%)	33	42
1	B	412/416 (99%)	391 (95%)	21 (5%)	29	36
All	All	824/832 (99%)	784 (95%)	40 (5%)	31	39

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	LEU
1	A	111	THR
1	A	118	LYS
1	A	136	ARG
1	A	174	ARG
1	A	175	ARG
1	A	177	VAL
1	A	179	LEU
1	A	224	LYS
1	A	262	ARG
1	A	297	GLN
1	A	298	MET
1	A	367	LYS
1	A	371	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	404	ARG
1	A	435	GLN
1	A	453	LYS
1	A	479	ILE
1	A	490	ASN
1	B	17	ASN
1	B	26	THR
1	B	34	VAL
1	B	40	LEU
1	B	111	THR
1	B	118	LYS
1	B	136	ARG
1	B	174	ARG
1	B	175	ARG
1	B	177	VAL
1	B	179	LEU
1	B	212	PHE
1	B	262	ARG
1	B	264	ASP
1	B	307	ARG
1	B	314	SER
1	B	371	HIS
1	B	435	GLN
1	B	446	LYS
1	B	454	GLU
1	B	479	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	HIS
1	A	178	ASN
1	A	490	ASN
1	B	17	ASN
1	B	242	HIS
1	B	305	ASN
1	B	364	ASN
1	B	455	HIS
1	B	471	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	PTK	A	499	-	35,35,35	1.57	7 (20%)	50,60,60	1.62	12 (24%)
3	GOL	A	500	-	5,5,5	0.23	0	5,5,5	0.59	0
5	SVR	B	499	-	44,44,93	1.26	5 (11%)	62,68,145	1.50	10 (16%)
3	GOL	B	502	-	5,5,5	0.26	0	5,5,5	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PTK	A	499	-	-	0/24/24/24	0/4/4/4
3	GOL	A	500	-	-	0/4/4/4	0/0/0/0
5	SVR	B	499	-	-	0/34/34/76	0/4/4/8
3	GOL	B	502	-	-	0/4/4/4	0/0/0/0

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	499	SVR	C55-N53	-3.02	1.36	1.41
2	A	499	PTK	CAV-CAZ	-2.84	1.39	1.43
2	A	499	PTK	CAT-CAX	-2.81	1.39	1.43
2	A	499	PTK	CAV-SBF	-2.67	1.74	1.78
2	A	499	PTK	CAS-CAW	-2.40	1.39	1.43
2	A	499	PTK	CAU-CAY	-2.33	1.39	1.43
2	A	499	PTK	CAS-SBC	-2.09	1.75	1.78
2	A	499	PTK	CAT-SBD	-2.07	1.75	1.78
5	B	499	SVR	C71-C68	-2.02	1.40	1.43
5	B	499	SVR	C70-C71	2.12	1.40	1.37
5	B	499	SVR	C49-C51	2.71	1.55	1.50
5	B	499	SVR	C58-C61	3.06	1.56	1.50

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	499	SVR	C72-C68-C71	-4.51	118.03	123.23
2	A	499	PTK	CAQ-CAX-CAT	-3.47	119.22	123.83
2	A	499	PTK	CAR-CAZ-CAV	-3.15	119.65	123.83
5	B	499	SVR	C70-C71-C68	-2.83	118.52	121.04
2	A	499	PTK	CAO-CAW-CAS	-2.61	120.37	123.83
2	A	499	PTK	CAP-CAY-CAU	-2.45	120.58	123.83
5	B	499	SVR	C57-C55-N53	2.40	122.46	118.71
5	B	499	SVR	O77-S73-C69	2.43	108.95	106.20
5	B	499	SVR	C70-C71-S75	2.55	121.26	117.51
2	A	499	PTK	OAH-SBF-CAV	2.72	109.28	106.20
5	B	499	SVR	C71-C68-C66	2.73	121.41	118.11
2	A	499	PTK	OAD-SBD-CAT	2.80	109.37	106.20
2	A	499	PTK	OAG-SBF-CAV	2.82	109.39	106.20
2	A	499	PTK	OAI-SBC-CAS	2.85	109.43	106.20
2	A	499	PTK	OAA-SBC-CAS	2.86	109.44	106.20
2	A	499	PTK	OAJ-SBD-CAT	2.88	109.47	106.20
2	A	499	PTK	OAE-SBE-CAU	2.98	109.58	106.20
5	B	499	SVR	C60-C57-C55	3.02	119.76	117.39
2	A	499	PTK	OAF-SBE-CAU	3.17	109.79	106.20
5	B	499	SVR	O78-S73-C69	3.58	110.25	106.20
5	B	499	SVR	O81-S75-C71	3.74	110.44	106.20
5	B	499	SVR	O80-S75-C71	4.47	111.27	106.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	500	GOL	1	0
5	B	499	SVR	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	492/498 (98%)	0.41	24 (4%)	33 48	38, 55, 94, 109	0
1	B	494/498 (99%)	1.79	118 (23%)	1 1	37, 56, 215, 243	0
All	All	986/996 (98%)	1.10	142 (14%)	3 6	37, 55, 196, 243	0

All (142) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	109	VAL	22.9
1	B	171	ILE	22.2
1	B	124	PHE	17.9
1	B	126	ILE	17.7
1	B	172	SER	16.4
1	B	170	THR	15.4
1	B	92	GLY	15.1
1	B	94	PHE	15.0
1	B	106	THR	14.3
1	B	104	GLY	13.7
1	B	95	VAL	12.4
1	B	138	GLY	12.2
1	B	169	HIS	12.0
1	B	108	TYR	11.0
1	B	137	PRO	10.8
1	B	89	ILE	10.8
1	B	91	THR	9.8
1	B	107	CYS	9.6
1	B	163	CYS	9.5
1	B	164	THR	9.4
1	B	119	GLY	9.4
1	B	175	ARG	9.3
1	B	149	ILE	9.3
1	B	105	ALA	9.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	93	GLN	9.1
1	B	165	VAL	9.0
1	B	142	TYR	8.9
1	B	101	MET	8.8
1	B	167	ASN	8.8
1	B	143	ILE	8.7
1	B	122	ASP	8.3
1	B	99	ALA	8.2
1	B	103	ARG	8.2
1	B	90	ARG	8.1
1	B	133	LYS	8.0
1	B	152	VAL	7.9
1	B	134	VAL	7.9
1	B	150	LEU	7.9
1	B	125	TYR	7.7
1	B	140	TYR	7.6
1	B	96	GLY	7.5
1	B	181	GLY	7.5
1	B	168	SER	7.4
1	B	156	GLU	7.3
1	B	161	LEU	7.2
1	B	131	LEU	7.2
1	B	151	GLN	7.1
1	B	148	LEU	7.0
1	B	139	ASN	7.0
1	B	102	GLU	6.8
1	B	141	ILE	6.7
1	B	160	THR	6.7
1	B	136	ARG	6.6
1	B	135	VAL	6.4
1	B	173	ASP	6.4
1	B	162	GLU	6.4
1	B	153	GLN	6.1
1	B	155	HIS	6.1
1	B	121	LYS	6.0
1	B	166	THR	6.0
1	B	158	GLU	5.9
1	B	154	SER	5.9
1	B	100	VAL	5.8
1	B	146	GLY	5.8
1	B	157	ASP	5.8
1	B	115	PHE	5.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	113	PRO	5.5
1	B	110	THR	5.4
1	B	174	ARG	5.3
1	B	114	ALA	5.2
1	B	182	CYS	5.0
1	B	177	VAL	5.0
1	B	117	ASP	5.0
1	B	116	ALA	5.0
1	B	179	LEU	4.8
1	B	132	SER	4.6
1	B	159	GLN	4.5
1	B	97	GLY	4.5
1	B	180	PRO	4.4
1	B	147	ILE	4.3
1	B	98	ASP	4.1
1	B	188	ALA	3.9
1	B	120	THR	3.8
1	B	129	GLN	3.8
1	B	488	TYR	3.7
1	B	189	VAL	3.6
1	B	118	LYS	3.6
1	B	144	ASP	3.5
1	B	123	LYS	3.3
1	B	130	ASN	3.3
1	B	88	GLU	3.2
1	B	319	ALA	3.0
1	B	447	LEU	3.0
1	B	183	ASP	3.0
1	A	488	TYR	3.0
1	B	372	ILE	3.0
1	B	128	TYR	2.9
1	B	316	VAL	2.9
1	B	112	ASP	2.9
1	A	282	ILE	2.8
1	B	317	ALA	2.8
1	B	444	ALA	2.8
1	B	127	ASP	2.8
1	A	327	VAL	2.6
1	B	327	VAL	2.6
1	A	12	PHE	2.6
1	A	98	ASP	2.6
1	A	317	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	313	VAL	2.6
1	B	176	GLY	2.5
1	A	326	CYS	2.5
1	A	292	VAL	2.5
1	B	145	ASP	2.5
1	A	294	CYS	2.4
1	A	94	PHE	2.4
1	A	311	ALA	2.4
1	B	371	HIS	2.4
1	B	318	ASN	2.4
1	A	275	VAL	2.3
1	B	87	PRO	2.3
1	B	486	LYS	2.3
1	B	321	PHE	2.3
1	B	324	ALA	2.2
1	A	324	ALA	2.2
1	B	186	LEU	2.2
1	A	320	VAL	2.2
1	B	322	ASN	2.1
1	A	174	ARG	2.1
1	B	446	LYS	2.1
1	A	97	GLY	2.1
1	B	17	ASN	2.1
1	B	293	ILE	2.1
1	B	449	HIS	2.1
1	A	23	ILE	2.1
1	A	433	ILE	2.1
1	A	321	PHE	2.0
1	B	350	CYS	2.0
1	A	283	SER	2.0
1	A	293	ILE	2.0
1	A	135	VAL	2.0
1	A	316	VAL	2.0
1	A	410	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GOL	A	500	6/6	0.72	0.25	3.98	61,62,62,62	0
5	SVR	B	499	41/86	0.79	0.26	3.03	104,106,108,108	0
3	GOL	B	502	6/6	0.74	0.25	1.90	80,81,82,82	0
4	K	A	502	1/1	0.88	0.11	-0.54	81,81,81,81	0
4	K	B	500	1/1	0.96	0.10	-2.09	66,66,66,66	0
4	K	A	501	1/1	0.94	0.06	-3.02	55,55,55,55	0
4	K	B	501	1/1	0.99	0.05	-5.51	58,58,58,58	0
2	PTK	A	499	32/32	0.78	0.20	-	118,118,119,119	32

6.5 Other polymers [i](#)

There are no such residues in this entry.