



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 09:38 PM GMT

PDB ID : 1PXX
Title : CRYSTAL STRUCTURE OF DICLOFENAC BOUND TO THE CYCLOOXYGENASE ACTIVE SITE OF COX-2
Authors : Kiefer, J.R.; Rowlinson, S.W.; Prusakiewicz, J.J.; Pawlitz, J.L.; Kozak, K.R.; Kalgutkar, A.S.; Stallings, W.C.; Marnett, L.J.; Kurumbail, R.G.
Deposited on : 2003-07-07
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	604	<div> <div>2%</div> <div>50% 35% 6% 9%</div> </div>
1	B	604	<div> <div>2%</div> <div>49% 36% 6% 9%</div> </div>
1	C	604	<div> <div>%</div> <div>50% 36% 6% 9%</div> </div>
1	D	604	<div> <div>2%</div> <div>50% 36% 6% 9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	661	-	-	-	X
2	NAG	B	1661	-	-	-	X
2	NAG	C	2661	-	-	X	X
2	NAG	D	3661	-	-	-	X
2	NAG	D	3681	-	-	-	X
4	BOG	D	3703	-	-	X	-
6	DIF	C	2701	-	-	-	X

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 18778 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Prostaglandin G/H synthase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	552	Total	C	N	O	S	0	0	0
			4472	2885	748	814	25			
1	B	552	Total	C	N	O	S	0	0	0
			4474	2885	750	814	25			
1	C	552	Total	C	N	O	S	0	0	0
			4474	2885	750	814	25			
1	D	552	Total	C	N	O	S	0	0	0
			4474	2885	750	814	25			

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is a polymer of unknown type called SUGAR (3-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	3	Total	C	N	O	0	0
			42	24	3	15		
3	B	3	Total	C	N	O	0	0
			42	24	3	15		
3	C	3	Total	C	N	O	0	0
			42	24	3	15		
3	D	3	Total	C	N	O	0	0
			42	24	3	15		

- Molecule 4 is SUGAR (B-OCTYLGLUCOSIDE) (three-letter code: BOG) (formula: C₁₄H₂₈O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			20	14	6		
4	D	1	Total	C	O	0	0
			19	13	6		

- Molecule 5 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
5	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 6 is 2-[2,6-DICHLOROPHENYL)AMINO]BENZENEACETIC ACID (three-letter code: DIF) (formula: $C_{14}H_{11}Cl_2NO_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	Cl	N	O	0	0
			19	14	2	1	2		
6	B	1	Total	C	Cl	N	O	0	0
			19	14	2	1	2		
6	C	1	Total	C	Cl	N	O	0	0
			19	14	2	1	2		
6	D	1	Total	C	Cl	N	O	0	0
			19	14	2	1	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	60	Total	O	0	0
			60	60		
7	B	93	Total	O	0	0
			93	93		
7	C	69	Total	O	0	0
			69	69		
7	D	95	Total	O	0	0
			95	95		



• Molecule 1: Prostaglandin G/H synthase 2





4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	181.15Å 135.09Å 124.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.90 19.97 – 2.90	Depositor EDS
% Data completeness (in resolution range)	83.6 (20.00-2.90) 83.3 (19.97-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.16	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.15 (at 2.88Å)	Xtriage
Refinement program	X-PLOR 98.1	Depositor
R, R_{free}	0.254 , 0.302 0.247 , 0.293	Depositor DCC
R_{free} test set	5472 reflections (10.66%)	DCC
Wilson B-factor (Å ²)	21.5	Xtriage
Anisotropy	0.578	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 52.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	13 of 56840 reflections (0.023%)	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	18778	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 64.30 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 8.4672e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, DIF, NAG, BOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/4598	0.66	1/6232 (0.0%)
1	B	0.47	0/4601	0.66	1/6239 (0.0%)
1	C	0.47	0/4601	0.66	1/6239 (0.0%)
1	D	0.47	0/4601	0.66	1/6239 (0.0%)
All	All	0.47	0/18401	0.66	4/24949 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	D	0	1
All	All	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1148	TYR	N-CA-C	-5.71	95.57	111.00
1	D	3148	TYR	N-CA-C	-5.42	96.38	111.00
1	C	2148	TYR	N-CA-C	-5.38	96.47	111.00
1	A	148	TYR	N-CA-C	-5.31	96.67	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	1348	TYR	Sidechain
1	D	3348	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4472	0	4366	231	0
1	B	4474	0	4372	234	1
1	C	4474	0	4372	235	1
1	D	4474	0	4372	227	0
2	A	28	0	26	5	0
2	B	28	0	26	6	0
2	C	28	0	26	7	0
2	D	28	0	26	4	0
3	A	42	0	37	3	0
3	B	42	0	37	1	0
3	C	42	0	37	2	0
3	D	42	0	37	3	0
4	A	20	0	28	5	1
4	D	19	0	25	9	1
5	A	43	0	30	4	0
5	B	43	0	30	7	0
5	C	43	0	30	4	0
5	D	43	0	30	4	0
6	A	19	0	10	3	0
6	B	19	0	10	3	0
6	C	19	0	10	4	0
6	D	19	0	10	1	0
7	A	60	0	0	9	0
7	B	93	0	0	9	0
7	C	69	0	0	8	0
7	D	95	0	0	7	0
All	All	18778	0	17947	928	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 928 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:3294:LEU:HD22	1:D:3409:TYR:HD2	1.03	1.15
1:A:294:LEU:HD22	1:A:409:TYR:CD2	1.80	1.14
1:D:3294:LEU:HD22	1:D:3409:TYR:CD2	1.84	1.12
1:A:294:LEU:HD22	1:A:409:TYR:HD2	1.00	1.11
1:D:3442:ILE:HD13	4:D:3703:BOG:H62	1.34	1.08

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1490:GLU:OE1	4:A:703:BOG:O2[4_555]	2.18	0.02
1:C:2490:GLU:OE1	4:D:3703:BOG:O2[4_456]	2.18	0.02

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	549/604 (91%)	516 (94%)	31 (6%)	2 (0%)	39	74
1	B	550/604 (91%)	513 (93%)	36 (6%)	1 (0%)	52	84
1	C	550/604 (91%)	510 (93%)	37 (7%)	3 (0%)	34	71
1	D	550/604 (91%)	510 (93%)	40 (7%)	0	100	100
All	All	2199/2416 (91%)	2049 (93%)	144 (6%)	6 (0%)	46	79

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	67	GLU
1	C	2106	PRO
1	C	2074	PHE
1	C	2438	ARG
1	A	106	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	492/537 (92%)	432 (88%)	60 (12%)	6	18
1	B	493/537 (92%)	430 (87%)	63 (13%)	5	16
1	C	493/537 (92%)	439 (89%)	54 (11%)	8	23
1	D	493/537 (92%)	430 (87%)	63 (13%)	5	16
All	All	1971/2148 (92%)	1731 (88%)	240 (12%)	6	18

5 of 240 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1463	LEU
1	C	2122	TYR
1	D	3409	TYR
1	B	1484	GLU
1	B	1579	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 57 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	1454	GLN
1	C	2131	ASN
1	D	3370	GLN
1	B	1464	ASN
1	C	2034	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

12 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	671	1,3	14,14,15	0.46	0	15,19,21	0.65	0
3	NAG	A	672	3	14,14,15	0.79	0	15,19,21	0.61	0
3	NAG	A	673	3	14,14,15	0.83	0	15,19,21	0.93	1 (6%)
3	NAG	B	1671	1,3	14,14,15	0.51	0	15,19,21	0.72	1 (6%)
3	NAG	B	1672	3	14,14,15	0.75	0	15,19,21	0.76	1 (6%)
3	NAG	B	1673	3	14,14,15	0.97	1 (7%)	15,19,21	0.89	1 (6%)
3	NAG	C	2671	1,3	14,14,15	0.45	0	15,19,21	0.70	1 (6%)
3	NAG	C	2672	3	14,14,15	0.68	0	15,19,21	0.79	1 (6%)
3	NAG	C	2673	3	14,14,15	0.93	1 (7%)	15,19,21	0.76	0
3	NAG	D	3671	1,3	14,14,15	0.58	0	15,19,21	0.67	0
3	NAG	D	3672	3	14,14,15	0.83	0	15,19,21	0.66	0
3	NAG	D	3673	3	14,14,15	0.93	1 (7%)	15,19,21	0.94	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	671	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	672	3	-	0/6/23/26	0/1/1/1
3	NAG	A	673	3	-	0/6/23/26	0/1/1/1
3	NAG	B	1671	1,3	-	0/6/23/26	0/1/1/1
3	NAG	B	1672	3	-	0/6/23/26	0/1/1/1
3	NAG	B	1673	3	-	0/6/23/26	0/1/1/1
3	NAG	C	2671	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2672	3	-	0/6/23/26	0/1/1/1
3	NAG	C	2673	3	-	0/6/23/26	0/1/1/1
3	NAG	D	3671	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	3672	3	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	D	3673	3	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	3673	NAG	C1-C2	2.31	1.55	1.52
3	C	2673	NAG	C1-C2	2.62	1.56	1.52
3	B	1673	NAG	C1-C2	2.66	1.56	1.52

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1671	NAG	C2-N2-C7	-2.33	120.05	123.04
3	C	2672	NAG	C4-C3-C2	-2.26	107.72	111.23
3	B	1672	NAG	C4-C3-C2	-2.22	107.78	111.23
3	C	2671	NAG	C2-N2-C7	-2.11	120.33	123.04
3	A	673	NAG	C1-O5-C5	2.14	114.97	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	672	NAG	2	0
3	A	673	NAG	3	0
3	B	1672	NAG	1	0
3	B	1673	NAG	1	0
3	C	2672	NAG	1	0
3	C	2673	NAG	2	0
3	D	3671	NAG	1	0
3	D	3672	NAG	2	0
3	D	3673	NAG	2	0

5.6 Ligand geometry

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	661	1	14,14,15	0.53	0	15,19,21	0.74	0
2	NAG	A	681	1	14,14,15	0.57	0	15,19,21	0.77	0
6	DIF	A	701	-	17,20,20	1.10	1 (5%)	24,27,27	2.03	5 (20%)
4	BOG	A	703	-	20,20,20	0.70	1 (5%)	25,25,25	0.63	0
5	HEM	A	704	1	30,50,50	2.78	9 (30%)	24,82,82	2.17	7 (29%)
5	HEM	B	1601	1	30,50,50	2.99	12 (40%)	24,82,82	2.11	8 (33%)
2	NAG	B	1661	1	14,14,15	0.46	0	15,19,21	0.76	0
2	NAG	B	1681	1	14,14,15	0.45	0	15,19,21	0.85	1 (6%)
6	DIF	B	1701	-	17,20,20	0.91	1 (5%)	24,27,27	1.70	4 (16%)
5	HEM	C	2601	1	30,50,50	2.96	10 (33%)	24,82,82	2.00	7 (29%)
2	NAG	C	2661	1	14,14,15	0.58	0	15,19,21	0.69	0
2	NAG	C	2681	1	14,14,15	0.47	0	15,19,21	1.03	1 (6%)
6	DIF	C	2701	-	17,20,20	1.11	1 (5%)	24,27,27	1.51	5 (20%)
5	HEM	D	3601	1	30,50,50	2.88	11 (36%)	24,82,82	2.17	8 (33%)
2	NAG	D	3661	1	14,14,15	0.70	0	15,19,21	0.86	1 (6%)
2	NAG	D	3681	1	14,14,15	0.46	0	15,19,21	0.89	0
6	DIF	D	3701	-	17,20,20	1.12	1 (5%)	24,27,27	1.73	4 (16%)
4	BOG	D	3703	-	17,17,20	0.35	0	17,18,25	0.73	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	661	1	-	0/6/23/26	0/1/1/1
2	NAG	A	681	1	-	0/6/23/26	0/1/1/1
6	DIF	A	701	-	-	0/6/8/8	0/2/2/2
4	BOG	A	703	-	-	0/11/31/31	0/1/1/1
5	HEM	A	704	1	-	0/10/54/54	0/0/8/8
5	HEM	B	1601	1	-	0/10/54/54	0/0/8/8
2	NAG	B	1661	1	-	0/6/23/26	0/1/1/1
2	NAG	B	1681	1	-	0/6/23/26	0/1/1/1
6	DIF	B	1701	-	-	0/6/8/8	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HEM	C	2601	1	-	0/10/54/54	0/0/8/8
2	NAG	C	2661	1	-	0/6/23/26	0/1/1/1
2	NAG	C	2681	1	-	0/6/23/26	0/1/1/1
6	DIF	C	2701	-	-	0/6/8/8	0/2/2/2
5	HEM	D	3601	1	-	0/10/54/54	0/0/8/8
2	NAG	D	3661	1	-	0/6/23/26	0/1/1/1
2	NAG	D	3681	1	-	0/6/23/26	0/1/1/1
6	DIF	D	3701	-	-	0/6/8/8	0/2/2/2
4	BOG	D	3703	-	-	0/18/18/31	0/0/0/1

The worst 5 of 47 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1601	HEM	C3B-C4B	-7.33	1.45	1.51
5	C	2601	HEM	C3B-CAB	-6.84	1.38	1.51
5	C	2601	HEM	C3B-C4B	-6.81	1.45	1.51
5	B	1601	HEM	C3B-CAB	-6.76	1.38	1.51
5	C	2601	HEM	C2D-C3D	-6.68	1.34	1.54

The worst 5 of 52 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	701	DIF	C4-C3-N1	-4.32	115.08	122.09
6	D	3701	DIF	C4-C3-N1	-4.16	115.33	122.09
6	B	1701	DIF	C4-C3-N1	-3.74	116.01	122.09
5	A	704	HEM	CBA-CAA-C2A	-3.34	106.55	112.53
5	B	1601	HEM	CBA-CAA-C2A	-3.18	106.83	112.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

14 monomers are involved in 68 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	661	NAG	5	0
6	A	701	DIF	3	0
4	A	703	BOG	5	1
5	A	704	HEM	4	0
5	B	1601	HEM	7	0
2	B	1661	NAG	6	0
6	B	1701	DIF	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	2601	HEM	4	0
2	C	2661	NAG	7	0
6	C	2701	DIF	4	0
5	D	3601	HEM	4	0
2	D	3661	NAG	4	0
6	D	3701	DIF	1	0
4	D	3703	BOG	9	1

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	552/604 (91%)	0.09	15 (2%) 58 52	5, 21, 42, 53	0
1	B	552/604 (91%)	0.13	15 (2%) 58 52	5, 21, 42, 58	0
1	C	552/604 (91%)	0.05	9 (1%) 74 72	3, 21, 41, 62	0
1	D	552/604 (91%)	0.18	15 (2%) 58 52	5, 21, 42, 55	0
All	All	2208/2416 (91%)	0.12	54 (2%) 62 57	3, 21, 42, 62	0

The worst 5 of 54 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1583	GLN	5.6
1	C	2583	GLN	4.9
1	D	3368	ASN	3.5
1	A	82	LEU	3.0
1	D	3401	GLU	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAG	B	1671	14/15	0.85	0.23	1.22	17,25,39,43	0
3	NAG	D	3671	14/15	0.88	0.22	1.21	18,23,26,36	0
3	NAG	A	671	14/15	0.86	0.20	0.74	18,26,30,38	0
3	NAG	C	2671	14/15	0.91	0.19	-0.46	22,27,40,46	0
3	NAG	A	672	14/15	0.73	0.38	-	44,51,56,57	0
3	NAG	C	2672	14/15	0.75	0.34	-	50,55,62,65	0
3	NAG	B	1673	14/15	0.41	0.61	-	41,65,67,69	0
3	NAG	D	3672	14/15	0.82	0.31	-	44,50,55,59	0
3	NAG	D	3673	14/15	0.35	0.51	-	51,66,71,72	0
3	NAG	A	673	14/15	0.55	0.54	-	58,64,72,73	0
3	NAG	C	2673	14/15	0.37	0.65	-	66,71,73,74	0
3	NAG	B	1672	14/15	0.81	0.32	-	51,57,61,61	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAG	D	3661	14/15	0.79	0.46	6.05	51,55,59,61	0
2	NAG	C	2661	14/15	0.77	0.33	4.23	45,54,61,63	0
2	NAG	A	661	14/15	0.79	0.42	3.25	55,57,61,61	0
2	NAG	D	3681	14/15	0.84	0.36	2.72	47,51,59,62	0
6	DIF	C	2701	19/19	0.89	0.25	2.17	9,29,57,58	0
2	NAG	B	1661	14/15	0.67	0.35	2.13	55,58,61,62	0
6	DIF	A	701	19/19	0.89	0.24	1.68	28,30,46,48	0
6	DIF	B	1701	19/19	0.88	0.23	1.51	13,31,51,52	0
6	DIF	D	3701	19/19	0.88	0.21	1.49	17,38,44,55	0
2	NAG	B	1681	14/15	0.78	0.38	1.44	46,48,56,56	0
2	NAG	C	2681	14/15	0.83	0.33	1.17	44,46,56,57	0
4	BOG	D	3703	19/20	0.89	0.24	0.66	19,22,30,30	0
5	HEM	B	1601	43/43	0.91	0.21	0.39	4,22,48,54	0
5	HEM	D	3601	43/43	0.90	0.21	0.35	13,21,41,43	0
2	NAG	A	681	14/15	0.84	0.27	0.15	44,48,51,52	0
4	BOG	A	703	20/20	0.90	0.23	0.11	17,20,24,25	0
5	HEM	A	704	43/43	0.94	0.17	-0.39	2,15,36,43	0
5	HEM	C	2601	43/43	0.95	0.15	-1.05	8,20,35,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.