



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:48 PM GMT

PDB ID : 4Q0R
Title : The catalytic core of Rad2 (complex I)
Authors : Mietus, M.; Nowak, E.; Jaciuk, M.; Kustosz, P.; Nowotny, M.
Deposited on : 2014-04-02
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3340 (2.80-2.72)
Clashscore	102246	3829 (2.80-2.72)
Ramachandran outliers	100387	3767 (2.80-2.72)
Sidechain outliers	100360	3770 (2.80-2.72)
RSRZ outliers	91569	3352 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	367	<div> <div>2%</div> <div>58%</div> <div>20%</div> <div>•</div> <div>20%</div> </div>
1	B	367	<div> <div>3%</div> <div>61%</div> <div>14%</div> <div>•</div> <div>25%</div> </div>
2	D	15	<div> <div>13%</div> <div>13%</div> <div>73%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4556 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA repair protein RAD2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	S	Se	0	1	0
			2291	1480	373	426	2	10			
1	B	277	Total	C	N	O	S	Se	0	0	0
			2104	1367	334	391	2	10			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	EXPRESSION TAG	UNP P07276
A	1	MSE	-	EXPRESSION TAG	UNP P07276
B	0	SER	-	EXPRESSION TAG	UNP P07276
B	1	MSE	-	EXPRESSION TAG	UNP P07276

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*TP*GP*AP*GP*TP*CP*AP*GP*AP*GP*CP*AP*AP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	4	Total	C	N	O	P	0	0	0
			82	39	15	24	4			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	59	Total	O	0	0
			59	59		
3	B	20	Total	O	0	0
			20	20		



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	94.08 Å 94.08 Å 155.26 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.41 – 2.75 39.41 – 2.75	Depositor EDS
% Data completeness (in resolution range)	97.8 (39.41-2.75) 98.8 (39.41-2.75)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.05 (at 2.77 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.245 , 0.310 0.267 , 0.320	Depositor DCC
R_{free} test set	1087 reflections (5.45%)	DCC
Wilson B-factor (Å ²)	58.4	Xtriage
Anisotropy	0.418	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 36.1	EDS
Estimated twinning fraction	0.028 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 21057 reflections	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	4556	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.45	0/2335	0.58	0/3145
1	B	0.45	0/2140	0.57	0/2890
2	D	1.11	0/91	0.93	0/138
All	All	0.48	0/4566	0.59	0/6173

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2291	0	2165	46	0
1	B	2104	0	1966	27	0
2	D	82	0	46	2	0
3	A	59	0	0	5	0
3	B	20	0	0	0	0
All	All	4556	0	4177	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:8:DT:H2"	2:D:9:DC:H5"	1.64	0.78
1:A:876:SER:OG	2:D:8:DT:OP1	2.08	0.66
1:B:777:GLU:HG3	1:B:781:ARG:HE	1.62	0.64
1:B:884:PHE:O	1:B:886:ASN:N	2.31	0.64
1:A:957:LEU:HD23	1:A:957:LEU:H	1.65	0.62
1:B:3:VAL:HA	1:B:862:GLY:O	2.02	0.60
1:A:963:PHE:HD1	1:A:964:MSE:HE3	1.68	0.57
1:A:26:ARG:NH1	1:A:804:ASN:O	2.31	0.57
1:A:963:PHE:CD1	1:A:964:MSE:HE3	2.40	0.56
1:A:5:SER:HB3	1:A:874:PRO:HG3	1.88	0.56
1:B:863:SER:HB2	1:B:865:TYR:H	1.73	0.54
1:B:19:LEU:HD21	1:B:835:VAL:HG13	1.89	0.54
1:B:818:LEU:HD13	1:B:938:TYR:CE2	2.43	0.53
1:A:59:PHE:CZ	1:A:979:LEU:HD21	2.44	0.53
1:A:956:ASP:CG	1:A:959:MSE:HE3	2.28	0.53
1:A:926:ASP:HB2	3:A:1042:HOH:O	2.08	0.52
1:B:889:ASN:OD1	1:B:889:ASN:N	2.41	0.52
1:A:3:VAL:HA	1:A:862:GLY:O	2.10	0.51
1:B:965:LYS:HA	1:B:970:TRP:H	1.76	0.51
1:A:826:LYS:O	1:A:835:VAL:HG22	2.11	0.51
1:A:871:GLY:O	1:A:917:LYS:HD3	2.11	0.51
1:B:913:ASP:OD1	1:B:913:ASP:N	2.44	0.51
1:A:781:ARG:HG2	1:A:959:MSE:HB3	1.93	0.50
1:A:886:ASN:ND2	1:A:889:ASN:OD1	2.44	0.50
1:A:14:ALA:HB2	1:A:838:TYR:CZ	2.47	0.50
1:A:926:ASP:HB3	1:A:928:ASP:H	1.77	0.49
1:B:980:ILE:O	1:B:982:LEU:N	2.40	0.49
1:A:802:GLN:HG3	3:A:1006:HOH:O	2.12	0.49
1:B:14:ALA:HB1	1:B:836:GLU:HG2	1.95	0.49
1:A:908:ASN:HB2	1:A:911:GLU:OE2	2.13	0.49
1:A:769:MSE:O	1:A:772:ILE:HG22	2.13	0.49
1:A:956:ASP:OD2	1:A:959:MSE:HE3	2.13	0.49
1:A:856:GLU:HB3	1:A:887:LEU:HG	1.94	0.48
1:B:891:LYS:HG3	1:B:925:LEU:HB3	1.95	0.48
1:A:851:ARG:HA	1:A:854:MSE:HE3	1.96	0.48
1:A:776:GLN:HB3	1:A:786:TYR:CZ	2.49	0.47
1:A:801:LEU:HD22	1:A:809:ILE:HG13	1.96	0.47
1:A:815:ASP:OD2	1:A:864:ASP:HB2	2.16	0.46
1:A:80:VAL:HG13	1:A:84:LYS:HD3	1.98	0.46
1:B:70:ILE:O	1:B:72:PRO:HD3	2.16	0.46
1:A:34:TRP:O	1:A:37:GLN:HG3	2.16	0.46
1:A:853:ASN:ND2	3:A:1026:HOH:O	2.49	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:962:SER:O	1:B:966:THR:HG23	2.16	0.45
1:A:3:VAL:HA	1:A:863:SER:HB3	1.97	0.45
1:B:30:ASP:HB2	1:B:794:GLU:OE2	2.16	0.45
1:B:3:VAL:HB	1:B:6:PHE:HB2	1.99	0.45
1:B:6:PHE:HD2	1:B:814:SER:HB2	1.82	0.45
1:A:872:MSE:HE1	1:A:914:LEU:HD11	1.99	0.44
1:B:5:SER:HB3	1:B:874:PRO:HG3	1.99	0.44
1:B:859:GLN:HE21	1:B:934:VAL:HB	1.82	0.44
1:A:947:THR:O	1:A:947:THR:OG1	2.36	0.43
1:A:947:THR:O	1:A:949:PRO:HD3	2.18	0.43
1:A:35:ILE:CD1	1:A:772:ILE:HG13	2.49	0.43
1:A:6:PHE:HD2	1:A:814:SER:HB2	1.84	0.43
1:A:935:TYR:CZ	1:A:939:MSE:HG3	2.54	0.43
1:A:54:HIS:HD2	1:A:771:MSE:CB	2.32	0.42
1:A:59:PHE:HZ	1:A:979:LEU:HD21	1.82	0.42
3:A:1031:HOH:O	1:B:18:ARG:HG2	2.19	0.42
1:A:82:VAL:O	1:A:86:GLU:HG3	2.19	0.42
1:A:2:GLY:HA3	1:A:814:SER:OG	2.19	0.42
1:B:968:LEU:HD23	1:B:970:TRP:NE1	2.35	0.41
1:A:62:ILE:HD11	1:A:779:LEU:HD23	2.02	0.41
1:B:3:VAL:HB	1:B:6:PHE:CB	2.50	0.41
1:A:4:HIS:O	1:A:861:LEU:HB3	2.21	0.41
1:A:836:GLU:OE2	1:B:832:LYS:HE3	2.20	0.41
1:B:894:TYR:HD2	1:B:925:LEU:HD22	1.84	0.41
1:A:852:LYS:HD3	3:A:1049:HOH:O	2.21	0.41
1:B:968:LEU:HD23	1:B:970:TRP:HE1	1.86	0.40
1:A:5:SER:O	1:A:8:ASP:HB2	2.21	0.40
1:B:894:TYR:CD2	1:B:925:LEU:HD22	2.56	0.40
1:B:873:GLY:H	1:B:876:SER:HB2	1.85	0.40
1:A:957:LEU:CD2	1:A:957:LEU:H	2.30	0.40
1:A:890:PHE:HE1	1:A:914:LEU:HD21	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/367 (78%)	269 (93%)	18 (6%)	1 (0%)	46	77
1	B	269/367 (73%)	249 (93%)	16 (6%)	4 (2%)	13	36
All	All	557/734 (76%)	518 (93%)	34 (6%)	5 (1%)	21	52

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	11	GLY
1	B	885	GLY
1	B	926	ASP
1	B	81	PRO
1	B	981	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/313 (74%)	213 (92%)	19 (8%)	14	35
1	B	210/313 (67%)	197 (94%)	13 (6%)	23	51
All	All	442/626 (71%)	410 (93%)	32 (7%)	17	42

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MSE
1	A	4	HIS
1	A	24	ASP
1	A	54	HIS
1	A	56	THR
1	A	82	VAL
1	A	775	VAL
1	A	777	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	788	THR
1	A	792	GLU
1	A	804	ASN
1	A	832	LYS
1	A	835	VAL
1	A	857	LEU
1	A	926	ASP
1	A	943	VAL
1	A	957	LEU
1	A	958	ASP
1	A	964	MSE
1	B	21	SER
1	B	32	SER
1	B	56	THR
1	B	87	THR
1	B	768	THR
1	B	782	PHE
1	B	832	LYS
1	B	866	THR
1	B	875	VAL
1	B	889	ASN
1	B	913	ASP
1	B	933	MSE
1	B	958	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	281/367 (76%)	0.32	8 (2%) 56 50	40, 48, 74, 79	1 (0%)
1	B	265/367 (72%)	0.39	10 (3%) 44 37	42, 55, 67, 70	0
2	D	4/15 (26%)	2.22	2 (50%) 0 0	58, 59, 62, 67	0
All	All	550/749 (73%)	0.37	20 (3%) 46 40	40, 51, 68, 79	1 (0%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	36	TYR	4.0
1	B	865	TYR	4.0
1	B	3	VAL	3.9
2	D	10	DA	3.7
1	A	3	VAL	3.4
1	A	4	HIS	3.3
1	B	895	ASN	3.2
1	A	88	ILE	3.1
1	B	918	LEU	3.1
1	B	963	PHE	2.8
1	A	972	HIS	2.8
1	A	2	GLY	2.8
1	B	58	PHE	2.6
1	B	4	HIS	2.6
1	B	87	THR	2.4
1	A	912	LYS	2.2
1	A	917	LYS	2.2
2	D	7	DG	2.2
1	B	910	PHE	2.2
1	B	968	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.