



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:24 AM GMT

PDB ID : 2Q4E
Title : Ensemble refinement of the protein crystal structure of gene product from Arabidopsis thaliana At4g09670
Authors : Levin, E.J.; Kondrashov, D.A.; Wesenberg, G.E.; Phillips Jr., G.N.; Center for Eukaryotic Structural Genomics (CESG)
Deposited on : 2007-05-31
Resolution : 2.49 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1-A	362	<div> <div>22%</div> <div>87% 7% 5%</div> </div>
1	1-B	362	<div> <div>15%</div> <div>90% 6% .</div> </div>
1	10-A	362	<div> <div>22%</div> <div>88% 7% 5%</div> </div>
1	10-B	362	<div> <div>15%</div> <div>91% 6% .</div> </div>
1	11-A	362	<div> <div>22%</div> <div>88% 6% 5%</div> </div>
1	11-B	362	<div> <div>15%</div> <div>89% 7% .</div> </div>
1	12-A	362	<div> <div>22%</div> <div>85% 10% 5%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	12-B	362	
1	13-A	362	
1	13-B	362	
1	14-A	362	
1	14-B	362	
1	15-A	362	
1	15-B	362	
1	16-A	362	
1	16-B	362	
1	2-A	362	
1	2-B	362	
1	3-A	362	
1	3-B	362	
1	4-A	362	
1	4-B	362	
1	5-A	362	
1	5-B	362	
1	6-A	362	
1	6-B	362	
1	7-A	362	
1	7-B	362	
1	8-A	362	
1	8-B	362	
1	9-A	362	
1	9-B	362	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 87408 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable oxidoreductase At4g09670.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	1-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			
1	2-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			
1	3-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			
1	4-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			
1	5-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			
1	6-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			
1	7-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			
1	8-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			
1	9-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			
1	10-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			
1	11-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			
1	12-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			
1	13-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			
1	14-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			
1	15-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			
1	16-A	343	Total	C	N	O	S	Se	0	0	0
			2648	1687	445	504	6	6			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	1-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			
1	2-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			
1	3-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			
1	4-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			
1	5-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			
1	6-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			
1	7-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			
1	8-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			
1	9-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			
1	10-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			
1	11-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			
1	12-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			
1	13-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			
1	14-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			
1	15-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			
1	16-B	350	Total	C	N	O	S	Se	0	0	0
			2692	1712	453	515	6	6			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	EXPRESSION TAG	UNP Q9SZ83
A	12	MSE	MET	MODIFIED RESIDUE	UNP Q9SZ83
A	106	MSE	MET	MODIFIED RESIDUE	UNP Q9SZ83
A	126	MSE	MET	MODIFIED RESIDUE	UNP Q9SZ83
A	130	MSE	MET	MODIFIED RESIDUE	UNP Q9SZ83
A	249	MSE	MET	MODIFIED RESIDUE	UNP Q9SZ83
A	310	MSE	MET	MODIFIED RESIDUE	UNP Q9SZ83

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	SER	-	EXPRESSION TAG	UNP Q9SZ83
B	12	MSE	MET	MODIFIED RESIDUE	UNP Q9SZ83
B	106	MSE	MET	MODIFIED RESIDUE	UNP Q9SZ83
B	126	MSE	MET	MODIFIED RESIDUE	UNP Q9SZ83
B	130	MSE	MET	MODIFIED RESIDUE	UNP Q9SZ83
B	249	MSE	MET	MODIFIED RESIDUE	UNP Q9SZ83
B	310	MSE	MET	MODIFIED RESIDUE	UNP Q9SZ83

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	1-A	60	Total O 60 60	0	0
2	2-A	58	Total O 58 58	0	0
2	3-A	62	Total O 62 62	0	0
2	4-A	60	Total O 60 60	0	0
2	5-A	59	Total O 59 59	0	0
2	6-A	60	Total O 60 60	0	0
2	7-A	61	Total O 61 61	0	0
2	8-A	56	Total O 56 56	0	0
2	9-A	60	Total O 60 60	0	0
2	10-A	59	Total O 59 59	0	0
2	11-A	60	Total O 60 60	0	0
2	12-A	61	Total O 61 61	0	0
2	13-A	62	Total O 62 62	0	0
2	14-A	59	Total O 59 59	0	0
2	15-A	59	Total O 59 59	0	0
2	16-A	60	Total O 60 60	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	1-B	63	Total 63	O 63	0	0
2	2-B	65	Total 65	O 65	0	0
2	3-B	61	Total 61	O 61	0	0
2	4-B	63	Total 63	O 63	0	0
2	5-B	64	Total 64	O 64	0	0
2	6-B	63	Total 63	O 63	0	0
2	7-B	62	Total 62	O 62	0	0
2	8-B	67	Total 67	O 67	0	0
2	9-B	63	Total 63	O 63	0	0
2	10-B	64	Total 64	O 64	0	0
2	11-B	63	Total 63	O 63	0	0
2	12-B	62	Total 62	O 62	0	0
2	13-B	61	Total 61	O 61	0	0
2	14-B	64	Total 64	O 64	0	0
2	15-B	64	Total 64	O 64	0	0
2	16-B	63	Total 63	O 63	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Probable oxidoreductase At4g09670



- Molecule 1: Probable oxidoreductase At4g09670




- Molecule 1: Probable oxidoreductase At4g09670









- Molecule 1: Probable oxidoreductase At4g09670





- Molecule 1: Probable oxidoreductase At4g09670




- Molecule 1: Probable oxidoreductase At4g09670




- Molecule 1: Probable oxidoreductase At4g09670






- Molecule 1: Probable oxidoreductase At4g09670

Chain 4-B: 15% 93%




- Molecule 1: Probable oxidoreductase At4g09670

Chain 5-A: 22% 88% 7% 5%




- Molecule 1: Probable oxidoreductase At4g09670

Chain 5-B: 15% 91% 6%




- Molecule 1: Probable oxidoreductase At4g09670

Chain 6-A: 22% 89% 5% 5%




- Molecule 1: Probable oxidoreductase At4g09670

Chain 6-B: 15% 91% 6%




- Molecule 1: Probable oxidoreductase At4g09670


Chain 7-A: 22% 86% 9% 5%




- Molecule 1: Probable oxidoreductase At4g09670

Chain 7-B: 15% 90% 6%







- Molecule 1: Probable oxidoreductase At4g09670




- Molecule 1: Probable oxidoreductase At4g09670




- Molecule 1: Probable oxidoreductase At4g09670




- Molecule 1: Probable oxidoreductase At4g09670




- Molecule 1: Probable oxidoreductase At4g09670




- Molecule 1: Probable oxidoreductase At4g09670




- Molecule 1: Probable oxidoreductase At4g09670







- Molecule 1: Probable oxidoreductase At4g09670




- Molecule 1: Probable oxidoreductase At4g09670




- Molecule 1: Probable oxidoreductase At4g09670





- Molecule 1: Probable oxidoreductase At4g09670









- Molecule 1: Probable oxidoreductase At4g09670




- Molecule 1: Probable oxidoreductase At4g09670



- Molecule 1: Probable oxidoreductase At4g09670





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.20Å 107.66Å 129.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.73 – 2.49 29.73 – 2.49	Depositor EDS
% Data completeness (in resolution range)	98.4 (29.73-2.49) 98.4 (29.73-2.49)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.60 (at 2.48Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.215 , 0.285 0.234 , 0.299	Depositor DCC
R_{free} test set	1472 reflections (5.38%)	DCC
Wilson B-factor (Å ²)	36.7	Xtriage
Anisotropy	0.414	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 92.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 28862 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	87408	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	1-A	0.34	0/2698	0.59	0/3656
1	1-B	0.35	0/2743	0.60	1/3716 (0.0%)
1	2-A	0.33	0/2698	0.57	0/3656
1	2-B	0.35	0/2743	0.60	0/3716
1	3-A	0.33	0/2698	0.59	0/3656
1	3-B	0.34	0/2743	0.58	0/3716
1	4-A	0.34	0/2698	0.59	0/3656
1	4-B	0.34	0/2743	0.60	0/3716
1	5-A	0.34	0/2698	0.59	0/3656
1	5-B	0.34	0/2743	0.58	0/3716
1	6-A	0.34	0/2698	0.59	0/3656
1	6-B	0.35	0/2743	0.60	1/3716 (0.0%)
1	7-A	0.33	0/2698	0.56	0/3656
1	7-B	0.34	0/2743	0.59	0/3716
1	8-A	0.34	0/2698	0.59	0/3656
1	8-B	0.35	0/2743	0.60	0/3716
1	9-A	0.33	0/2698	0.59	0/3656
1	9-B	0.35	0/2743	0.60	0/3716
1	10-A	0.34	0/2698	0.58	0/3656
1	10-B	0.35	0/2743	0.59	0/3716
1	11-A	0.34	0/2698	0.58	0/3656
1	11-B	0.35	0/2743	0.60	0/3716
1	12-A	0.34	0/2698	0.58	0/3656
1	12-B	0.34	0/2743	0.59	0/3716
1	13-A	0.36	0/2698	0.62	0/3656
1	13-B	0.37	0/2743	0.65	1/3716 (0.0%)
1	14-A	0.36	0/2698	0.62	0/3656
1	14-B	0.38	0/2743	0.64	1/3716 (0.0%)
1	15-A	0.37	0/2698	0.63	0/3656
1	15-B	0.38	0/2743	0.66	1/3716 (0.0%)
1	16-A	0.37	0/2698	0.62	0/3656
1	16-B	0.38	0/2743	0.63	1/3716 (0.0%)
All	All	0.35	0/87056	0.60	6/117952 (0.0%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	15-B	27	LEU	N-CA-C	-6.95	92.23	111.00
1	16-B	27	LEU	N-CA-C	-6.42	93.67	111.00
1	13-B	27	LEU	N-CA-C	-5.91	95.04	111.00
1	14-B	27	LEU	N-CA-C	-5.71	95.58	111.00
1	6-B	27	LEU	N-CA-C	-5.42	96.37	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1-A	2648	0	2642	0	0
1	1-B	2692	0	2680	0	0
1	2-A	2648	0	2642	0	0
1	2-B	2692	0	2680	0	0
1	3-A	2648	0	2642	0	0
1	3-B	2692	0	2680	0	0
1	4-A	2648	0	2642	0	0
1	4-B	2692	0	2680	0	0
1	5-A	2648	0	2642	0	0
1	5-B	2692	0	2680	0	0
1	6-A	2648	0	2642	0	0
1	6-B	2692	0	2680	0	0
1	7-A	2648	0	2642	0	0
1	7-B	2692	0	2680	0	0
1	8-A	2648	0	2642	0	0
1	8-B	2692	0	2680	0	0
1	9-A	2648	0	2642	0	0
1	9-B	2692	0	2680	0	0
1	10-A	2648	0	2642	0	0
1	10-B	2692	0	2680	0	0
1	11-A	2648	0	2642	0	0
1	11-B	2692	0	2680	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	12-A	2648	0	2642	0	0
1	12-B	2692	0	2680	0	0
1	13-A	2648	0	2642	0	0
1	13-B	2692	0	2680	0	0
1	14-A	2648	0	2642	0	0
1	14-B	2692	0	2680	0	0
1	15-A	2648	0	2642	0	0
1	15-B	2692	0	2680	0	0
1	16-A	2648	0	2642	0	0
1	16-B	2692	0	2680	0	0
2	1-A	60	0	0	0	0
2	1-B	63	0	0	0	0
2	2-A	58	0	0	0	0
2	2-B	65	0	0	0	0
2	3-A	62	0	0	0	0
2	3-B	61	0	0	0	0
2	4-A	60	0	0	0	0
2	4-B	63	0	0	0	0
2	5-A	59	0	0	0	0
2	5-B	64	0	0	0	0
2	6-A	60	0	0	0	0
2	6-B	63	0	0	0	0
2	7-A	61	0	0	0	0
2	7-B	62	0	0	0	0
2	8-A	56	0	0	0	0
2	8-B	67	0	0	0	0
2	9-A	60	0	0	0	0
2	9-B	63	0	0	0	0
2	10-A	59	0	0	0	0
2	10-B	64	0	0	0	0
2	11-A	60	0	0	0	0
2	11-B	63	0	0	0	0
2	12-A	61	0	0	0	0
2	12-B	62	0	0	0	0
2	13-A	62	0	0	0	0
2	13-B	61	0	0	0	0
2	14-A	59	0	0	0	0
2	14-B	64	0	0	0	0
2	15-A	59	0	0	0	0
2	15-B	64	0	0	0	0
2	16-A	60	0	0	0	0
2	16-B	63	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	87408	0	85152	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). Clashscore could not be calculated for this entry.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1-A	339/362 (94%)	294 (87%)	37 (11%)	8 (2%)	7	11
1	1-B	346/362 (96%)	300 (87%)	36 (10%)	10 (3%)	6	8
1	2-A	339/362 (94%)	300 (88%)	34 (10%)	5 (2%)	13	22
1	2-B	346/362 (96%)	307 (89%)	32 (9%)	7 (2%)	9	15
1	3-A	339/362 (94%)	276 (81%)	49 (14%)	14 (4%)	3	4
1	3-B	346/362 (96%)	286 (83%)	52 (15%)	8 (2%)	8	12
1	4-A	339/362 (94%)	287 (85%)	37 (11%)	15 (4%)	3	3
1	4-B	346/362 (96%)	315 (91%)	26 (8%)	5 (1%)	14	24
1	5-A	339/362 (94%)	295 (87%)	37 (11%)	7 (2%)	9	14
1	5-B	346/362 (96%)	298 (86%)	37 (11%)	11 (3%)	5	6
1	6-A	339/362 (94%)	298 (88%)	35 (10%)	6 (2%)	11	18
1	6-B	346/362 (96%)	309 (89%)	29 (8%)	8 (2%)	8	12
1	7-A	339/362 (94%)	291 (86%)	37 (11%)	11 (3%)	5	6
1	7-B	346/362 (96%)	294 (85%)	40 (12%)	12 (4%)	4	6
1	8-A	339/362 (94%)	280 (83%)	48 (14%)	11 (3%)	5	6
1	8-B	346/362 (96%)	303 (88%)	37 (11%)	6 (2%)	11	19

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	9-A	339/362 (94%)	299 (88%)	37 (11%)	3 (1%)	21	37
1	9-B	346/362 (96%)	309 (89%)	32 (9%)	5 (1%)	14	24
1	10-A	339/362 (94%)	294 (87%)	37 (11%)	8 (2%)	7	11
1	10-B	346/362 (96%)	296 (86%)	40 (12%)	10 (3%)	6	8
1	11-A	339/362 (94%)	284 (84%)	44 (13%)	11 (3%)	5	6
1	11-B	346/362 (96%)	288 (83%)	43 (12%)	15 (4%)	3	4
1	12-A	339/362 (94%)	272 (80%)	49 (14%)	18 (5%)	2	2
1	12-B	346/362 (96%)	294 (85%)	46 (13%)	6 (2%)	11	19
1	13-A	339/362 (94%)	281 (83%)	43 (13%)	15 (4%)	3	3
1	13-B	346/362 (96%)	309 (89%)	27 (8%)	10 (3%)	6	8
1	14-A	339/362 (94%)	288 (85%)	36 (11%)	15 (4%)	3	3
1	14-B	346/362 (96%)	302 (87%)	33 (10%)	11 (3%)	5	6
1	15-A	339/362 (94%)	265 (78%)	53 (16%)	21 (6%)	2	1
1	15-B	346/362 (96%)	282 (82%)	48 (14%)	16 (5%)	3	3
1	16-A	339/362 (94%)	270 (80%)	50 (15%)	19 (6%)	2	2
1	16-B	346/362 (96%)	291 (84%)	42 (12%)	13 (4%)	4	5
All	All	10960/11584 (95%)	9357 (85%)	1263 (12%)	340 (3%)	5	7

5 of 340 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	1-A	140	LEU
1	1-A	317	LEU
1	1-B	28	ALA
1	1-B	179	LEU
1	1-B	246	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1-A	286/293 (98%)	266 (93%)	20 (7%)	19	34
1	1-B	290/293 (99%)	276 (95%)	14 (5%)	31	55
1	2-A	286/293 (98%)	269 (94%)	17 (6%)	24	44
1	2-B	290/293 (99%)	280 (97%)	10 (3%)	44	72
1	3-A	286/293 (98%)	263 (92%)	23 (8%)	15	28
1	3-B	290/293 (99%)	275 (95%)	15 (5%)	29	51
1	4-A	286/293 (98%)	267 (93%)	19 (7%)	21	38
1	4-B	290/293 (99%)	280 (97%)	10 (3%)	44	72
1	5-A	286/293 (98%)	268 (94%)	18 (6%)	22	40
1	5-B	290/293 (99%)	278 (96%)	12 (4%)	37	63
1	6-A	286/293 (98%)	270 (94%)	16 (6%)	26	47
1	6-B	290/293 (99%)	277 (96%)	13 (4%)	34	59
1	7-A	286/293 (98%)	264 (92%)	22 (8%)	16	30
1	7-B	290/293 (99%)	276 (95%)	14 (5%)	31	55
1	8-A	286/293 (98%)	273 (96%)	13 (4%)	34	59
1	8-B	290/293 (99%)	272 (94%)	18 (6%)	23	41
1	9-A	286/293 (98%)	265 (93%)	21 (7%)	17	32
1	9-B	290/293 (99%)	276 (95%)	14 (5%)	31	55
1	10-A	286/293 (98%)	268 (94%)	18 (6%)	22	40
1	10-B	290/293 (99%)	279 (96%)	11 (4%)	40	67
1	11-A	286/293 (98%)	273 (96%)	13 (4%)	34	59
1	11-B	290/293 (99%)	276 (95%)	14 (5%)	31	55
1	12-A	286/293 (98%)	267 (93%)	19 (7%)	21	38
1	12-B	290/293 (99%)	272 (94%)	18 (6%)	23	41
1	13-A	286/293 (98%)	262 (92%)	24 (8%)	14	25
1	13-B	290/293 (99%)	275 (95%)	15 (5%)	29	51
1	14-A	286/293 (98%)	262 (92%)	24 (8%)	14	25
1	14-B	290/293 (99%)	270 (93%)	20 (7%)	19	35
1	15-A	286/293 (98%)	261 (91%)	25 (9%)	13	24
1	15-B	290/293 (99%)	264 (91%)	26 (9%)	12	22
1	16-A	286/293 (98%)	261 (91%)	25 (9%)	13	24
1	16-B	290/293 (99%)	259 (89%)	31 (11%)	8	15

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	9216/9376 (98%)	8644 (94%)	572 (6%)	23	41

5 of 572 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	9-A	58	LYS
1	11-A	99	LEU
1	16-A	159	CYS
1	9-A	180	ASP
1	10-A	33	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 116 such sidechains are listed below:

Mol	Chain	Res	Type
1	9-A	124	GLN
1	11-A	60	HIS
1	16-A	246	ASN
1	9-A	200	ASN
1	10-A	51	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	1-A	337/362 (93%)	1.21	81 (24%) 1 1	15, 33, 53, 80	337 (100%)
1	1-B	344/362 (95%)	0.94	56 (16%) 2 2	16, 31, 46, 53	344 (100%)
1	2-A	337/362 (93%)	1.21	81 (24%) 1 1	15, 33, 53, 80	337 (100%)
1	2-B	344/362 (95%)	0.94	56 (16%) 2 2	16, 31, 46, 53	344 (100%)
1	3-A	337/362 (93%)	1.21	81 (24%) 1 1	15, 33, 53, 80	337 (100%)
1	3-B	344/362 (95%)	0.94	56 (16%) 2 2	16, 31, 46, 53	344 (100%)
1	4-A	337/362 (93%)	1.21	81 (24%) 1 1	15, 33, 53, 80	337 (100%)
1	4-B	344/362 (95%)	0.94	56 (16%) 2 2	16, 31, 46, 53	344 (100%)
1	5-A	337/362 (93%)	1.21	81 (24%) 1 1	15, 33, 53, 80	337 (100%)
1	5-B	344/362 (95%)	0.94	56 (16%) 2 2	16, 31, 46, 53	344 (100%)
1	6-A	337/362 (93%)	1.21	81 (24%) 1 1	15, 33, 53, 80	337 (100%)
1	6-B	344/362 (95%)	0.94	56 (16%) 2 2	16, 31, 46, 53	344 (100%)
1	7-A	337/362 (93%)	1.21	81 (24%) 1 1	15, 33, 53, 80	337 (100%)
1	7-B	344/362 (95%)	0.94	56 (16%) 2 2	16, 31, 46, 53	344 (100%)
1	8-A	337/362 (93%)	1.21	81 (24%) 1 1	15, 33, 53, 80	337 (100%)
1	8-B	344/362 (95%)	0.94	56 (16%) 2 2	16, 31, 46, 53	344 (100%)
1	9-A	337/362 (93%)	1.21	81 (24%) 1 1	15, 33, 53, 80	337 (100%)
1	9-B	344/362 (95%)	0.94	56 (16%) 2 2	16, 31, 46, 53	344 (100%)
1	10-A	337/362 (93%)	1.21	81 (24%) 1 1	15, 33, 53, 80	337 (100%)
1	10-B	344/362 (95%)	0.94	56 (16%) 2 2	16, 31, 46, 53	344 (100%)
1	11-A	337/362 (93%)	1.21	81 (24%) 1 1	15, 33, 53, 80	337 (100%)
1	11-B	344/362 (95%)	0.94	56 (16%) 2 2	16, 31, 46, 53	344 (100%)
1	12-A	337/362 (93%)	1.21	81 (24%) 1 1	15, 33, 53, 80	337 (100%)
1	12-B	344/362 (95%)	0.94	56 (16%) 2 2	16, 31, 46, 53	344 (100%)

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	13-A	337/362 (93%)	1.21	81 (24%)	1	1	15, 33, 53, 80	337 (100%)
1	13-B	344/362 (95%)	0.94	56 (16%)	2	2	16, 31, 46, 53	344 (100%)
1	14-A	337/362 (93%)	1.21	81 (24%)	1	1	15, 33, 53, 80	337 (100%)
1	14-B	344/362 (95%)	0.94	56 (16%)	2	2	16, 31, 46, 53	344 (100%)
1	15-A	337/362 (93%)	1.21	81 (24%)	1	1	15, 33, 53, 80	337 (100%)
1	15-B	344/362 (95%)	0.94	56 (16%)	2	2	16, 31, 46, 53	344 (100%)
1	16-A	337/362 (93%)	1.21	81 (24%)	1	1	15, 33, 53, 80	337 (100%)
1	16-B	344/362 (95%)	0.94	56 (16%)	2	2	16, 31, 46, 53	344 (100%)
All	All	10896/11584 (94%)	1.07	2192 (20%)	2	1	15, 32, 51, 80	10896 (100%)

The worst 5 of 2192 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	1-A	317	LEU	9.8
1	2-A	317	LEU	9.8
1	3-A	317	LEU	9.8
1	4-A	317	LEU	9.8
1	5-A	317	LEU	9.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.