



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:51 PM GMT

PDB ID : 4Q44  
Title : Polymerase-Damaged DNA Complex  
Authors : Nair, D.T.; Kottur, J.; Sharma, A.  
Deposited on : 2014-04-13  
Resolution : 2.71 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2439 (2.74-2.70)
Clashscore	102246	2771 (2.74-2.70)
Ramachandran outliers	100387	2726 (2.74-2.70)
Sidechain outliers	100360	2727 (2.74-2.70)
RSRZ outliers	91569	2443 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 22%, green 75%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>75%</span> <span>22%</span> </div> </div>
1	F	342	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, yellow 24%, green 73%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>73%</span> <span>24%</span> </div> </div>
2	B	18	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, yellow 50%, green 50%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>50%</span> <span>50%</span> </div> </div>
2	C	18	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 6%, orange 1%, yellow 17%, green 78%, grey 6%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>78%</span> <span>17%</span> </div> </div>
2	G	18	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 6%, orange 1%, yellow 22%, green 78%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>78%</span> <span>22%</span> </div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	H	18	 67% 11% 22%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6904 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase IV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	342	Total	C	N	O	S	0	0	0
			2684	1692	493	485	14			
1	F	342	Total	C	N	O	S	0	0	0
			2688	1695	494	485	14			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	EXPRESSION TAG	UNP Q47155
A	1	SER	-	EXPRESSION TAG	UNP Q47155
A	64	ALA	LYS	CONFLICT	UNP Q47155
A	205	ALA	LYS	CONFLICT	UNP Q47155
F	0	GLY	-	EXPRESSION TAG	UNP Q47155
F	1	SER	-	EXPRESSION TAG	UNP Q47155
F	64	ALA	LYS	CONFLICT	UNP Q47155
F	205	ALA	LYS	CONFLICT	UNP Q47155

- Molecule 2 is a DNA chain called DNA (5'-D(\*TP\*CP\*TP\*AP\*(RDG)P\*GP\*GP\*TP\*CP\*CP\*TP\*AP\*GP\*GP\*AP\*CP\*CP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	18	Total	C	N	O	P	0	0	0
			370	179	66	108	17			
2	C	17	Total	C	N	O	P	0	0	0
			350	169	64	101	16			
2	G	18	Total	C	N	O	P	0	0	0
			370	179	66	108	17			
2	H	14	Total	C	N	O	P	0	0	0
			293	140	54	85	14			

- Molecule 3 is 5'-O-[(R)-HYDROXY{[(R)-HYDROXY(PHOSPHONOOXY)PHOSPHORYL]AMINO}PHOSPHORYL]THYMIDINE (three-letter code: 1FZ) (formula:



- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

- Molecule 5 is water.

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	8	Total	O	0	0
			8	8		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: DNA polymerase IV




#### • Molecule 1: DNA polymerase IV




#### • Molecule 2: DNA (5'-D(\*TP\*CP\*TP\*AP\*(RDG)P\*GP\*GP\*TP\*CP\*CP\*TP\*AP\*GP\*GP\*A P\*CP\*CP\*C)-3')





#### • Molecule 2: DNA (5'-D(\*TP\*CP\*TP\*AP\*(RDG)P\*GP\*GP\*TP\*CP\*CP\*TP\*AP\*GP\*GP\*A P\*CP\*CP\*C)-3')





- Molecule 2: DNA (5'-D(\*TP\*CP\*TP\*AP\*(RDG)P\*GP\*GP\*TP\*CP\*CP\*TP\*AP\*GP\*GP\*A P\*CP\*CP\*C)-3')



- Molecule 2: DNA (5'-D(\*TP\*CP\*TP\*AP\*(RDG)P\*GP\*GP\*TP\*CP\*CP\*TP\*AP\*GP\*GP\*A P\*CP\*CP\*C)-3')





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.66Å 56.93Å 110.40Å 90.00° 94.66° 90.00°	Depositor
Resolution (Å)	48.21 – 2.71 48.21 – 2.71	Depositor EDS
% Data completeness (in resolution range)	99.6 (48.21-2.71) 99.6 (48.21-2.71)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.54 (at 2.69Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.8_1069)	Depositor
R, $R_{free}$	0.225 , 0.270 0.233 , 0.277	Depositor DCC
$R_{free}$ test set	1495 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	47.7	Xtriage
Anisotropy	0.291	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 29.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 29597 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6904	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.44% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: RDG, MG, 1FZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.24	0/2735	0.47	1/3695 (0.0%)
1	F	0.23	0/2739	0.44	0/3699
2	B	0.41	0/381	0.99	0/583
2	C	0.42	0/359	1.05	0/549
2	G	0.41	0/381	1.04	0/583
2	H	0.40	0/296	0.96	0/454
All	All	0.28	0/6891	0.63	1/9563 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	282	LEU	CB-CG-CD1	-5.63	101.42	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	F	340	ASP	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2684	0	2734	63	0
1	F	2688	0	2745	51	0
2	B	370	0	208	7	0
2	C	350	0	196	4	0
2	G	370	0	208	5	0
2	H	293	0	161	2	0
3	A	29	0	14	3	0
3	F	29	0	14	2	0
4	A	2	0	0	0	0
4	F	2	0	0	0	0
5	A	18	0	0	5	0
5	B	4	0	0	0	0
5	C	6	0	0	0	0
5	F	40	0	0	9	0
5	G	11	0	0	2	0
5	H	8	0	0	1	0
All	All	6904	0	6280	126	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 126 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:235:ASN:O	5:F:520:HOH:O	1.81	0.99
2:H:863:DT:O4	5:H:907:HOH:O	1.83	0.97
3:F:401:1FZ:O1G	5:F:524:HOH:O	1.90	0.90
1:A:3:LYS:H	1:A:110:THR:HG22	1.44	0.82
1:A:154:ASP:OD2	5:A:504:HOH:O	1.98	0.79

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/342 (99%)	316 (93%)	18 (5%)	6 (2%)	11	25
1	F	340/342 (99%)	318 (94%)	18 (5%)	4 (1%)	16	38
All	All	680/684 (99%)	634 (93%)	36 (5%)	10 (2%)	13	31

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	251	GLU
1	A	282	LEU
1	F	340	ASP
1	A	24	LEU
1	A	253	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/286 (100%)	264 (93%)	21 (7%)	17	38
1	F	286/286 (100%)	267 (93%)	19 (7%)	21	44
All	All	571/572 (100%)	531 (93%)	40 (7%)	19	41

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	296	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	F	42	SER
1	F	295	PHE
1	A	340	ASP
1	F	65	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	286	GLN
1	F	300	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	RDG	B	841	2	21,31,32	2.84	8 (38%)	26,44,47	2.12	10 (38%)
2	RDG	C	860	2	21,31,32	2.84	9 (42%)	26,44,47	2.09	10 (38%)
2	RDG	G	841	2	21,31,32	2.83	9 (42%)	26,44,47	2.21	11 (42%)
2	RDG	H	860	2	21,31,32	2.84	8 (38%)	26,44,47	2.14	11 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RDG	B	841	2	-	0/6/26/27	0/3/4/4
2	RDG	C	860	2	-	0/6/26/27	0/3/4/4
2	RDG	G	841	2	-	0/6/26/27	0/3/4/4
2	RDG	H	860	2	-	0/6/26/27	0/3/4/4

The worst 5 of 34 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	841	RDG	O3'-C3'	-3.47	1.35	1.43
2	C	860	RDG	O3'-C3'	-3.46	1.35	1.43
2	H	860	RDG	O3'-C3'	-3.45	1.35	1.43
2	B	841	RDG	O3'-C3'	-3.44	1.35	1.43
2	G	841	RDG	C5'-C4'	-2.87	1.42	1.51

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	841	RDG	C7-C9-C10	-3.88	103.75	106.75
2	C	860	RDG	C7-C9-C10	-3.80	103.81	106.75
2	H	860	RDG	C7-C9-C10	-3.79	103.81	106.75
2	B	841	RDG	C7-C9-C10	-3.77	103.83	106.75
2	G	841	RDG	C11-N2-C2	-3.70	118.18	123.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	841	RDG	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	1FZ	A	401	4	24,30,30	1.58	2 (8%)	31,47,47	1.84	5 (16%)
3	1FZ	F	401	4	24,30,30	1.57	2 (8%)	31,47,47	1.91	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1FZ	A	401	4	-	0/13/34/34	0/2/2/2
3	1FZ	F	401	4	-	0/13/34/34	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	401	1FZ	PB-O2B	4.62	1.51	1.46
3	A	401	1FZ	PB-O2B	4.64	1.51	1.46
3	A	401	1FZ	PA-O1A	4.70	1.51	1.46
3	F	401	1FZ	PA-O1A	4.71	1.51	1.46

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	401	1FZ	C5-C4-N3	-5.47	119.04	125.14
3	A	401	1FZ	C5-C4-N3	-5.34	119.19	125.14
3	F	401	1FZ	PG-O3B-PB	-4.00	119.27	132.67
3	A	401	1FZ	PG-O3B-PB	-3.40	121.26	132.67
3	A	401	1FZ	C2'-C1'-N1	-2.38	108.36	114.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	1FZ	3	0
3	F	401	1FZ	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	342/342 (100%)	0.19	4 (1%) 81 81	49, 64, 81, 101	0
1	F	342/342 (100%)	-0.00	0 100 100	44, 57, 72, 83	0
2	B	17/18 (94%)	-0.17	0 100 100	51, 66, 87, 90	0
2	C	16/18 (88%)	-0.08	1 (6%) 23 23	51, 67, 91, 116	0
2	G	17/18 (94%)	-0.07	1 (5%) 26 25	47, 62, 85, 93	0
2	H	13/18 (72%)	-0.40	0 100 100	49, 59, 73, 78	0
All	All	747/756 (98%)	0.07	6 (0%) 87 88	44, 60, 79, 116	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	0	GLY	6.0
2	C	857	DC	4.7
1	A	252	ASP	3.7
2	G	837	DT	3.3
1	A	1	SER	3.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	RDG	C	860	28/29	0.94	0.18	-	63,68,73,78	0
2	RDG	G	841	28/29	0.94	0.20	-	43,50,87,95	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	RDG	B	841	28/29	0.95	0.18	-	46,52,59,61	0
2	RDG	H	860	28/29	0.86	0.16	-	64,77,95,95	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	MG	A	402	1/1	0.95	0.17	0.75	46,46,46,46	0
4	MG	F	402	1/1	0.64	0.19	0.56	39,39,39,39	0
3	1FZ	F	401	29/29	0.96	0.14	-0.72	39,51,61,63	0
3	1FZ	A	401	29/29	0.97	0.15	-0.96	47,54,63,67	0
4	MG	F	403	1/1	0.81	0.19	-	63,63,63,63	0
4	MG	A	403	1/1	0.93	0.09	-	60,60,60,60	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.