



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:52 PM GMT

PDB ID : 4Q72
Title : Crystal Structure of Bradyrhizobium japonicum Proline Utilization A (PutA) Mutant D779Y
Authors : Tanner, J.J.; Pemberton, T.A.; Luo, M.
Deposited on : 2014-04-23
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1001	<div> <div>5%</div> <div>89%</div> <div>7%</div> <div>.</div> </div>
1	B	1001	<div> <div>7%</div> <div>89%</div> <div>7%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	2004	-	-	-	X
4	SO4	A	2006	-	-	-	X
4	SO4	B	1004	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 14836 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proline dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	971	Total	C	N	O	S	0	0	0
			7217	4562	1295	1337	23			
1	B	969	Total	C	N	O	S	0	0	0
			7169	4533	1286	1327	23			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	EXPRESSION TAG	UNP Q89E26
A	0	HIS	-	EXPRESSION TAG	UNP Q89E26
A	779	TYR	ASP	ENGINEERED MUTATION	UNP Q89E26
B	-1	GLY	-	EXPRESSION TAG	UNP Q89E26
B	0	HIS	-	EXPRESSION TAG	UNP Q89E26
B	779	TYR	ASP	ENGINEERED MUTATION	UNP Q89E26

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	186	Total	O	0	0
			186	186		
5	B	110	Total	O	0	0
			110	110		


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Proline dehydrogenase



• Molecule 1: Proline dehydrogenase





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	167.11Å 195.99Å 108.70Å 90.00° 121.44° 90.00°	Depositor
Resolution (Å)	31.99 – 2.30 31.99 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.6 (31.99-2.30) 98.6 (31.99-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.17 (at 2.29Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.215 , 0.251 0.229 , 0.264	Depositor DCC
R_{free} test set	6545 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	34.1	Xtriage
Anisotropy	0.527	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 130021 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14836	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/7359	0.56	0/10013
1	B	0.40	0/7309	0.54	0/9949
All	All	0.42	0/14668	0.55	0/19962

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7217	0	7155	36	0
1	B	7169	0	7083	35	0
2	A	53	0	31	1	0
2	B	53	0	31	0	0
3	A	18	0	24	0	0
4	A	15	0	0	0	0
4	B	15	0	0	1	0
5	A	186	0	0	1	0
5	B	110	0	0	0	0
All	All	14836	0	14324	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:491:ARG:HD2	1:B:492:PRO:HD2	1.65	0.77
1:A:491:ARG:HD2	1:A:492:PRO:HD2	1.69	0.73
1:B:197:GLU:OE1	1:B:456:ARG:NH1	2.33	0.61
1:A:197:GLU:OE1	1:A:456:ARG:NH1	2.38	0.56
1:B:475:ARG:NH2	4:B:1003:SO4:O2	2.38	0.55
1:A:158:GLN:NE2	5:A:2186:HOH:O	2.29	0.55
1:A:108:GLU:HG3	1:A:921:ARG:HH11	1.72	0.55
1:B:482:LEU:O	1:B:484:ARG:NH1	2.40	0.53
1:A:138:GLY:HA2	1:A:922:ILE:HG23	1.92	0.52
1:B:108:GLU:HG3	1:B:921:ARG:HH11	1.74	0.51
1:A:463:ALA:O	1:A:466:SER:HB3	2.11	0.51
1:A:772:LEU:HD12	1:A:773:PRO:HD2	1.93	0.51
1:A:482:LEU:O	1:A:484:ARG:NH1	2.44	0.51
1:A:192:PHE:CE2	1:A:219:ILE:HG12	2.47	0.50
1:A:308:GLY:HA3	1:A:337:MET:O	2.12	0.50
1:B:168:VAL:HB	1:B:444:VAL:HG22	1.94	0.50
1:A:870:PRO:HA	1:A:888:PRO:O	2.13	0.49
1:A:368:ALA:HB3	1:A:489:ILE:HD11	1.94	0.49
1:B:694:VAL:HG21	1:B:709:LEU:HD13	1.93	0.49
1:B:138:GLY:HA2	1:B:922:ILE:HG23	1.95	0.48
1:B:953:PRO:HB2	1:B:965:LYS:HD3	1.94	0.48
1:B:660:PRO:HB2	1:B:689:ILE:HG21	1.96	0.48
1:A:372:LEU:HD23	1:A:485:PRO:HB2	1.95	0.48
1:B:192:PHE:CE2	1:B:219:ILE:HG12	2.49	0.48
1:A:824:PRO:HG3	1:A:833:PRO:HD2	1.94	0.48
1:A:491:ARG:HE	1:A:493:GLN:HB3	1.79	0.48
1:B:792:CYS:O	1:B:914:LEU:HD23	2.14	0.47
1:B:200:ARG:NH1	1:B:778:ASP:OD2	2.37	0.47
1:A:856:LEU:HD11	1:A:859:ALA:HB2	1.98	0.46
1:A:29:LEU:HB2	1:A:34:GLU:OE2	2.16	0.45
1:A:957:ASN:O	1:A:960:SER:HB2	2.17	0.45
1:A:108:GLU:HG3	1:A:921:ARG:NH1	2.32	0.45
1:B:109:THR:HG23	1:B:769:ALA:HB3	1.98	0.45
1:B:497:HIS:CE1	1:B:499:ARG:HB2	2.52	0.45
1:A:497:HIS:HE1	1:A:499:ARG:HB2	1.82	0.44
1:B:372:LEU:HD23	1:B:485:PRO:HB2	2.00	0.44
1:A:109:THR:HG23	1:A:769:ALA:HB3	1.99	0.44
1:B:368:ALA:HB3	1:B:489:ILE:HD11	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:590:PHE:HD2	1:B:689:ILE:HD11	1.83	0.44
1:A:497:HIS:CE1	1:A:499:ARG:HB2	2.52	0.44
1:B:451:LEU:HB3	1:B:772:LEU:HD13	2.00	0.44
1:A:238:LEU:HD12	1:A:288:SER:HB2	1.99	0.44
1:B:856:LEU:HD11	1:B:859:ALA:HB2	1.99	0.44
1:B:285:LEU:HD23	1:B:317:ARG:CZ	2.47	0.44
1:B:854:ALA:HB1	1:B:874:GLU:O	2.17	0.44
1:B:396:ASN:HA	1:B:467:PHE:CD2	2.53	0.43
1:B:317:ARG:O	1:B:321:VAL:HG23	2.19	0.43
1:A:170:GLY:O	1:A:446:SER:HA	2.18	0.43
1:A:436:ILE:HD13	1:A:436:ILE:HA	1.82	0.43
1:A:754:PRO:HB3	1:A:974:ARG:NH2	2.34	0.42
1:B:143:LEU:HA	1:B:143:LEU:HD12	1.90	0.42
1:A:660:PRO:HB2	1:A:689:ILE:HG21	2.01	0.42
1:A:193:ASP:O	1:A:443:PRO:HA	2.20	0.42
1:A:104:PHE:CE2	1:A:136:PRO:HA	2.54	0.42
1:B:87:ASP:HB3	1:B:90:THR:OG1	2.20	0.42
1:B:193:ASP:O	1:B:443:PRO:HA	2.20	0.41
1:A:811:MET:HE2	1:A:811:MET:HB2	1.90	0.41
1:A:896:ARG:HA	1:A:897:PRO:HD3	1.88	0.41
1:B:584:GLU:OE2	1:B:617:ARG:NH2	2.44	0.41
1:B:793:SER:HB2	1:B:915:THR:HG21	2.02	0.41
1:A:285:LEU:HD23	1:A:317:ARG:CZ	2.50	0.41
1:A:317:ARG:O	1:A:321:VAL:HG23	2.21	0.41
1:B:104:PHE:CE2	1:B:136:PRO:HA	2.56	0.41
1:B:904:LEU:HD13	1:B:932:ARG:HG2	2.02	0.41
1:B:181:LYS:HA	1:B:182:PRO:HD3	1.93	0.41
1:B:298:ASP:HA	1:B:299:PRO:HD3	1.89	0.41
2:A:2001:FAD:H4'	2:A:2001:FAD:H1'1	1.91	0.41
1:A:953:PRO:HB2	1:A:965:LYS:HD3	2.03	0.40
1:B:308:GLY:HA3	1:B:337:MET:O	2.20	0.40
1:A:110:LYS:HB3	1:A:121:TRP:HE3	1.86	0.40
1:A:904:LEU:HD13	1:A:932:ARG:HG2	2.02	0.40
1:B:146:ARG:HH21	1:B:146:ARG:HD3	1.76	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	961/1001 (96%)	940 (98%)	21 (2%)	0	100	100
1	B	959/1001 (96%)	941 (98%)	18 (2%)	0	100	100
All	All	1920/2002 (96%)	1881 (98%)	39 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	706/758 (93%)	690 (98%)	16 (2%)	58	75
1	B	695/758 (92%)	683 (98%)	12 (2%)	68	83
All	All	1401/1516 (92%)	1373 (98%)	28 (2%)	63	79

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	ASP
1	A	92	ASP
1	A	119	SER
1	A	186	GLN
1	A	235	SER
1	A	255	VAL
1	A	314	TYR
1	A	372	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	522	ARG
1	A	651	PHE
1	A	786	ARG
1	A	811	MET
1	A	853	GLU
1	A	866	CYS
1	A	883	GLU
1	A	957	ASN
1	B	16	ASP
1	B	92	ASP
1	B	119	SER
1	B	235	SER
1	B	314	TYR
1	B	372	LEU
1	B	786	ARG
1	B	811	MET
1	B	853	GLU
1	B	866	CYS
1	B	883	GLU
1	B	957	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	A	2001	-	48,58,58	1.87	14 (29%)	54,89,89	2.42	11 (20%)
3	GOL	A	2002	-	5,5,5	0.40	0	5,5,5	0.20	0
3	GOL	A	2003	-	5,5,5	0.29	0	5,5,5	0.64	0
3	GOL	A	2004	-	5,5,5	0.36	0	5,5,5	0.67	0
4	SO4	A	2005	-	4,4,4	0.21	0	6,6,6	0.12	0
4	SO4	A	2006	-	4,4,4	0.21	0	6,6,6	0.24	0
4	SO4	A	2007	-	4,4,4	0.17	0	6,6,6	0.60	0
2	FAD	B	1001	-	48,58,58	1.94	15 (31%)	54,89,89	2.37	13 (24%)
4	SO4	B	1002	-	4,4,4	0.18	0	6,6,6	0.15	0
4	SO4	B	1003	-	4,4,4	0.18	0	6,6,6	0.19	0
4	SO4	B	1004	-	4,4,4	0.25	0	6,6,6	0.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	2001	-	-	0/30/50/50	0/6/6/6
3	GOL	A	2002	-	-	0/4/4/4	0/0/0/0
3	GOL	A	2003	-	-	0/4/4/4	0/0/0/0
3	GOL	A	2004	-	-	0/4/4/4	0/0/0/0
4	SO4	A	2005	-	-	0/0/0/0	0/0/0/0
4	SO4	A	2006	-	-	0/0/0/0	0/0/0/0
4	SO4	A	2007	-	-	0/0/0/0	0/0/0/0
2	FAD	B	1001	-	-	0/30/50/50	0/6/6/6
4	SO4	B	1002	-	-	0/0/0/0	0/0/0/0
4	SO4	B	1003	-	-	0/0/0/0	0/0/0/0
4	SO4	B	1004	-	-	0/0/0/0	0/0/0/0

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	FAD	O4B-C4B	-2.64	1.38	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	FAD	PA-O2A	-2.63	1.43	1.54
2	B	1001	FAD	O4B-C4B	-2.52	1.39	1.45
2	B	1001	FAD	PA-O2A	-2.43	1.44	1.54
2	A	2001	FAD	O2'-C2'	-2.42	1.37	1.43
2	A	2001	FAD	O3'-C3'	-2.40	1.37	1.43
2	B	1001	FAD	O3'-C3'	-2.28	1.37	1.43
2	B	1001	FAD	O2'-C2'	-2.27	1.38	1.43
2	A	2001	FAD	PA-O5B	-2.23	1.48	1.59
2	B	1001	FAD	O4'-C4'	-2.15	1.38	1.43
2	B	1001	FAD	O2B-C2B	-2.15	1.37	1.43
2	B	1001	FAD	PA-O5B	-2.14	1.49	1.59
2	A	2001	FAD	O4'-C4'	-2.09	1.38	1.43
2	A	2001	FAD	C2A-N3A	2.35	1.36	1.32
2	B	1001	FAD	C2A-N3A	2.50	1.36	1.32
2	A	2001	FAD	C5X-N5	2.66	1.39	1.35
2	A	2001	FAD	C6A-N6A	2.67	1.43	1.34
2	B	1001	FAD	C5X-N5	2.83	1.39	1.35
2	B	1001	FAD	C6A-N6A	3.01	1.44	1.34
2	A	2001	FAD	C4-N3	3.20	1.39	1.33
2	B	1001	FAD	C4-N3	3.40	1.39	1.33
2	A	2001	FAD	C4X-N5	3.82	1.39	1.33
2	B	1001	FAD	C9A-N10	3.85	1.44	1.38
2	B	1001	FAD	C4X-N5	3.99	1.39	1.33
2	A	2001	FAD	C4-C4X	3.99	1.49	1.41
2	A	2001	FAD	C9A-N10	4.17	1.44	1.38
2	B	1001	FAD	C4-C4X	4.29	1.49	1.41
2	A	2001	FAD	O4-C4	4.80	1.36	1.24
2	B	1001	FAD	O4-C4	5.33	1.37	1.24

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	FAD	N3A-C2A-N1A	-12.67	119.19	128.89
2	B	1001	FAD	N3A-C2A-N1A	-12.21	119.54	128.89
2	A	2001	FAD	C4-C4X-C10	-6.20	115.97	119.94
2	B	1001	FAD	C4-C4X-C10	-4.83	116.85	119.94
2	B	1001	FAD	O2P-P-O3P	-3.42	89.56	105.09
2	B	1001	FAD	O2A-PA-O5B	-2.69	94.89	108.46
2	A	2001	FAD	C4A-C5A-N7A	-2.57	107.11	109.48
2	A	2001	FAD	O2A-PA-O5B	-2.55	95.61	108.46
2	B	1001	FAD	C4X-C10-N10	-2.55	119.02	120.52
2	A	2001	FAD	O2P-P-O3P	-2.45	93.96	105.09

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1001	FAD	C4A-C5A-N7A	-2.09	107.56	109.48
2	B	1001	FAD	C4X-C4-N3	-2.05	120.78	123.59
2	B	1001	FAD	O5'-P-O1P	2.01	117.40	109.62
2	A	2001	FAD	O5B-PA-O1A	2.19	118.12	109.62
2	A	2001	FAD	O5'-P-O1P	2.31	118.58	109.62
2	B	1001	FAD	O5B-PA-O1A	2.38	118.86	109.62
2	A	2001	FAD	C5X-C9A-N10	2.62	119.61	117.62
2	A	2001	FAD	C4X-N5-C5X	2.68	119.85	116.76
2	B	1001	FAD	C4-C4X-N5	3.14	122.52	118.72
2	B	1001	FAD	C4X-N5-C5X	3.21	120.45	116.76
2	B	1001	FAD	C5X-C9A-N10	3.27	120.10	117.62
2	A	2001	FAD	C4-C4X-N5	3.73	123.24	118.72
2	A	2001	FAD	C4-N3-C2	5.97	120.40	115.25
2	B	1001	FAD	C4-N3-C2	6.05	120.47	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2001	FAD	1	0
4	B	1003	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	971/1001 (97%)	0.33	50 (5%)	32	41	20, 34, 57, 84	0
1	B	969/1001 (96%)	0.45	74 (7%)	17	24	26, 39, 61, 85	0
All	All	1940/2002 (96%)	0.39	124 (6%)	23	31	20, 37, 59, 85	0

All (124) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	434	ALA	7.8
1	A	129	VAL	6.1
1	B	129	VAL	6.1
1	B	2	PRO	5.7
1	B	409	GLY	5.5
1	B	311	ILE	4.9
1	B	224	GLY	4.9
1	B	509	PRO	4.9
1	B	3	ASN	4.8
1	A	224	GLY	4.7
1	B	226	HIS	4.7
1	B	310	ALA	4.6
1	B	133	GLY	4.3
1	B	225	ASN	4.3
1	A	188	THR	4.1
1	A	2	PRO	4.1
1	B	223	ALA	4.0
1	B	340	LEU	4.0
1	B	188	THR	3.8
1	A	186	GLN	3.7
1	A	509	PRO	3.7
1	A	30	SER	3.7
1	B	131	GLN	3.7
1	B	227	ALA	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	132	PRO	3.6
1	A	133	GLY	3.4
1	A	3	ASN	3.4
1	B	338	VAL	3.4
1	A	546	THR	3.4
1	A	542	ILE	3.3
1	B	4	ILE	3.3
1	B	865	GLY	3.2
1	A	26	ALA	3.2
1	B	31	PRO	3.2
1	B	270	ALA	3.2
1	B	462	GLY	3.2
1	A	661	LEU	3.2
1	B	87	ASP	3.2
1	B	546	THR	3.1
1	A	132	PRO	3.1
1	A	131	GLN	3.1
1	A	434	ALA	3.1
1	B	663	ILE	3.0
1	A	87	ASP	3.0
1	B	548	ASP	3.0
1	B	30	SER	3.0
1	A	311	ILE	3.0
1	A	548	ASP	3.0
1	A	310	ALA	3.0
1	A	226	HIS	2.9
1	A	508	ALA	2.9
1	B	634	PRO	2.9
1	B	542	ILE	2.9
1	A	493	GLN	2.9
1	A	31	PRO	2.8
1	B	901	GLU	2.8
1	B	55	LEU	2.8
1	B	393	ALA	2.8
1	A	32	PRO	2.8
1	B	490	VAL	2.8
1	A	663	ILE	2.8
1	A	92	ASP	2.7
1	B	25	PRO	2.7
1	A	393	ALA	2.7
1	A	463	ALA	2.7
1	A	559	ALA	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	435	ASP	2.7
1	B	189	ARG	2.7
1	B	318	ALA	2.7
1	B	756	ILE	2.6
1	B	28	HIS	2.6
1	B	309	LEU	2.6
1	A	668	VAL	2.6
1	A	898	GLU	2.6
1	A	680	ALA	2.6
1	B	629	SER	2.5
1	A	534	ALA	2.5
1	B	341	VAL	2.5
1	B	543	ALA	2.5
1	B	222	ALA	2.4
1	A	185	GLY	2.4
1	B	864	GLU	2.4
1	B	575	ALA	2.4
1	B	878	ALA	2.4
1	A	665	LEU	2.4
1	B	504	CYS	2.4
1	A	749	ASP	2.3
1	A	659	PHE	2.3
1	A	225	ASN	2.3
1	A	28	HIS	2.3
1	B	665	LEU	2.3
1	B	464	ASN	2.3
1	B	90	THR	2.3
1	B	628	GLY	2.3
1	B	661	LEU	2.3
1	B	523	THR	2.2
1	A	21	ALA	2.2
1	B	21	ALA	2.2
1	B	122	ALA	2.2
1	B	670	ALA	2.2
1	B	668	VAL	2.2
1	A	338	VAL	2.2
1	B	392	PHE	2.2
1	A	901	GLU	2.2
1	B	899	ASN	2.2
1	A	15	ASP	2.1
1	A	25	PRO	2.1
1	B	408	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	387	ARG	2.1
1	B	669	THR	2.1
1	B	331	ALA	2.1
1	A	309	LEU	2.1
1	B	35	ALA	2.1
1	B	97	ASP	2.1
1	A	834	VAL	2.1
1	A	93	GLN	2.0
1	B	333	ASP	2.0
1	B	32	PRO	2.0
1	B	492	PRO	2.0
1	B	279	ALA	2.0
1	A	460	GLU	2.0
1	A	464	ASN	2.0
1	B	93	GLN	2.0
1	B	700	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	SO4	B	1004	5/5	0.94	0.26	3.19	58,60,64,68	0
4	SO4	A	2006	5/5	0.95	0.20	2.44	48,52,59,63	0
3	GOL	A	2004	6/6	0.93	0.13	2.08	32,43,46,48	0
4	SO4	B	1002	5/5	0.97	0.19	0.25	57,61,65,67	0
4	SO4	A	2007	5/5	0.97	0.16	-0.24	45,48,53,53	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FAD	B	1001	53/53	0.95	0.15	-0.58	20,32,41,42	0
2	FAD	A	2001	53/53	0.96	0.14	-0.87	17,23,31,44	0
3	GOL	A	2003	6/6	0.93	0.12	-1.08	42,46,48,50	0
4	SO4	B	1003	5/5	0.89	0.27	-	84,85,90,92	0
4	SO4	A	2005	5/5	0.88	0.32	-	82,82,86,87	0
3	GOL	A	2002	6/6	0.88	0.13	-	51,53,53,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.