



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:54 AM GMT

PDB ID : 3Q8U  
Title : Crystal structure of Staphylococcus aureus nucleoside diphosphate kinase complexed with ADP  
Authors : Srivastava, S.K.; Rajasree, K.; Gopal, B.  
Deposited on : 2011-01-07  
Resolution : 2.22 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4405 (2.24-2.20)
Clashscore	102246	5146 (2.24-2.20)
Ramachandran outliers	100387	5065 (2.24-2.20)
Sidechain outliers	100360	5066 (2.24-2.20)
RSRZ outliers	91569	4414 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	157	<div> <div>73%</div> <div>17%</div> <div>• • 5%</div> </div>
1	B	157	<div> <div>75%</div> <div>17%</div> <div>• • 5%</div> </div>
1	C	157	<div> <div>82%</div> <div>10%</div> <div>• • 5%</div> </div>
1	D	157	<div> <div>%</div> <div>83%</div> <div>10%</div> <div>• • 5%</div> </div>
1	E	157	<div> <div>%</div> <div>69%</div> <div>24%</div> <div>• 5%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	157	<div><div></div><div>3%</div><div>77%</div><div>14%</div><div>• 5%</div></div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7566 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoside diphosphate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	149	Total	C	N	O	S	0	0	0
			1165	733	202	225	5			
1	B	149	Total	C	N	O	S	0	0	0
			1158	730	202	221	5			
1	C	149	Total	C	N	O	S	0	0	0
			1158	730	202	221	5			
1	D	149	Total	C	N	O	S	0	0	0
			1161	731	202	223	5			
1	E	149	Total	C	N	O	S	0	0	0
			1158	730	202	221	5			
1	F	149	Total	C	N	O	S	0	0	0
			1158	730	202	221	5			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	150	LEU	-	EXPRESSION TAG	UNP Q5HFBV4
A	151	GLU	-	EXPRESSION TAG	UNP Q5HFBV4
A	152	HIS	-	EXPRESSION TAG	UNP Q5HFBV4
A	153	HIS	-	EXPRESSION TAG	UNP Q5HFBV4
A	154	HIS	-	EXPRESSION TAG	UNP Q5HFBV4
A	155	HIS	-	EXPRESSION TAG	UNP Q5HFBV4
A	156	HIS	-	EXPRESSION TAG	UNP Q5HFBV4
A	157	HIS	-	EXPRESSION TAG	UNP Q5HFBV4
B	150	LEU	-	EXPRESSION TAG	UNP Q5HFBV4
B	151	GLU	-	EXPRESSION TAG	UNP Q5HFBV4
B	152	HIS	-	EXPRESSION TAG	UNP Q5HFBV4
B	153	HIS	-	EXPRESSION TAG	UNP Q5HFBV4
B	154	HIS	-	EXPRESSION TAG	UNP Q5HFBV4
B	155	HIS	-	EXPRESSION TAG	UNP Q5HFBV4
B	156	HIS	-	EXPRESSION TAG	UNP Q5HFBV4
B	157	HIS	-	EXPRESSION TAG	UNP Q5HFBV4
C	150	LEU	-	EXPRESSION TAG	UNP Q5HFBV4

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	151	GLU	-	EXPRESSION TAG	UNP Q5HFV4
C	152	HIS	-	EXPRESSION TAG	UNP Q5HFV4
C	153	HIS	-	EXPRESSION TAG	UNP Q5HFV4
C	154	HIS	-	EXPRESSION TAG	UNP Q5HFV4
C	155	HIS	-	EXPRESSION TAG	UNP Q5HFV4
C	156	HIS	-	EXPRESSION TAG	UNP Q5HFV4
C	157	HIS	-	EXPRESSION TAG	UNP Q5HFV4
D	150	LEU	-	EXPRESSION TAG	UNP Q5HFV4
D	151	GLU	-	EXPRESSION TAG	UNP Q5HFV4
D	152	HIS	-	EXPRESSION TAG	UNP Q5HFV4
D	153	HIS	-	EXPRESSION TAG	UNP Q5HFV4
D	154	HIS	-	EXPRESSION TAG	UNP Q5HFV4
D	155	HIS	-	EXPRESSION TAG	UNP Q5HFV4
D	156	HIS	-	EXPRESSION TAG	UNP Q5HFV4
D	157	HIS	-	EXPRESSION TAG	UNP Q5HFV4
E	150	LEU	-	EXPRESSION TAG	UNP Q5HFV4
E	151	GLU	-	EXPRESSION TAG	UNP Q5HFV4
E	152	HIS	-	EXPRESSION TAG	UNP Q5HFV4
E	153	HIS	-	EXPRESSION TAG	UNP Q5HFV4
E	154	HIS	-	EXPRESSION TAG	UNP Q5HFV4
E	155	HIS	-	EXPRESSION TAG	UNP Q5HFV4
E	156	HIS	-	EXPRESSION TAG	UNP Q5HFV4
E	157	HIS	-	EXPRESSION TAG	UNP Q5HFV4
F	150	LEU	-	EXPRESSION TAG	UNP Q5HFV4
F	151	GLU	-	EXPRESSION TAG	UNP Q5HFV4
F	152	HIS	-	EXPRESSION TAG	UNP Q5HFV4
F	153	HIS	-	EXPRESSION TAG	UNP Q5HFV4
F	154	HIS	-	EXPRESSION TAG	UNP Q5HFV4
F	155	HIS	-	EXPRESSION TAG	UNP Q5HFV4
F	156	HIS	-	EXPRESSION TAG	UNP Q5HFV4
F	157	HIS	-	EXPRESSION TAG	UNP Q5HFV4

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	D	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	E	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	F	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		
3	F	1	Total	Mg	0	0
			1	1		
3	E	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	85	Total 85	O 85	0	0
4	B	81	Total 81	O 81	0	0
4	C	76	Total 76	O 76	0	0
4	D	53	Total 53	O 53	0	0
4	E	95	Total 95	O 95	0	0
4	F	51	Total 51	O 51	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nucleoside diphosphate kinase




- Molecule 1: Nucleoside diphosphate kinase



- Molecule 1: Nucleoside diphosphate kinase




- Molecule 1: Nucleoside diphosphate kinase





- Molecule 1: Nucleoside diphosphate kinase







● Molecule 1: Nucleoside diphosphate kinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.27Å 103.06Å 127.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.52 – 2.22 34.52 – 2.22	Depositor EDS
% Data completeness (in resolution range)	97.8 (34.52-2.22) 97.8 (34.52-2.22)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.86 (at 2.22Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.180 , 0.258 0.180 , 0.256	Depositor DCC
$R_{free}$ test set	2221 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.1	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 37.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 44308 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7566	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.41% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.01	3/1188 (0.3%)	1.07	7/1607 (0.4%)
1	B	0.98	0/1181	0.99	5/1598 (0.3%)
1	C	1.00	1/1181 (0.1%)	0.98	5/1598 (0.3%)
1	D	0.97	0/1184	1.02	5/1602 (0.3%)
1	E	1.00	1/1181 (0.1%)	0.98	3/1598 (0.2%)
1	F	0.95	1/1181 (0.1%)	0.97	4/1598 (0.3%)
All	All	0.98	6/7096 (0.1%)	1.00	29/9601 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	133	GLU	CD-OE1	-6.02	1.19	1.25
1	F	124	GLU	CG-CD	5.49	1.60	1.51
1	A	145	ALA	CA-CB	5.47	1.64	1.52
1	A	133	GLU	CD-OE2	-5.37	1.19	1.25
1	E	124	GLU	CG-CD	5.17	1.59	1.51

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	15	ARG	NE-CZ-NH1	-13.36	113.62	120.30
1	E	15	ARG	NE-CZ-NH1	-11.79	114.41	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	15	ARG	NE-CZ-NH2	10.90	125.75	120.30
1	F	102	ARG	NE-CZ-NH2	10.45	125.53	120.30
1	F	102	ARG	NE-CZ-NH1	-10.45	115.08	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	54	GLY	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1165	0	1149	32	1
1	B	1158	0	1143	28	0
1	C	1158	0	1143	17	0
1	D	1161	0	1145	12	1
1	E	1158	0	1143	43	0
1	F	1158	0	1142	20	0
2	A	27	0	12	0	0
2	B	27	0	12	0	0
2	C	27	0	12	2	0
2	D	27	0	12	0	0
2	E	27	0	12	0	0
2	F	27	0	12	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
4	A	85	0	0	19	0
4	B	81	0	0	16	2
4	C	76	0	0	9	0
4	D	53	0	0	1	0
4	E	95	0	0	14	2
4	F	51	0	0	5	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	7566	0	6937	144	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 144 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:122:SER:HB3	4:F:430:HOH:O	1.15	1.32
4:A:168:HOH:O	1:E:60:ASP:HB3	1.29	1.31
1:C:82:ASN:HB3	4:C:401:HOH:O	1.28	1.27
1:A:82:ASN:HB3	4:A:414:HOH:O	1.22	1.27
1:E:145:ALA:HB2	4:E:368:HOH:O	1.32	1.25

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:426:HOH:O	4:E:363:HOH:O[4_455]	1.12	1.08
4:B:172:HOH:O	4:E:364:HOH:O[4_455]	1.28	0.92
1:A:133:GLU:OE1	1:D:86:HIS:ND1[3_544]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	147/157 (94%)	142 (97%)	5 (3%)	0	100	100
1	B	147/157 (94%)	143 (97%)	4 (3%)	0	100	100
1	C	147/157 (94%)	143 (97%)	3 (2%)	1 (1%)	26	25
1	D	147/157 (94%)	144 (98%)	3 (2%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	147/157 (94%)	144 (98%)	3 (2%)	0	100	100
1	F	147/157 (94%)	143 (97%)	4 (3%)	0	100	100
All	All	882/942 (94%)	859 (97%)	22 (2%)	1 (0%)	56	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	53	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	127/135 (94%)	118 (93%)	9 (7%)	18	18
1	B	125/135 (93%)	118 (94%)	7 (6%)	26	29
1	C	125/135 (93%)	121 (97%)	4 (3%)	46	57
1	D	126/135 (93%)	120 (95%)	6 (5%)	31	36
1	E	125/135 (93%)	119 (95%)	6 (5%)	31	36
1	F	125/135 (93%)	119 (95%)	6 (5%)	31	36
All	All	753/810 (93%)	715 (95%)	38 (5%)	30	34

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	61	LEU
1	D	47	THR
1	F	61	LEU
1	C	134	ASN
1	D	61	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	16	ASN
1	D	39	GLN
1	E	134	ASN
1	C	134	ASN
1	E	16	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ADP	A	158	3	22,29,29	1.20	2 (9%)	27,45,45	2.09	5 (18%)
2	ADP	B	158	3	22,29,29	0.93	1 (4%)	27,45,45	2.01	5 (18%)
2	ADP	C	158	-	22,29,29	1.05	2 (9%)	27,45,45	2.18	7 (25%)
2	ADP	D	158	3	22,29,29	1.11	1 (4%)	27,45,45	2.15	5 (18%)
2	ADP	E	158	3	22,29,29	0.92	1 (4%)	27,45,45	1.87	3 (11%)
2	ADP	F	158	3	22,29,29	1.25	3 (13%)	27,45,45	2.47	8 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	158	3	-	0/12/32/32	0/3/3/3
2	ADP	B	158	3	-	0/12/32/32	0/3/3/3
2	ADP	C	158	-	-	0/12/32/32	0/3/3/3
2	ADP	D	158	3	-	0/12/32/32	0/3/3/3
2	ADP	E	158	3	-	0/12/32/32	0/3/3/3
2	ADP	F	158	3	-	0/12/32/32	0/3/3/3

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	158	ADP	O4'-C1'	2.06	1.43	1.41
2	C	158	ADP	C2-N3	2.21	1.36	1.32
2	A	158	ADP	C2-N3	2.27	1.36	1.32
2	F	158	ADP	C2-N3	2.28	1.36	1.32
2	B	158	ADP	C5-C4	2.45	1.46	1.40

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	158	ADP	N3-C2-N1	-7.98	122.79	128.89
2	E	158	ADP	N3-C2-N1	-7.75	122.96	128.89
2	A	158	ADP	N3-C2-N1	-7.64	123.04	128.89
2	C	158	ADP	N3-C2-N1	-6.95	123.57	128.89
2	B	158	ADP	N3-C2-N1	-6.76	123.72	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	158	ADP	2	0

## 5.7 Other polymers

There are no such residues in this entry.



## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	149/157 (94%)	-0.33	0	100 100	8, 15, 29, 36	0
1	B	149/157 (94%)	-0.50	0	100 100	7, 16, 29, 34	0
1	C	149/157 (94%)	-0.32	0	100 100	8, 17, 30, 36	0
1	D	149/157 (94%)	-0.33	2 (1%)	79 78	8, 17, 30, 36	0
1	E	149/157 (94%)	-0.46	2 (1%)	79 78	6, 16, 28, 35	0
1	F	149/157 (94%)	-0.25	4 (2%)	58 57	8, 17, 31, 38	0
All	All	894/942 (94%)	-0.37	8 (0%)	85 85	6, 16, 30, 38	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	148	TYR	3.9
1	E	147	LEU	3.1
1	F	53	GLN	2.9
1	D	54	GLY	2.8
1	D	149	GLU	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	ADP	C	158	27/27	0.96	0.15	0.93	23,33,37,38	0
2	ADP	F	158	27/27	0.93	0.14	0.47	27,38,43,44	0
2	ADP	A	158	27/27	0.98	0.11	0.10	18,22,24,25	0
2	ADP	D	158	27/27	0.95	0.11	-0.30	21,29,36,37	0
2	ADP	E	158	27/27	0.98	0.08	-0.58	8,13,17,18	0
2	ADP	B	158	27/27	0.98	0.07	-0.91	9,14,21,22	0
3	MG	E	159	1/1	0.96	0.05	-	24,24,24,24	0
3	MG	B	159	1/1	0.98	0.08	-	2,2,2,2	0
3	MG	F	159	1/1	0.82	0.08	-	43,43,43,43	0
3	MG	A	159	1/1	0.86	0.17	-	48,48,48,48	0
3	MG	D	159	1/1	0.95	0.05	-	40,40,40,40	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.