



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:56 PM GMT

PDB ID : 4QEY
Title : Crystal structure of a DUF4425 family protein (BACOVA_05332) from *Bacteroides ovatus* ATCC 8483 at 2.52 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2014-05-19
Resolution : 2.52 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4241 (2.54-2.50)
Clashscore	102246	4968 (2.54-2.50)
Ramachandran outliers	100387	4873 (2.54-2.50)
Sidechain outliers	100360	4875 (2.54-2.50)
RSRZ outliers	91569	4253 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	132	<div> <div>3%</div> <div>86%</div> <div>9%</div> <div>5%</div> </div>
1	B	132	<div> <div>7%</div> <div>86%</div> <div>8%</div> <div>6%</div> </div>
1	C	132	<div> <div>6%</div> <div>86%</div> <div>8%</div> <div>6%</div> </div>
1	D	132	<div> <div>3%</div> <div>86%</div> <div>7%</div> <div>7%</div> </div>
1	E	132	<div> <div>2%</div> <div>88%</div> <div>6%</div> <div>6%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	132	
1	G	132	
1	H	132	
1	I	132	
1	J	132	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PEG	A	201	-	-	-	X
2	PEG	D	201	-	-	-	X
2	PEG	E	201	-	-	-	X
2	PEG	J	201	-	-	-	X
3	EDO	A	202	-	-	X	X
3	EDO	A	206	-	-	-	X
3	EDO	A	207	-	-	-	X
3	EDO	C	201	-	-	-	X
3	EDO	C	202	-	-	-	X
3	EDO	D	202	-	-	-	X
3	EDO	E	203	-	-	-	X
3	EDO	G	201	-	-	-	X
3	EDO	G	203	-	-	-	X
3	EDO	H	202	-	-	-	X
3	EDO	J	202	-	-	-	X
3	EDO	J	204	-	-	-	X
4	PGE	H	201	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9789 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	125	Total	C	N	O	S	Se	0	3	0
			940	578	154	204	2	2			
1	B	124	Total	C	N	O	S	Se	0	5	0
			958	588	156	210	2	2			
1	C	124	Total	C	N	O	S	Se	0	3	0
			931	573	153	201	2	2			
1	D	123	Total	C	N	O	S	Se	0	4	0
			919	566	150	199	2	2			
1	E	124	Total	C	N	O	S	Se	0	5	0
			945	582	153	206	2	2			
1	F	124	Total	C	N	O	S	Se	0	6	0
			963	594	157	208	2	2			
1	G	125	Total	C	N	O	S	Se	0	4	0
			957	591	158	204	2	2			
1	H	125	Total	C	N	O	S	Se	0	5	0
			953	586	157	206	2	2			
1	I	125	Total	C	N	O	S	Se	0	4	0
			937	577	154	202	2	2			
1	J	124	Total	C	N	O	S	Se	0	3	0
			926	569	152	201	2	2			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	LEADER SEQUENCE	UNP A7M5D7
B	0	GLY	-	LEADER SEQUENCE	UNP A7M5D7
C	0	GLY	-	LEADER SEQUENCE	UNP A7M5D7
D	0	GLY	-	LEADER SEQUENCE	UNP A7M5D7
E	0	GLY	-	LEADER SEQUENCE	UNP A7M5D7
F	0	GLY	-	LEADER SEQUENCE	UNP A7M5D7
G	0	GLY	-	LEADER SEQUENCE	UNP A7M5D7
H	0	GLY	-	LEADER SEQUENCE	UNP A7M5D7
I	0	GLY	-	LEADER SEQUENCE	UNP A7M5D7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
J	0	GLY	-	LEADER SEQUENCE	UNP A7M5D7

- Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		
2	B	1	Total	C	O	0	0
			7	4	3		
2	D	1	Total	C	O	0	0
			7	4	3		
2	E	1	Total	C	O	0	0
			7	4	3		
2	I	1	Total	C	O	0	0
			7	4	3		
2	J	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		
3	E	1	Total	C	O	0	0
			4	2	2		
3	E	1	Total	C	O	0	0
			4	2	2		
3	E	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	E	1	Total 4	C 2	O 2	0	0
3	E	1	Total 4	C 2	O 2	0	0
3	E	1	Total 4	C 2	O 2	0	0
3	F	1	Total 4	C 2	O 2	0	0
3	G	1	Total 4	C 2	O 2	0	0
3	G	1	Total 4	C 2	O 2	0	0
3	G	1	Total 4	C 2	O 2	0	0
3	H	1	Total 4	C 2	O 2	0	0
3	H	1	Total 4	C 2	O 2	0	0
3	J	1	Total 4	C 2	O 2	0	0
3	J	1	Total 4	C 2	O 2	0	0
3	J	1	Total 4	C 2	O 2	0	0
3	J	1	Total 4	C 2	O 2	0	0
3	J	1	Total 4	C 2	O 2	0	0

- Molecule 4 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	H	1	Total	C	O	0	0
			10	6	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	21	Total	O	0	0
			21	21		
5	B	17	Total	O	0	0
			17	17		
5	C	21	Total	O	0	0
			21	21		
5	D	16	Total	O	0	0
			16	16		
5	E	28	Total	O	0	0
			28	28		
5	F	16	Total	O	0	1
			17	17		
5	G	14	Total	O	0	1
			15	15		
5	H	21	Total	O	0	0
			21	21		
5	I	20	Total	O	0	0
			20	20		
5	J	20	Total	O	0	0
			20	20		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Uncharacterized protein




- Molecule 1: Uncharacterized protein





- Molecule 1: Uncharacterized protein





- Molecule 1: Uncharacterized protein





- Molecule 1: Uncharacterized protein




- Molecule 1: Uncharacterized protein





- Molecule 1: Uncharacterized protein




- Molecule 1: Uncharacterized protein



- Molecule 1: Uncharacterized protein



- Molecule 1: Uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	90.59Å 101.34Å 91.42Å 90.00° 107.91° 90.00°	Depositor
Resolution (Å)	47.35 – 2.52 47.34 – 2.52	Depositor EDS
% Data completeness (in resolution range)	98.5 (47.35-2.52) 98.6 (47.34-2.52)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 2.51Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.196 , 0.221 0.202 , 0.223	Depositor DCC
R_{free} test set	2670 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	51.3	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 54.2	EDS
Estimated twinning fraction	0.000 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 52414 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9789	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 54.89 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.4226e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, PGE, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/946	0.68	0/1275
1	B	0.44	0/964	0.68	0/1297
1	C	0.43	0/937	0.68	0/1263
1	D	0.43	0/928	0.68	0/1253
1	E	0.46	0/957	0.69	0/1289
1	F	0.45	0/978	0.69	0/1314
1	G	0.46	0/966	0.67	0/1297
1	H	0.45	0/965	0.69	0/1300
1	I	0.45	0/946	0.68	0/1276
1	J	0.47	0/932	0.70	0/1258
All	All	0.45	0/9519	0.68	0/12822

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	940	0	903	11	0
1	B	958	0	919	7	0
1	C	931	0	897	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	919	0	872	5	0
1	E	945	0	908	4	0
1	F	963	0	950	7	0
1	G	957	0	947	7	0
1	H	953	0	924	7	0
1	I	937	0	899	7	0
1	J	926	0	884	8	0
2	A	7	0	10	1	0
2	B	7	0	10	1	0
2	D	7	0	10	1	0
2	E	7	0	10	0	0
2	I	7	0	10	0	0
2	J	7	0	10	3	0
3	A	24	0	36	4	0
3	B	8	0	12	0	0
3	C	8	0	12	0	0
3	D	4	0	6	0	0
3	E	24	0	36	0	0
3	F	4	0	6	0	0
3	G	12	0	18	0	0
3	H	8	0	12	0	0
3	J	20	0	30	0	0
4	H	10	0	14	1	0
5	A	21	0	0	0	0
5	B	17	0	0	0	0
5	C	21	0	0	0	0
5	D	16	0	0	0	0
5	E	28	0	0	0	0
5	F	17	0	0	0	0
5	G	15	0	0	0	0
5	H	21	0	0	0	0
5	I	20	0	0	0	0
5	J	20	0	0	1	0
All	All	9789	0	9345	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 63 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:59:ASN:HD22	1:B:154:VAL:HB	1.58	0.69
1:A:51:LEU:H	3:A:202:EDO:H21	1.58	0.68
1:F:154:VAL:HB	1:G:59:ASN:HD22	1.62	0.65
1:A:102:ALA:HB1	2:A:201:PEG:H11	1.79	0.64
1:A:51:LEU:H	3:A:202:EDO:C2	2.17	0.56

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	126/132 (96%)	117 (93%)	9 (7%)	0	100	100
1	B	127/132 (96%)	122 (96%)	5 (4%)	0	100	100
1	C	125/132 (95%)	123 (98%)	2 (2%)	0	100	100
1	D	125/132 (95%)	120 (96%)	5 (4%)	0	100	100
1	E	127/132 (96%)	122 (96%)	5 (4%)	0	100	100
1	F	128/132 (97%)	123 (96%)	5 (4%)	0	100	100
1	G	127/132 (96%)	123 (97%)	4 (3%)	0	100	100
1	H	128/132 (97%)	123 (96%)	5 (4%)	0	100	100
1	I	127/132 (96%)	121 (95%)	6 (5%)	0	100	100
1	J	125/132 (95%)	118 (94%)	7 (6%)	0	100	100
All	All	1265/1320 (96%)	1212 (96%)	53 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	104/110 (94%)	104 (100%)	0	100	100
1	B	107/110 (97%)	106 (99%)	1 (1%)	84	95
1	C	103/110 (94%)	103 (100%)	0	100	100
1	D	100/110 (91%)	100 (100%)	0	100	100
1	E	105/110 (96%)	105 (100%)	0	100	100
1	F	110/110 (100%)	109 (99%)	1 (1%)	84	95
1	G	108/110 (98%)	108 (100%)	0	100	100
1	H	107/110 (97%)	106 (99%)	1 (1%)	84	95
1	I	103/110 (94%)	102 (99%)	1 (1%)	82	94
1	J	102/110 (93%)	102 (100%)	0	100	100
All	All	1049/1100 (95%)	1045 (100%)	4 (0%)	93	98

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	81	ASN
1	F	47	ASP
1	H	47	ASP
1	I	117	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

35 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PEG	A	201	-	6,6,6	0.42	0	5,5,5	0.21	0
3	EDO	A	202	-	3,3,3	0.47	0	2,2,2	0.63	0
3	EDO	A	203	-	3,3,3	0.74	0	2,2,2	0.23	0
3	EDO	A	204	-	3,3,3	0.64	0	2,2,2	0.49	0
3	EDO	A	205	-	3,3,3	0.46	0	2,2,2	0.64	0
3	EDO	A	206	-	3,3,3	0.58	0	2,2,2	0.55	0
3	EDO	A	207	-	3,3,3	0.69	0	2,2,2	0.39	0
2	PEG	B	201	-	6,6,6	0.24	0	5,5,5	0.13	0
3	EDO	B	202	-	3,3,3	0.58	0	2,2,2	0.37	0
3	EDO	B	203	-	3,3,3	0.54	0	2,2,2	0.53	0
3	EDO	C	201	-	3,3,3	0.69	0	2,2,2	0.27	0
3	EDO	C	202	-	3,3,3	0.67	0	2,2,2	0.30	0
2	PEG	D	201	-	6,6,6	0.28	0	5,5,5	0.08	0
3	EDO	D	202	-	3,3,3	0.75	0	2,2,2	0.30	0
2	PEG	E	201	-	6,6,6	0.19	0	5,5,5	0.11	0
3	EDO	E	202	-	3,3,3	0.61	0	2,2,2	0.40	0
3	EDO	E	203	-	3,3,3	0.49	0	2,2,2	0.48	0
3	EDO	E	204	-	3,3,3	0.56	0	2,2,2	0.47	0
3	EDO	E	205	-	3,3,3	0.47	0	2,2,2	0.51	0
3	EDO	E	206	-	3,3,3	0.63	0	2,2,2	0.33	0
3	EDO	E	207	-	3,3,3	0.55	0	2,2,2	0.52	0
3	EDO	F	201	-	3,3,3	0.70	0	2,2,2	0.17	0
3	EDO	G	201	-	3,3,3	0.70	0	2,2,2	0.19	0
3	EDO	G	202	-	3,3,3	0.48	0	2,2,2	0.55	0
3	EDO	G	203	-	3,3,3	0.56	0	2,2,2	0.42	0
4	PGE	H	201	-	9,9,9	0.29	0	8,8,8	0.33	0
3	EDO	H	202	-	3,3,3	0.47	0	2,2,2	0.51	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	H	203	-	3,3,3	0.52	0	2,2,2	0.59	0
2	PEG	I	201	-	6,6,6	0.30	0	5,5,5	0.19	0
2	PEG	J	201	-	6,6,6	0.28	0	5,5,5	0.23	0
3	EDO	J	202	-	3,3,3	0.76	0	2,2,2	0.04	0
3	EDO	J	203	-	3,3,3	0.64	0	2,2,2	0.41	0
3	EDO	J	204	-	3,3,3	0.44	0	2,2,2	0.55	0
3	EDO	J	205	-	3,3,3	0.54	0	2,2,2	0.55	0
3	EDO	J	206	-	3,3,3	0.58	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	A	201	-	-	0/4/4/4	0/0/0/0
3	EDO	A	202	-	-	0/1/1/1	0/0/0/0
3	EDO	A	203	-	-	0/1/1/1	0/0/0/0
3	EDO	A	204	-	-	0/1/1/1	0/0/0/0
3	EDO	A	205	-	-	0/1/1/1	0/0/0/0
3	EDO	A	206	-	-	0/1/1/1	0/0/0/0
3	EDO	A	207	-	-	0/1/1/1	0/0/0/0
2	PEG	B	201	-	-	0/4/4/4	0/0/0/0
3	EDO	B	202	-	-	0/1/1/1	0/0/0/0
3	EDO	B	203	-	-	0/1/1/1	0/0/0/0
3	EDO	C	201	-	-	0/1/1/1	0/0/0/0
3	EDO	C	202	-	-	0/1/1/1	0/0/0/0
2	PEG	D	201	-	-	0/4/4/4	0/0/0/0
3	EDO	D	202	-	-	0/1/1/1	0/0/0/0
2	PEG	E	201	-	-	0/4/4/4	0/0/0/0
3	EDO	E	202	-	-	0/1/1/1	0/0/0/0
3	EDO	E	203	-	-	0/1/1/1	0/0/0/0
3	EDO	E	204	-	-	0/1/1/1	0/0/0/0
3	EDO	E	205	-	-	0/1/1/1	0/0/0/0
3	EDO	E	206	-	-	0/1/1/1	0/0/0/0
3	EDO	E	207	-	-	0/1/1/1	0/0/0/0
3	EDO	F	201	-	-	0/1/1/1	0/0/0/0
3	EDO	G	201	-	-	0/1/1/1	0/0/0/0
3	EDO	G	202	-	-	0/1/1/1	0/0/0/0
3	EDO	G	203	-	-	0/1/1/1	0/0/0/0
4	PGE	H	201	-	-	0/7/7/7	0/0/0/0
3	EDO	H	202	-	-	0/1/1/1	0/0/0/0
3	EDO	H	203	-	-	0/1/1/1	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	I	201	-	-	0/4/4/4	0/0/0/0
2	PEG	J	201	-	-	0/4/4/4	0/0/0/0
3	EDO	J	202	-	-	0/1/1/1	0/0/0/0
3	EDO	J	203	-	-	0/1/1/1	0/0/0/0
3	EDO	J	204	-	-	0/1/1/1	0/0/0/0
3	EDO	J	205	-	-	0/1/1/1	0/0/0/0
3	EDO	J	206	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	PEG	1	0
3	A	202	EDO	4	0
3	A	203	EDO	1	0
2	B	201	PEG	1	0
2	D	201	PEG	1	0
4	H	201	PGE	1	0
2	J	201	PEG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	123/132 (93%)	0.37	4 (3%)	50	55	34, 52, 83, 102	0
1	B	122/132 (92%)	0.51	9 (7%)	17	20	38, 59, 86, 94	0
1	C	122/132 (92%)	0.58	8 (6%)	22	24	39, 55, 80, 97	0
1	D	121/132 (91%)	0.47	4 (3%)	50	55	38, 56, 80, 95	0
1	E	122/132 (92%)	0.43	3 (2%)	61	65	31, 48, 75, 87	0
1	F	122/132 (92%)	0.49	8 (6%)	22	24	37, 54, 80, 95	0
1	G	123/132 (93%)	0.39	4 (3%)	50	55	36, 53, 83, 98	0
1	H	123/132 (93%)	0.55	9 (7%)	18	20	34, 52, 82, 114	0
1	I	123/132 (93%)	0.44	5 (4%)	41	46	35, 56, 82, 100	0
1	J	122/132 (92%)	0.45	2 (1%)	74	78	36, 50, 72, 87	0
All	All	1223/1320 (92%)	0.47	56 (4%)	36	42	31, 54, 82, 114	0

The worst 5 of 56 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	154	VAL	5.2
1	F	62	GLY	4.6
1	C	154	VAL	4.4
1	C	108[A]	THR	4.3
1	F	63	VAL	4.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	EDO	A	207	4/4	0.81	0.35	17.62	65,66,67,67	0
3	EDO	A	206	4/4	0.82	0.56	16.18	66,66,67,68	0
3	EDO	J	204	4/4	0.84	0.49	15.93	78,78,79,80	0
3	EDO	G	203	4/4	0.79	0.29	15.85	72,75,77,78	0
2	PEG	E	201	7/7	0.78	0.56	10.04	64,70,81,82	0
3	EDO	A	202	4/4	0.96	0.40	9.85	43,44,47,47	0
3	EDO	D	202	4/4	0.63	0.39	9.24	68,72,73,74	0
3	EDO	C	201	4/4	0.91	0.23	7.04	53,55,59,61	0
3	EDO	H	202	4/4	0.91	0.31	6.49	67,68,69,69	0
3	EDO	C	202	4/4	0.57	0.36	5.52	77,77,78,78	0
2	PEG	A	201	7/7	0.82	0.30	4.58	47,55,57,58	0
4	PGE	H	201	10/10	0.88	0.29	4.08	52,56,59,60	0
3	EDO	J	202	4/4	0.90	0.34	3.95	56,58,59,60	0
2	PEG	J	201	7/7	0.64	0.28	3.80	63,69,77,78	0
3	EDO	G	201	4/4	0.77	0.30	3.72	58,58,59,59	0
2	PEG	D	201	7/7	0.80	0.28	3.68	65,71,73,73	0
3	EDO	E	203	4/4	0.90	0.26	3.31	69,72,75,76	0
3	EDO	G	202	4/4	0.97	0.29	1.96	62,62,65,68	0
3	EDO	E	202	4/4	0.91	0.20	1.69	48,49,52,53	0
3	EDO	A	204	4/4	0.83	0.27	1.45	56,57,61,63	0
2	PEG	I	201	7/7	0.90	0.21	1.19	62,66,68,69	0
3	EDO	J	205	4/4	0.73	0.23	0.41	70,72,74,75	0
2	PEG	B	201	7/7	0.89	0.21	0.22	58,58,64,64	0
3	EDO	E	205	4/4	0.93	0.20	0.20	72,72,72,73	0
3	EDO	A	203	4/4	0.85	0.15	-0.71	58,60,62,63	0
3	EDO	F	201	4/4	0.88	0.14	-1.48	60,62,63,63	0
3	EDO	H	203	4/4	0.89	0.14	-1.65	69,69,69,70	0
3	EDO	B	202	4/4	0.80	0.14	-1.95	84,84,84,85	0
3	EDO	J	203	4/4	0.95	0.12	-1.95	53,54,55,55	0
3	EDO	A	205	4/4	0.80	0.42	-	65,66,66,70	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	EDO	E	204	4/4	0.88	0.30	-	64,66,67,67	0
3	EDO	E	206	4/4	0.69	0.19	-	87,87,88,89	0
3	EDO	B	203	4/4	0.84	0.22	-	76,76,77,77	0
3	EDO	E	207	4/4	0.84	0.25	-	75,75,76,77	0
3	EDO	J	206	4/4	0.86	0.22	-	72,73,75,77	0

6.5 Other polymers [i](#)

There are no such residues in this entry.