



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 09:45 PM GMT

PDB ID : 1QGW
Title : CRYSTAL STRUCTURE OF PHYCOERYTHRIN 545 FROM THE MARINE CRYPTOPHYTE RHODOMONAS CS24
Authors : Harrop, S.J.; Wilk, K.E.; Hiller, R.G.; Curmi, P.M.G.
Deposited on : 1999-05-10
Resolution : 1.63 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2091 (1.66-1.62)
Ramachandran outliers	100387	2052 (1.66-1.62)
Sidechain outliers	100360	2052 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	76	 96% .
2	B	67	 94% 6%
3	C	177	 90% 5% • 5%
3	D	177	 94% 5% •

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 4788 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (CRYPTOPHYTAN PHYCOERYTHRIN (ALPHA-1 CHAIN)).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	76	Total	C	N	O	S	6	0	0
			567	348	98	117	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	4	LYZ	LYS	HYDROXYLATION	UNP Q00433
A	10	GLN	LEU	CONFLICT	UNP Q00433

- Molecule 2 is a protein called PROTEIN (CRYPTOPHYTAN PHYCOERYTHRIN (ALPHA-2 CHAIN)).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	67	Total	C	N	O	S	4	2	0
			493	303	86	98	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	4	LYZ	LYS	HYDROXYLATION	UNP Q00433

- Molecule 3 is a protein called PROTEIN (CRYPTOPHYTAN PHYCOERYTHRIN (BETA CHAIN)).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	169	Total	C	N	O	S	28	4	0
			1236	759	215	252	10			
3	D	177	Total	C	N	O	S	19	9	0
			1300	794	225	270	11			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	50	CYS	VAL	CONFLICT	UNP P27198
C	56	VAL	TYR	CONFLICT	UNP P27198
C	61	CYS	GLU	CONFLICT	UNP P27198
C	65	SER	HIS	CONFLICT	UNP P27198
C	72	MEN	ASN	METHYLATION	UNP P27198
C	73	CYS	GLU	CONFLICT	UNP P27198
D	50	CYS	VAL	CONFLICT	UNP P27198
D	56	VAL	TYR	CONFLICT	UNP P27198
D	61	CYS	GLU	CONFLICT	UNP P27198
D	65	SER	HIS	CONFLICT	UNP P27198
D	72	MEN	ASN	METHYLATION	UNP P27198
D	73	CYS	GLU	CONFLICT	UNP P27198


- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total Cl 1 1	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mg 1 1	0	0
5	D	1	Total Mg 1 1	0	0

- Molecule 6 is 15,16-DIHYDROBILIVERDIN (three-letter code: DBV) (formula: C₃₃H₃₆N₄O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			43	33	4	6		
6	B	1	Total	C	N	O	0	0
			43	33	4	6		

- Molecule 7 is PHYCOERYTHROBILIN (three-letter code: PEB) (formula: C₃₃H₄₀N₄O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	C	1	Total	C	N	O	0	0
			43	33	4	6		
7	C	1	Total	C	N	O	0	0
			43	33	4	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	C	1	Total	C	N	O	0	0
			43	33	4	6		
7	D	1	Total	C	N	O	0	0
			43	33	4	6		
7	D	1	Total	C	N	O	0	0
			43	33	4	6		
7	D	1	Total	C	N	O	0	0
			43	33	4	6		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	168	Total	O	0	0
			168	168		
8	B	135	Total	O	0	0
			135	135		
8	C	217	Total	O	0	0
			217	217		
8	D	325	Total	O	0	0
			325	325		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: PROTEIN (CRYPTOPHYTAN PHYCOERYTHRIN (ALPHA-1 CHAIN))

Chain A:  96% 



- Molecule 2: PROTEIN (CRYPTOPHYTAN PHYCOERYTHRIN (ALPHA-2 CHAIN))

Chain B:  94% 6%



- Molecule 3: PROTEIN (CRYPTOPHYTAN PHYCOERYTHRIN (BETA CHAIN))

Chain C:  90% 5% 5%



- Molecule 3: PROTEIN (CRYPTOPHYTAN PHYCOERYTHRIN (BETA CHAIN))

Chain D:  94% 5%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.03Å 82.63Å 89.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.63	Depositor
% Data completeness (in resolution range)	95.4 (20.00-1.63)	Depositor
R_{merge}	0.04	Depositor
R_{sym}	0.04	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.149 , 0.188	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4788	wwPDB-VP
Average B, all atoms (Å ²)	8.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CL, MEN, LYZ, PEB, DBV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/563	0.86	0/749
2	B	0.41	0/498	0.81	0/661
3	C	0.44	0/1258	0.90	3/1695 (0.2%)
3	D	0.45	1/1348 (0.1%)	0.88	5/1814 (0.3%)
All	All	0.44	1/3667 (0.0%)	0.88	8/4919 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	7	ARG	CD-NE	-5.34	1.37	1.46

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	91	ARG	NE-CZ-NH2	-10.33	115.13	120.30
3	D	108	ARG	NE-CZ-NH2	7.33	123.96	120.30
3	C	7	ARG	NE-CZ-NH2	7.29	123.94	120.30
3	D	1	MET	CG-SD-CE	5.97	109.75	100.20
3	D	7	ARG	CD-NE-CZ	5.55	131.37	123.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	567	0	564	3	0
2	B	493	0	511	5	2
3	C	1236	0	1231	5	2
3	D	1300	0	1300	5	0
4	C	1	0	0	0	0
5	A	1	0	0	0	0
5	D	1	0	0	0	0
6	A	43	0	33	3	0
6	B	43	0	33	4	0
7	C	129	0	110	4	0
7	D	129	0	110	3	0
8	A	168	0	0	1	1
8	B	135	0	0	1	1
8	C	217	0	0	2	4
8	D	325	0	0	2	4
All	All	4788	0	3892	28	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 28 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:115:GLU:HG3	8:C:1103:HOH:O	1.76	0.85
2:B:51:THR:HG21	8:D:1087:HOH:O	1.82	0.80
1:A:37:GLU:OE2	8:A:929:HOH:O	2.09	0.70
1:A:9:PRO:O	3:C:91:ARG:HD3	1.94	0.68
3:C:125:ASN:OD1	8:C:1115:HOH:O	2.14	0.64

The worst 5 of 7 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:34:LYS:NZ	3:C:157:ASP:OD1[4_456]	1.73	0.47
8:A:1028:HOH:O	8:D:1103:HOH:O[3_555]	1.91	0.29
8:C:1047:HOH:O	8:D:912:HOH:O[3_555]	2.03	0.17
8:B:302:HOH:O	8:C:1097:HOH:O[1_455]	2.10	0.10
2:B:34:LYS:NZ	3:C:157:ASP:CG[4_456]	2.10	0.10

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	73/76 (96%)	72 (99%)	1 (1%)	0	100	100
2	B	66/67 (98%)	64 (97%)	2 (3%)	0	100	100
3	C	168/177 (95%)	167 (99%)	1 (1%)	0	100	100
3	D	182/177 (103%)	181 (100%)	1 (0%)	0	100	100
All	All	489/497 (98%)	484 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	60/60 (100%)	60 (100%)	0	100	100
2	B	52/50 (104%)	52 (100%)	0	100	100
3	C	138/141 (98%)	136 (99%)	2 (1%)	74	52
3	D	150/141 (106%)	147 (98%)	3 (2%)	63	35
All	All	400/392 (102%)	395 (99%)	5 (1%)	76	55

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	111	ASN
3	C	127	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	D	111	ASN
3	D	127	ASN
3	D	144	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
3	C	111	ASN
3	C	127	ASN
3	D	111	ASN
3	D	127	ASN
3	D	144	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	LYZ	A	4	1	8,9,10	0.66	0	6,10,12	2.03	2 (33%)
2	LYZ	B	4	2	8,9,10	0.77	0	6,10,12	3.47	3 (50%)
3	MEN	C	72	3	7,8,9	0.63	0	5,9,11	1.26	1 (20%)
3	MEN	D	72	3	7,8,9	0.40	0	5,9,11	1.51	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LYZ	A	4	1	-	0/7/9/11	0/0/0/0
2	LYZ	B	4	2	-	0/7/9/11	0/0/0/0
3	MEN	C	72	3	-	0/6/8/10	0/0/0/0
3	MEN	D	72	3	-	0/6/8/10	0/0/0/0

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	LYZ	OH-CD-CE	-6.06	88.08	109.18
2	B	4	LYZ	OH-CD-CG	-4.07	97.69	109.29
2	B	4	LYZ	CG-CD-CE	-3.91	107.95	112.12
1	A	4	LYZ	OH-CD-CG	-3.74	98.63	109.29
3	D	72	MEN	O-C-CA	-3.31	116.88	125.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 3 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	DBV	A	219	1	32,46,46	1.86	5 (15%)	34,67,67	1.78	8 (23%)
6	DBV	B	219	2	32,46,46	1.75	5 (15%)	34,67,67	1.72	8 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	PEB	C	250	3	36,46,46	1.83	7 (19%)	38,67,67	1.67	11 (28%)
7	PEB	C	258	3	36,46,46	1.84	6 (16%)	38,67,67	1.55	8 (21%)
7	PEB	C	282	3	36,46,46	1.87	6 (16%)	38,67,67	1.68	10 (26%)
7	PEB	D	250	3	36,46,46	1.93	7 (19%)	38,67,67	1.66	10 (26%)
7	PEB	D	258	3	36,46,46	1.78	5 (13%)	38,67,67	1.44	8 (21%)
7	PEB	D	282	3	36,46,46	1.94	6 (16%)	38,67,67	1.72	11 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	DBV	A	219	1	-	0/21/74/74	0/4/4/4
6	DBV	B	219	2	-	0/21/74/74	0/4/4/4
7	PEB	C	250	3	-	2/19/74/74	0/4/4/4
7	PEB	C	258	3	-	2/19/74/74	0/4/4/4
7	PEB	C	282	3	-	2/19/74/74	0/4/4/4
7	PEB	D	250	3	-	2/19/74/74	0/4/4/4
7	PEB	D	258	3	-	2/19/74/74	0/4/4/4
7	PEB	D	282	3	-	2/19/74/74	0/4/4/4

The worst 5 of 47 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	D	250	PEB	C2A-C1A	-6.32	1.46	1.52
7	D	282	PEB	C2A-C1A	-5.59	1.46	1.52
7	C	258	PEB	C2A-C1A	-5.57	1.46	1.52
7	C	282	PEB	C2A-C1A	-5.19	1.47	1.52
7	C	250	PEB	C2A-C1A	-5.11	1.47	1.52

The worst 5 of 74 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	219	DBV	CHC-C1D-ND	-5.25	107.91	113.99
6	A	219	DBV	CAB-C3B-C4B	-3.99	122.68	127.01
6	B	219	DBV	CHC-C1D-ND	-3.94	109.43	113.99
7	D	250	PEB	CBC-CAC-C2C	-3.91	105.52	112.53
7	C	282	PEB	CHC-C4C-C3C	-3.38	124.71	130.41

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	D	250	PEB	C4A-CHA-C1B-C2B
7	D	282	PEB	C4A-CHA-C1B-C2B
7	C	258	PEB	C4A-CHA-C1B-C2B
7	D	258	PEB	C4A-CHA-C1B-C2B
7	C	250	PEB	C4A-CHA-C1B-C2B

There are no ring outliers.

6 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	219	DBV	3	0
6	B	219	DBV	4	0
7	C	258	PEB	3	0
7	C	282	PEB	1	0
7	D	258	PEB	2	0
7	D	282	PEB	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.