



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 09:45 PM GMT

PDB ID : 1QH4
Title : CRYSTAL STRUCTURE OF CHICKEN BRAIN-TYPE CREATINE KINASE AT 1.41 ANGSTROM RESOLUTION
Authors : Eder, M.; Schlattner, U.; Becker, A.; Wallimann, T.; Kabsch, W.; Fritz-Wolf, K.
Deposited on : 1999-05-11
Resolution : 1.41 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1743 (1.44-1.40)
Ramachandran outliers	100387	1698 (1.44-1.40)
Sidechain outliers	100360	1697 (1.44-1.40)
RSRZ outliers	91569	1632 (1.44-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	380	<div> <div>8%</div> <div>84%</div> <div>15%</div> <div>.</div> </div>
1	B	380	<div> <div>4%</div> <div>89%</div> <div>8%</div> <div>.</div> </div>
1	C	380	<div> <div>4%</div> <div>87%</div> <div>12%</div> <div>.</div> </div>
1	D	380	<div> <div>6%</div> <div>82%</div> <div>14%</div> <div>..</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 13559 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CREATINE KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	380	Total	C	N	O	S	0	0	0
			3005	1890	529	570	16			
1	B	380	Total	C	N	O	S	0	7	0
			3030	1907	535	571	17			
1	C	380	Total	C	N	O	S	0	6	0
			3021	1901	530	572	18			
1	D	380	Total	C	N	O	S	0	2	0
			3011	1895	529	570	17			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total	Ca	0	0
			1	1		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	360	Total O 360 360	0	0
4	B	416	Total O 416 416	0	0
4	C	447	Total O 447 447	0	0
4	D	244	Total O 244 244	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CREATINE KINASE




• Molecule 1: CREATINE KINASE




• Molecule 1: CREATINE KINASE



• Molecule 1: CREATINE KINASE





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.43 Å 175.99 Å 95.40 Å 90.00° 95.85° 90.00°	Depositor
Resolution (Å)	6.00 – 1.41 6.00 – 1.41	Depositor EDS
% Data completeness (in resolution range)	99.1 (6.00-1.41) 97.9 (6.00-1.41)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.93 (at 1.41 Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.137 , 0.188 0.144 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	13.7	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.70 , 103.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 297347 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13559	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/3067	1.27	25/4136 (0.6%)
1	B	0.57	0/3127	1.26	22/4215 (0.5%)
1	C	0.59	0/3113	1.37	33/4197 (0.8%)
1	D	0.51	0/3083	1.26	23/4157 (0.6%)
All	All	0.55	0/12390	1.29	103/16705 (0.6%)

There are no bond length outliers.

The worst 5 of 103 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	130	ARG	CD-NE-CZ	29.40	164.76	123.60
1	C	292	ARG	NE-CZ-NH1	11.51	126.06	120.30
1	B	43	ARG	NE-CZ-NH2	-11.26	114.67	120.30
1	C	43	ARG	NE-CZ-NH1	10.65	125.62	120.30
1	A	252	ARG	NE-CZ-NH2	-10.57	115.02	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3005	0	2963	27	0
1	B	3030	0	3008	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3021	0	2990	25	0
1	D	3011	0	2975	26	0
2	D	1	0	0	0	0
3	A	4	0	3	0	0
3	B	8	0	6	1	0
3	C	8	0	6	0	0
3	D	4	0	3	0	0
4	A	360	0	0	4	0
4	B	416	0	0	5	0
4	C	447	0	0	11	0
4	D	244	0	0	3	0
All	All	13559	0	11954	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 90 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:16:VAL:H	3:B:1501:ACT:H2	1.32	0.92
1:A:184:GLN:O	1:A:188:ILE:HG13	1.85	0.76
1:D:255:THR:HG22	4:D:1723:HOH:O	1.83	0.76
1:A:186:GLN:HA	1:A:189:ASP:OD2	1.87	0.75
1:C:359[B]:LEU:HG	4:C:1775:HOH:O	1.87	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	378/380 (100%)	363 (96%)	13 (3%)	2 (0%)	34 9

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	385/380 (101%)	370 (96%)	10 (3%)	5 (1%)	15	2
1	C	384/380 (101%)	372 (97%)	10 (3%)	2 (0%)	34	9
1	D	380/380 (100%)	365 (96%)	12 (3%)	3 (1%)	24	4
All	All	1527/1520 (100%)	1470 (96%)	45 (3%)	12 (1%)	24	4

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	327	THR
1	A	331	GLY
1	B	325	VAL
1	C	323	GLY
1	C	322	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	330/330 (100%)	320 (97%)	10 (3%)	48	13
1	B	337/330 (102%)	326 (97%)	11 (3%)	45	11
1	C	336/330 (102%)	333 (99%)	3 (1%)	84	62
1	D	332/330 (101%)	311 (94%)	21 (6%)	22	2
All	All	1335/1320 (101%)	1290 (97%)	45 (3%)	45	10

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	326	ASP
1	D	32	LYS
1	D	325	VAL
1	C	322	THR
1	D	68	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	286	ASN
1	B	178	ASN
1	B	318	GLN
1	A	268	ASN
1	B	296	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ACT	A	1502	-	1,3,3	1.84	0	0,3,3	0.00	-
3	ACT	B	1501	-	1,3,3	1.69	0	0,3,3	0.00	-
3	ACT	B	1503	-	1,3,3	2.26	1 (100%)	0,3,3	0.00	-
3	ACT	C	1504	-	1,3,3	1.27	0	0,3,3	0.00	-
3	ACT	C	1505	-	1,3,3	2.71	1 (100%)	0,3,3	0.00	-
3	ACT	D	1506	-	1,3,3	1.80	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACT	A	1502	-	-	0/0/0/0	0/0/0/0
3	ACT	B	1501	-	-	0/0/0/0	0/0/0/0
3	ACT	B	1503	-	-	0/0/0/0	0/0/0/0
3	ACT	C	1504	-	-	0/0/0/0	0/0/0/0
3	ACT	C	1505	-	-	0/0/0/0	0/0/0/0
3	ACT	D	1506	-	-	0/0/0/0	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1503	ACT	CH3-C	2.26	1.52	1.48
3	C	1505	ACT	CH3-C	2.71	1.52	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1501	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	380/380 (100%)	0.28	32 (8%) 14 11	10, 25, 54, 103	0
1	B	380/380 (100%)	-0.20	17 (4%) 37 33	9, 19, 38, 101	0
1	C	380/380 (100%)	-0.25	14 (3%) 45 42	10, 18, 36, 98	0
1	D	380/380 (100%)	0.23	24 (6%) 23 20	9, 29, 58, 102	0
All	All	1520/1520 (100%)	0.01	87 (5%) 27 24	9, 21, 51, 103	0

The worst 5 of 87 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	328	ALA	23.5
1	B	325	VAL	19.6
1	C	325	VAL	18.2
1	A	330	VAL	17.9
1	A	328	ALA	15.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	ACT	D	1506	4/4	0.95	0.10	1.81	26,31,31,33	0
3	ACT	C	1505	4/4	0.91	0.11	1.36	49,52,52,58	0
3	ACT	B	1503	4/4	0.96	0.09	0.65	33,39,40,45	0
3	ACT	A	1502	4/4	0.97	0.07	0.37	27,32,34,41	0
3	ACT	C	1504	4/4	0.97	0.07	0.35	19,27,32,35	0
2	CA	D	382	1/1	1.00	0.02	-2.14	12,12,12,12	0
3	ACT	B	1501	4/4	0.88	0.16	-	27,34,40,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.