



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:45 PM GMT

PDB ID : 1QJA
Title : 14-3-3 ZETA/PHOSHOPEPTIDE COMPLEX (MODE 2)
Authors : Rittinger, K.; Budman, J.; Xu, J.; Volinia, S.; Cantley, L.C.; Smerdon, S.J.;
Gamblin, S.J.; Yaffe, M.B.
Deposited on : 1999-06-23
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	245	 66% 18% • • 11%
1	B	245	 65% 22% • • 8%
2	Q	8	 88% 13%
2	R	8	 75% 25%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3948 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 14-3-3 PROTEIN ZETA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	0	0
			1745	1098	291	346	10			
1	B	225	Total	C	N	O	S	0	0	0
			1811	1138	304	359	10			

- Molecule 2 is a protein called PHOSPHOPEPTIDE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	Q	8	Total	C	N	O	P	0	0	0
			71	44	13	13	1			
2	R	8	Total	C	N	O	P	0	0	0
			71	44	13	13	1			

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	110	Total	O	0	0
			110	110		
3	B	122	Total	O	0	0
			122	122		
3	Q	9	Total	O	0	0
			9	9		
3	R	9	Total	O	0	0
			9	9		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: 14-3-3 PROTEIN ZETA




• Molecule 1: 14-3-3 PROTEIN ZETA



• Molecule 2: PHOSHOPEPTIDE



• Molecule 2: PHOSHOPEPTIDE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.38 Å 70.96 Å 130.56 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.00	Depositor
% Data completeness (in resolution range)	95.2 (15.00-2.00)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.214 , 0.275	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3948	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/1768	1.61	29/2377 (1.2%)
1	B	0.60	0/1835	1.46	22/2465 (0.9%)
2	Q	0.56	0/62	1.32	1/82 (1.2%)
2	R	0.72	0/62	1.23	0/82
All	All	0.61	0/3727	1.53	52/5006 (1.0%)

There are no bond length outliers.

All (52) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	41	ARG	CD-NE-CZ	22.26	154.77	123.60
1	A	80	ARG	CD-NE-CZ	17.25	147.75	123.60
1	B	83	ARG	NE-CZ-NH2	-16.71	111.94	120.30
1	A	160	MET	C-N-CA	16.06	161.86	121.70
1	A	56	ARG	NE-CZ-NH1	14.49	127.55	120.30
1	B	83	ARG	CD-NE-CZ	13.87	143.02	123.60
1	B	83	ARG	NE-CZ-NH1	10.47	125.54	120.30
1	A	161	GLN	N-CA-CB	10.28	129.10	110.60
1	A	160	MET	O-C-N	-9.61	107.32	122.70
1	B	128	TYR	CB-CG-CD2	-9.22	115.47	121.00
1	A	160	MET	CA-C-O	9.10	139.20	120.10
1	B	80	ARG	NE-CZ-NH2	-8.99	115.81	120.30
1	A	60	ARG	CD-NE-CZ	8.47	135.46	123.60
1	B	60	ARG	CD-NE-CZ	8.23	135.12	123.60
1	A	161	GLN	CB-CA-C	-8.20	93.99	110.40
1	A	56	ARG	NE-CZ-NH2	-8.18	116.21	120.30
1	B	140	GLY	CA-C-O	7.92	134.86	120.60
1	B	78	MET	CA-CB-CG	7.68	126.36	113.30
1	B	60	ARG	NE-CZ-NH1	7.63	124.11	120.30
1	A	127	ARG	NE-CZ-NH2	-7.57	116.52	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	20	ASP	CB-CG-OD1	7.28	124.86	118.30
1	A	60	ARG	NE-CZ-NH1	7.16	123.88	120.30
1	A	124	ASP	CB-CG-OD1	7.10	124.69	118.30
1	B	20	ASP	CB-CG-OD2	-6.88	112.11	118.30
1	A	90	LEU	CA-CB-CG	6.66	130.62	115.30
1	A	127	ARG	NE-CZ-NH1	6.66	123.63	120.30
1	A	83	ARG	NE-CZ-NH2	-6.39	117.11	120.30
1	A	55	ARG	NE-CZ-NH2	-6.37	117.11	120.30
1	A	161	GLN	CA-CB-CG	6.26	127.18	113.40
1	A	180	GLU	CA-CB-CG	6.22	127.08	113.40
1	A	19	TYR	CB-CG-CD1	-6.18	117.29	121.00
1	B	140	GLY	O-C-N	-6.13	112.89	122.70
1	B	124	ASP	CB-CG-OD1	6.12	123.81	118.30
2	Q	3	ARG	NE-CZ-NH1	6.10	123.35	120.30
1	B	149	TYR	CB-CG-CD2	-6.05	117.37	121.00
1	A	128	TYR	CB-CG-CD1	6.00	124.60	121.00
1	B	96	ASP	CB-CG-OD1	5.97	123.68	118.30
1	A	222	ARG	CD-NE-CZ	5.93	131.90	123.60
1	A	161	GLN	O-C-N	5.87	132.26	121.10
1	B	149	TYR	CB-CG-CD1	5.73	124.44	121.00
1	B	57	SER	N-CA-CB	5.72	119.08	110.50
1	A	204	ASP	CB-CG-OD1	5.65	123.39	118.30
1	A	124	ASP	CB-CG-OD2	-5.62	113.24	118.30
1	B	18	ARG	NE-CZ-NH2	-5.41	117.59	120.30
1	B	128	TYR	CB-CG-CD1	5.35	124.21	121.00
1	B	80	ARG	CD-NE-CZ	5.34	131.07	123.60
1	A	63	SER	N-CA-CB	5.28	118.42	110.50
1	A	80	ARG	NE-CZ-NH2	5.26	122.93	120.30
1	A	20	ASP	CB-CG-OD1	5.22	123.00	118.30
1	B	139	LYS	CA-C-N	5.13	126.45	116.20
1	A	169	GLY	CA-C-O	5.03	129.65	120.60
1	B	48	TYR	CB-CG-CD1	-5.01	117.99	121.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1745	0	1733	28	1
1	B	1811	0	1811	38	1
2	Q	71	0	65	1	0
2	R	71	0	65	2	0
3	A	110	0	0	7	0
3	B	122	0	0	1	0
3	Q	9	0	0	0	0
3	R	9	0	0	0	0
All	All	3948	0	3674	64	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:8:GLN:HG3	3:A:2001:HOH:O	1.42	1.15
1:A:102:GLU:HG2	3:A:2054:HOH:O	1.59	1.01
1:A:3:LYS:HD2	1:A:29:VAL:HG13	1.60	0.83
1:B:164:HIS:HD2	1:B:166:ILE:H	1.33	0.76
1:A:42:ASN:HB3	3:A:2027:HOH:O	1.86	0.75
1:A:222:ARG:HG3	3:A:2106:HOH:O	1.88	0.73
1:B:26:MET:O	1:B:30:THR:HG23	1.93	0.68
1:B:1:MET:HG3	1:B:6:LEU:HG	1.75	0.66
1:B:133:ALA:HB3	1:B:138:LYS:N	2.13	0.64
1:A:66:GLU:OE1	1:A:83:ARG:HD3	1.98	0.63
1:B:139:LYS:HG3	1:B:140:GLY:N	2.14	0.63
1:B:30:THR:HG22	1:B:36:LEU:HD11	1.81	0.62
1:B:49:LYS:HE2	1:B:128:TYR:OH	1.99	0.61
1:B:3:LYS:HG2	1:B:32:GLN:OE1	2.01	0.61
1:A:160:MET:O	1:A:167:ARG:NH1	2.33	0.61
1:B:113:GLU:HA	1:B:160:MET:HE2	1.82	0.60
1:B:153:PHE:HA	1:B:170:LEU:HD21	1.85	0.58
1:B:60:ARG:NH2	3:B:2040:HOH:O	2.36	0.58
1:A:106:ILE:N	1:A:107:PRO:HD2	2.20	0.57
1:B:113:GLU:HA	1:B:160:MET:CE	2.35	0.56
1:A:227:LEU:HD23	2:R:5:TYR:HB3	1.88	0.56
1:B:164:HIS:CD2	1:B:166:ILE:H	2.21	0.56
1:A:91:ARG:NH1	3:A:2048:HOH:O	2.27	0.55
1:B:9:LYS:HG2	1:B:25:CYS:SG	2.47	0.54
1:A:8:GLN:HG2	3:A:2004:HOH:O	2.06	0.54
1:A:211:TYR:CE1	1:A:212:LYS:HG2	2.43	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:73:GLU:HG3	1:B:74:LYS:N	2.23	0.53
1:A:119:LEU:HB3	1:A:152:ALA:HB2	1.91	0.53
1:A:9:LYS:HD2	1:A:25:CYS:SG	2.49	0.53
1:B:116:VAL:HG21	1:B:160:MET:HE3	1.91	0.52
1:B:143:ASP:O	1:B:147:GLN:HG3	2.11	0.51
1:A:180:GLU:OE1	2:R:3:ARG:NH1	2.45	0.49
1:B:59:TRP:NE1	1:B:87:GLU:HG3	2.27	0.49
1:A:119:LEU:CB	1:A:152:ALA:HB2	2.43	0.48
1:B:59:TRP:CD1	1:B:87:GLU:HG3	2.49	0.47
1:A:18:ARG:NH2	1:B:89:GLU:OE2	2.39	0.47
1:B:117:PHE:HD2	1:B:166:ILE:HD12	1.80	0.47
1:A:109:ALA:O	1:A:110:SER:CB	2.62	0.47
1:B:164:HIS:HD2	1:B:166:ILE:N	2.08	0.47
1:B:73:GLU:HG3	1:B:74:LYS:HG3	1.98	0.46
1:B:56:ARG:HG3	1:B:90:LEU:HD11	1.97	0.46
1:B:55:ARG:CB	1:B:90:LEU:HD13	2.46	0.46
1:B:55:ARG:HB3	1:B:90:LEU:HD13	1.98	0.46
1:B:106:ILE:N	1:B:107:PRO:HD2	2.31	0.46
1:A:222:ARG:HG3	1:A:222:ARG:HH11	1.81	0.45
1:A:113:GLU:OE1	3:A:2057:HOH:O	2.20	0.45
1:B:59:TRP:CE2	1:B:132:VAL:HG12	2.51	0.45
1:A:9:LYS:HE2	1:B:82:TYR:CD1	2.51	0.45
1:A:113:GLU:HA	1:A:160:MET:CE	2.47	0.45
1:B:60:ARG:NH1	2:Q:3:ARG:HB2	2.32	0.44
1:A:206:LEU:HD21	1:A:214:SER:CB	2.48	0.44
1:B:37:SER:OG	1:B:38:ASN:N	2.50	0.44
1:A:180:GLU:HG3	1:A:181:ILE:HG12	2.00	0.44
1:B:119:LEU:HB2	1:B:152:ALA:HB2	2.00	0.43
1:B:27:LYS:O	1:B:31:GLU:HG3	2.18	0.43
1:B:73:GLU:CG	1:B:74:LYS:N	2.82	0.43
1:A:163:THR:HB	1:A:206:LEU:HG	2.00	0.43
1:B:153:PHE:CA	1:B:170:LEU:HD21	2.48	0.42
1:A:38:ASN:HD22	1:A:41:ARG:NH1	2.17	0.42
1:B:120:LYS:HD2	1:B:170:LEU:HA	2.01	0.41
1:B:126:TYR:HB3	1:B:145:SER:HB2	2.02	0.41
1:A:193:LYS:HA	1:A:225:LEU:HD11	2.02	0.41
1:B:73:GLU:HG3	1:B:74:LYS:H	1.86	0.41
1:A:115:LYS:HG2	1:A:155:ILE:HD13	2.03	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:161:GLN:N	1:B:73:GLU:OE2[3_545]	1.80	0.40

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/245 (86%)	207 (98%)	3 (1%)	1 (0%)	34	26
1	B	221/245 (90%)	216 (98%)	3 (1%)	2 (1%)	21	13
2	Q	5/8 (62%)	5 (100%)	0	0	100	100
2	R	5/8 (62%)	5 (100%)	0	0	100	100
All	All	442/506 (87%)	433 (98%)	6 (1%)	3 (1%)	26	19

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	140	GLY
1	A	109	ALA
1	B	73	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	190/209 (91%)	177 (93%)	13 (7%)	20	13
1	B	197/209 (94%)	183 (93%)	14 (7%)	18	12

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	Q	6/6 (100%)	6 (100%)	0	100	100
2	R	6/6 (100%)	6 (100%)	0	100	100
All	All	399/430 (93%)	372 (93%)	27 (7%)	20	13

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	9	LYS
1	A	41	ARG
1	A	78	MET
1	A	80	ARG
1	A	90	LEU
1	A	110	SER
1	A	137	ASP
1	A	147	GLN
1	A	158	LYS
1	A	174	PHE
1	A	211	TYR
1	A	222	ARG
1	B	1	MET
1	B	8	GLN
1	B	37	SER
1	B	57	SER
1	B	70	GLU
1	B	78	MET
1	B	100	LEU
1	B	111	GLN
1	B	138	LYS
1	B	139	LYS
1	B	157	LYS
1	B	174	PHE
1	B	208	GLU
1	B	225	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	8	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	38	ASN
1	B	164	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SEP	Q	7	2	8,9,10	3.55	1 (12%)	8,12,14	4.69	5 (62%)
2	SEP	R	7	2	8,9,10	3.32	1 (12%)	8,12,14	5.40	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SEP	Q	7	2	-	0/6/8/10	0/0/0/0
2	SEP	R	7	2	-	0/6/8/10	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	R	7	SEP	P-OG	9.08	1.90	1.60
2	Q	7	SEP	P-OG	9.77	1.93	1.60

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	R	7	SEP	OG-CB-CA	-14.68	95.75	108.27
2	Q	7	SEP	OG-CB-CA	-11.33	98.61	108.27
2	Q	7	SEP	O-C-CA	-4.28	114.33	125.49
2	Q	7	SEP	O2P-P-OG	-3.92	95.29	106.56
2	Q	7	SEP	OG-P-O1P	2.13	112.57	107.14
2	Q	7	SEP	O3P-P-OG	2.48	113.70	106.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.