



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:45 PM GMT

PDB ID : 1QJE
Title : ISOPENICILLIN N SYNTHASE FROM ASPERGILLUS NIDULANS (IP1 - FE COMPLEX)
Authors : Burzlaff, N.I.; Clifton, I.J.; Rutledge, P.J.; Roach, P.L.; Adlington, R.M.; Baldwin, J.E.
Deposited on : 1999-06-23
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2337 (1.40-1.32)
Ramachandran outliers	100387	2280 (1.40-1.32)
Sidechain outliers	100360	2279 (1.40-1.32)
RSRZ outliers	91569	2199 (1.40-1.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	<div> <div>68%</div> <div>83%</div> <div>13%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACV	A	1334[B]	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3181 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ISOPENICILLIN N SYNTHASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	327	2635	1684	444	502	5	0	1	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

- Molecule 3 is ISOPENICILLIN N (three-letter code: IP1) (formula: C₁₄H₂₁N₃O₆S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	1
			24	14	3	6	1		

- Molecule 4 is L-D-(A-AMINOADIPOYL)-L-CYSTEINYL-D-VALINE (three-letter code: ACV) (formula: $C_{14}H_{25}N_3O_6S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	1
			24	14	3	6	1		

- Molecule 5 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total 1	Fe 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	492	Total 492	O 492	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ISOPENICILLIN N SYNTHASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	46.67Å 71.91Å 101.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 1.35 23.34 – 1.35	Depositor EDS
% Data completeness (in resolution range)	92.2 (25.00-1.35) 91.2 (23.34-1.35)	Depositor EDS
R_{merge}	0.01	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.16 (at 1.35Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.131 , 0.187 0.371 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	12.0	Xtriage
Anisotropy	0.371	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.47 , 81.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 69132 reflections	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	3181	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IP1, FE2, ACV, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.04	3/2715 (0.1%)	1.84	64/3694 (1.7%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	309	GLU	CD-OE2	7.31	1.33	1.25
1	A	331	THR	C-OXT	7.27	1.37	1.23
1	A	306	SER	CB-OG	-5.61	1.34	1.42

All (64) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	173	ARG	CD-NE-CZ	26.79	161.10	123.60
1	A	300	ARG	CD-NE-CZ	19.52	150.93	123.60
1	A	116	ARG	NE-CZ-NH2	-15.96	112.32	120.30
1	A	308	ARG	NE-CZ-NH1	13.06	126.83	120.30
1	A	308	ARG	NE-CZ-NH2	-12.35	114.13	120.30
1	A	249	TYR	OH-CZ-CE2	12.07	152.70	120.10
1	A	72	ASP	CB-CG-OD1	12.04	129.13	118.30
1	A	249	TYR	CB-CG-CD1	12.00	128.20	121.00
1	A	76	ARG	NE-CZ-NH1	11.87	126.23	120.30
1	A	141	PHE	CB-CG-CD1	-10.67	113.33	120.80
1	A	141	PHE	CB-CG-CD2	10.10	127.87	120.80
1	A	249	TYR	CZ-CE2-CD2	9.78	128.60	119.80
1	A	37	ARG	NE-CZ-NH2	-9.46	115.57	120.30
1	A	76	ARG	NE-CZ-NH2	-8.69	115.95	120.30
1	A	173	ARG	NE-CZ-NH1	8.63	124.61	120.30
1	A	291	ASP	CB-CG-OD1	8.61	126.05	118.30
1	A	249	TYR	CE1-CZ-OH	-8.32	97.62	120.10
1	A	90	TYR	CB-CG-CD1	8.21	125.92	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	6	LYS	CG-CD-CE	8.19	136.47	111.90
1	A	145	TYR	CB-CG-CD2	7.86	125.72	121.00
1	A	116	ARG	NE-CZ-NH1	7.66	124.13	120.30
1	A	87	ARG	NE-CZ-NH1	7.61	124.11	120.30
1	A	275	VAL	CA-CB-CG1	-7.47	99.69	110.90
1	A	68	GLU	OE1-CD-OE2	-7.41	114.41	123.30
1	A	141	PHE	CZ-CE2-CD2	-7.32	111.32	120.10
1	A	188	ARG	NE-CZ-NH2	-7.32	116.64	120.30
1	A	224	TYR	CB-CG-CD1	-7.15	116.71	121.00
1	A	243	GLU	CB-CG-CD	6.92	132.89	114.20
1	A	116	ARG	CD-NE-CZ	-6.89	113.95	123.60
1	A	226	SER	CB-CA-C	-6.86	97.06	110.10
1	A	150	PHE	CG-CD1-CE1	6.64	128.10	120.80
1	A	239	TYR	CG-CD1-CE1	-6.53	116.08	121.30
1	A	81	GLU	OE1-CD-OE2	-6.40	115.62	123.30
1	A	237	ALA	CB-CA-C	-6.39	100.51	110.10
1	A	249	TYR	CE1-CZ-CE2	-6.37	109.62	119.80
1	A	141	PHE	CG-CD2-CE2	6.36	127.79	120.80
1	A	6	LYS	CB-CG-CD	6.31	128.01	111.60
1	A	84	ASP	CB-CG-OD2	-6.26	112.67	118.30
1	A	309	GLU	CB-CG-CD	6.24	131.05	114.20
1	A	249	TYR	CB-CG-CD2	-6.11	117.34	121.00
1	A	20	ASP	CB-CG-OD2	-5.96	112.93	118.30
1	A	84	ASP	CB-CG-OD1	5.94	123.64	118.30
1	A	72	ASP	CB-CG-OD2	-5.92	112.97	118.30
1	A	173	ARG	NH1-CZ-NH2	-5.64	113.20	119.40
1	A	245	ASP	CB-CG-OD2	5.60	123.34	118.30
1	A	264	TYR	CB-CG-CD1	5.57	124.34	121.00
1	A	68	GLU	CG-CD-OE1	5.53	129.36	118.30
1	A	21	ASP	CB-CG-OD2	5.51	123.26	118.30
1	A	265	TYR	CB-CG-CD1	5.51	124.30	121.00
1	A	271	ARG	NE-CZ-NH1	5.49	123.04	120.30
1	A	176	LYS	CD-CE-NZ	5.46	124.25	111.70
1	A	13	ASP	CB-CG-OD2	5.38	123.14	118.30
1	A	171	PHE	CD1-CE1-CZ	-5.37	113.65	120.10
1	A	201	LYS	CD-CE-NZ	5.36	124.02	111.70
1	A	224	TYR	CB-CG-CD2	5.27	124.16	121.00
1	A	197	GLU	OE1-CD-OE2	-5.26	116.99	123.30
1	A	239	TYR	CD1-CE1-CZ	5.24	124.52	119.80
1	A	205	ASP	CB-CG-OD1	5.22	123.00	118.30
1	A	266	LYS	CD-CE-NZ	5.17	123.60	111.70
1	A	171	PHE	CB-CG-CD2	-5.15	117.19	120.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	302	PRO	O-C-N	5.10	130.86	122.70
1	A	237	ALA	N-CA-CB	5.08	117.22	110.10
1	A	30	GLN	CG-CD-OE1	-5.07	111.46	121.60
1	A	189	TYR	CD1-CE1-CZ	-5.06	115.25	119.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2635	0	2518	11	0
2	A	5	0	0	1	0
3	A	24	0	19	0	0
4	A	24	0	21	2	0
5	A	1	0	0	0	0
6	A	492	0	0	8	2
All	All	3181	0	2558	12	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:ALA:HB1	6:A:2380:HOH:O	2.01	0.60
1:A:140:ASP:HB3	6:A:2482:HOH:O	2.09	0.52
4:A:1334[B]:ACV:H331	6:A:2358:HOH:O	2.11	0.50
1:A:309:GLU:HG2	6:A:2454:HOH:O	2.12	0.48
1:A:141:PHE:HA	2:A:1332:SO4:O1	2.15	0.46
1:A:80:LYS:NZ	6:A:2166:HOH:O	2.50	0.45
1:A:116:ARG:NH2	1:A:291:ASP:OD2	2.50	0.44
1:A:213:TRP:CZ2	1:A:327:LYS:HE3	2.54	0.43
1:A:68:GLU:HB3	6:A:2147:HOH:O	2.19	0.43
1:A:166:LYS:NZ	6:A:2303:HOH:O	2.52	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:245:ASP:HB2	6:A:2011:HOH:O	2.20	0.42
1:A:214:HIS:NE2	4:A:1334[B]:ACV:H162	2.36	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:2394:HOH:O	6:A:2482:HOH:O[4_455]	1.80	0.40
6:A:2101:HOH:O	6:A:2220:HOH:O[4_455]	1.91	0.29

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/331 (98%)	317 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/284 (99%)	279 (99%)	3 (1%)	80	53

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	SER
1	A	91	TYR
1	A	306	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	225	GLN
1	A	230	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	1332	-	4,4,4	0.82	0	6,6,6	1.05	0
3	IP1	A	1333[A]	-	19,25,25	0.79	1 (5%)	27,38,38	2.06	11 (40%)
4	ACV	A	1334[B]	5	17,23,23	0.55	0	20,30,30	2.66	9 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	1332	-	-	0/0/0/0	0/0/0/0
3	IP1	A	1333[A]	-	-	0/10/49/49	0/2/2/2
4	ACV	A	1334[B]	5	-	0/24/32/32	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1333[A]	IP1	C30-N29	-2.01	1.45	1.47

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1334[B]	ACV	C33-C32-C30	-6.69	105.19	111.30
4	A	1334[B]	ACV	C16-C12-N11	-5.41	103.80	111.40
3	A	1333[A]	IP1	S17-C16-N29	-3.72	100.05	105.09
3	A	1333[A]	IP1	C13-C12-N11	-3.04	106.51	115.36
3	A	1333[A]	IP1	O18-C13-N29	-2.82	127.58	131.64
3	A	1333[A]	IP1	C32-C30-N29	-2.45	102.77	106.45
4	A	1334[B]	ACV	O15-C10-N11	-2.39	118.95	123.01
4	A	1334[B]	ACV	C16-C12-C13	-2.27	104.54	109.66
3	A	1333[A]	IP1	C30-N29-C13	-2.17	120.89	125.99
4	A	1334[B]	ACV	O18-C13-N29	-2.11	118.79	122.93
3	A	1333[A]	IP1	C33-C32-C30	-2.01	107.06	111.56
3	A	1333[A]	IP1	O18-C13-C12	2.15	142.41	136.37
3	A	1333[A]	IP1	C37-C32-C33	2.59	115.02	110.88
3	A	1333[A]	IP1	C7-C10-N11	2.67	120.19	115.83
4	A	1334[B]	ACV	C37-C32-C33	2.73	118.70	110.67
4	A	1334[B]	ACV	C12-N11-C10	2.86	128.88	121.58
3	A	1333[A]	IP1	C37-C32-S17	2.96	114.12	109.17
4	A	1334[B]	ACV	C37-C32-C30	3.00	114.04	111.30
4	A	1334[B]	ACV	O18-C13-C12	3.16	127.35	120.36
3	A	1333[A]	IP1	C16-N29-C30	4.72	122.15	117.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1332	SO4	1	0
4	A	1334[B]	ACV	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	327/331 (98%)	2.90	224 (68%) 0 0	7, 14, 28, 41	0

All (224) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	325	ILE	12.2
1	A	239	TYR	8.6
1	A	237	ALA	8.4
1	A	322	VAL	8.1
1	A	96	GLY	7.3
1	A	238	GLY	6.9
1	A	323	SER	6.9
1	A	326	ASN	6.4
1	A	319	ASN	6.3
1	A	324	LEU	6.3
1	A	198	ALA	6.2
1	A	236	ALA	5.9
1	A	94	ILE	5.8
1	A	213	TRP	5.7
1	A	71	TRP	5.7
1	A	261	THR	5.6
1	A	327	LYS	5.6
1	A	256	TYR	5.5
1	A	293	VAL	5.5
1	A	162	LEU	5.4
1	A	307	ASP	5.3
1	A	300	ARG	5.3
1	A	235	THR	5.2
1	A	207	THR	5.2
1	A	295	ASP	5.1
1	A	92	LEU	5.1
1	A	328	ASN	5.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	164	LEU	5.1
1	A	220	ILE	5.0
1	A	264	TYR	4.9
1	A	316	TYR	4.9
1	A	193	ASP	4.9
1	A	297	PHE	4.8
1	A	161	ALA	4.8
1	A	306	SER	4.7
1	A	288	LEU	4.7
1	A	202	THR	4.6
1	A	313	TYR	4.6
1	A	294	ILE	4.5
1	A	250	LEU	4.5
1	A	65	ILE	4.5
1	A	9	VAL	4.5
1	A	189	TYR	4.4
1	A	285	PHE	4.4
1	A	265	TYR	4.4
1	A	206	GLY	4.4
1	A	156	LEU	4.4
1	A	19	GLY	4.3
1	A	86	VAL	4.3
1	A	83	GLN	4.3
1	A	91	TYR	4.3
1	A	17	LEU	4.3
1	A	18	PHE	4.3
1	A	275	VAL	4.3
1	A	308	ARG	4.3
1	A	205	ASP	4.2
1	A	29	ALA	4.2
1	A	296	PRO	4.2
1	A	320	GLY	4.2
1	A	23	ALA	4.1
1	A	201	LYS	4.1
1	A	160	TYR	4.1
1	A	20	ASP	4.0
1	A	39	THR	4.0
1	A	43	TYR	4.0
1	A	171	PHE	4.0
1	A	211	PHE	4.0
1	A	302	PRO	4.0
1	A	311	LEU	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	157	LEU	3.9
1	A	192	LEU	3.9
1	A	95	PRO	3.9
1	A	249	TYR	3.9
1	A	304	GLY	3.9
1	A	117	ILE	3.9
1	A	195	TYR	3.8
1	A	258	ALA	3.8
1	A	272	VAL	3.8
1	A	78	TYR	3.8
1	A	191	TYR	3.8
1	A	253	CYS	3.8
1	A	175	PHE	3.7
1	A	45	VAL	3.7
1	A	228	VAL	3.7
1	A	260	LEU	3.7
1	A	254	GLY	3.7
1	A	170	PHE	3.7
1	A	77	ALA	3.6
1	A	231	LEU	3.6
1	A	186	LEU	3.6
1	A	181	LEU	3.6
1	A	263	ASN	3.6
1	A	329	GLY	3.5
1	A	32	ILE	3.5
1	A	318	GLN	3.5
1	A	310[A]	PRO	3.5
1	A	305	LYS	3.5
1	A	185	VAL	3.5
1	A	233	VAL	3.5
1	A	73	LEU	3.5
1	A	309	GLU	3.5
1	A	204	ALA	3.5
1	A	290	TYR	3.5
1	A	141	PHE	3.5
1	A	243	GLU	3.4
1	A	103	PHE	3.4
1	A	222	VAL	3.4
1	A	242	ILE	3.4
1	A	262	ASN	3.4
1	A	42	PHE	3.3
1	A	321	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	12	ILE	3.3
1	A	106	LEU	3.3
1	A	163	ALA	3.3
1	A	203	ALA	3.3
1	A	5	SER	3.3
1	A	110	PHE	3.2
1	A	138	PHE	3.2
1	A	274	TRP	3.2
1	A	40	GLY	3.2
1	A	121	THR	3.2
1	A	197	GLU	3.2
1	A	59	LYS	3.2
1	A	35	ALA	3.2
1	A	129	TRP	3.2
1	A	314	GLY	3.1
1	A	49	ILE	3.1
1	A	75	ILE	3.1
1	A	187	ILE	3.1
1	A	269	ILE	3.1
1	A	184	VAL	3.1
1	A	76	ARG	3.1
1	A	315	ASP	3.1
1	A	331	THR	3.1
1	A	24	ALA	3.1
1	A	28	VAL	3.0
1	A	200	ILE	3.0
1	A	317	LEU	3.0
1	A	224	TYR	3.0
1	A	119	ALA	3.0
1	A	217	VAL	3.0
1	A	44	ALA	3.0
1	A	209	LEU	3.0
1	A	221	THR	3.0
1	A	100	VAL	3.0
1	A	283	PRO	2.9
1	A	26	MET	2.9
1	A	90	TYR	2.9
1	A	82	HIS	2.9
1	A	199	ALA	2.9
1	A	150	PHE	2.9
1	A	284	PHE	2.9
1	A	34	ALA	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	88	ALA	2.8
1	A	131	ASP	2.8
1	A	169	ASN	2.8
1	A	166	LYS	2.8
1	A	61	PHE	2.8
1	A	277	ALA	2.8
1	A	286	VAL	2.8
1	A	303	ASN	2.8
1	A	41	PHE	2.8
1	A	128	VAL	2.8
1	A	282	LEU	2.8
1	A	123	THR	2.8
1	A	251	ILE	2.8
1	A	16	PRO	2.7
1	A	299	PRO	2.7
1	A	22	GLN	2.7
1	A	247	THR	2.7
1	A	194	PRO	2.6
1	A	167	GLU	2.6
1	A	330	GLN	2.6
1	A	7	ALA	2.6
1	A	105	TYR	2.6
1	A	301	GLU	2.6
1	A	196	PRO	2.6
1	A	151	GLY	2.6
1	A	74	ALA	2.5
1	A	270	HIS	2.5
1	A	298	ASP	2.5
1	A	142	ALA	2.5
1	A	133	THR	2.5
1	A	146	TYR	2.5
1	A	11	LYS	2.5
1	A	38	ASP	2.5
1	A	126	VAL	2.5
1	A	182	ALA	2.4
1	A	219	LEU	2.4
1	A	97	LYS	2.4
1	A	180	THR	2.4
1	A	67	PRO	2.4
1	A	104	CYS	2.4
1	A	173	ARG	2.4
1	A	227	ASN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	37	ARG	2.4
1	A	6	LYS	2.3
1	A	14	VAL	2.3
1	A	33	ASP	2.3
1	A	214	HIS	2.3
1	A	165	GLY	2.3
1	A	102	SER	2.3
1	A	108	PRO	2.3
1	A	244	ALA	2.3
1	A	111	THR	2.2
1	A	21	ASP	2.2
1	A	230	ASN	2.2
1	A	79	ASN	2.2
1	A	218	SER	2.2
1	A	147	TRP	2.2
1	A	54	LEU	2.2
1	A	99	ALA	2.2
1	A	80	LYS	2.1
1	A	85	GLN	2.1
1	A	152	LEU	2.1
1	A	257	MET	2.1
1	A	232	GLN	2.1
1	A	72	ASP	2.1
1	A	58	THR	2.1
1	A	229	GLN	2.0
1	A	145	TYR	2.0
1	A	312	SER	2.0
1	A	240	GLN	2.0
1	A	98	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	ACV	A	1334[B]	24/24	0.70	0.25	2.11	11,17,19,19	24
3	IP1	A	1333[A]	24/24	0.73	0.20	0.74	11,18,21,27	24
5	FE2	A	1335	1/1	0.88	0.12	-	24,24,24,24	0
2	SO4	A	1332	5/5	0.79	0.27	-	39,44,47,61	0

6.5 Other polymers

There are no such residues in this entry.