



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 07:05 PM GMT

PDB ID : 4QJY
Title : Crystal structure of native Ara127N, a GH127 beta-L-arabinofuranosidase from *Geobacillus Stearothermophilus* T6
Authors : Lansky, S.; Salama, R.; Dann, R.; Shner, I.; Manjasetty, B.; Belrhali, H.; Shoham, Y.; Shoham, G.
Deposited on : 2014-06-05
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	648	 3% 85% 12% ..
1	B	648	 4% 83% 13% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	A	703	-	-	-	X
2	ACT	B	701	-	-	-	X
2	ACT	B	702	-	-	-	X
2	ACT	B	704	-	-	-	X
2	ACT	B	705	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11013 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GH127 beta-L-arabinofuranosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	637	Total	C	N	O	S	0	5	0
			5187	3318	885	956	28			
1	B	637	Total	C	N	O	S	0	3	0
			5178	3312	885	954	27			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	HIS	-	EXPRESSION TAG	UNP B3EYN9
A	-2	HIS	-	EXPRESSION TAG	UNP B3EYN9
A	-1	HIS	-	EXPRESSION TAG	UNP B3EYN9
A	0	HIS	-	EXPRESSION TAG	UNP B3EYN9
A	1	HIS	-	EXPRESSION TAG	UNP B3EYN9
A	2	HIS	-	EXPRESSION TAG	UNP B3EYN9
B	-3	HIS	-	EXPRESSION TAG	UNP B3EYN9
B	-2	HIS	-	EXPRESSION TAG	UNP B3EYN9
B	-1	HIS	-	EXPRESSION TAG	UNP B3EYN9
B	0	HIS	-	EXPRESSION TAG	UNP B3EYN9
B	1	HIS	-	EXPRESSION TAG	UNP B3EYN9
B	2	HIS	-	EXPRESSION TAG	UNP B3EYN9

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	324	Total	O	0	0
			324	324		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	280	Total 280	O 280	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GH127 beta-L-arabinofuranosidase



- Molecule 1: GH127 beta-L-arabinofuranosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.54Å 118.10Å 174.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.89 – 2.29 29.87 – 2.29	Depositor EDS
% Data completeness (in resolution range)	99.5 (29.89-2.29) 99.6 (29.87-2.29)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.150 , 0.214 0.160 , 0.217	Depositor DCC
R_{free} test set	3122 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	29.6	Xtriage
Anisotropy	0.740	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 61437 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11013	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.81	2/5331 (0.0%)	0.86	6/7235 (0.1%)
1	B	0.77	0/5316	0.89	11/7215 (0.2%)
All	All	0.79	2/10647 (0.0%)	0.88	17/14450 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	1
All	All	0	4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	183	ASP	CB-CG	5.56	1.63	1.51
1	A	290	TRP	CB-CG	-5.43	1.40	1.50

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	174	ARG	NE-CZ-NH1	7.02	123.81	120.30
1	A	480	ARG	NE-CZ-NH1	6.61	123.60	120.30
1	A	530	ARG	NE-CZ-NH2	-6.60	117.00	120.30
1	A	480	ARG	NE-CZ-NH2	-6.32	117.14	120.30
1	A	356	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	B	512	ARG	NE-CZ-NH1	5.64	123.12	120.30
1	B	316	ASP	CB-CG-OD1	5.64	123.38	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	453	GLN	CB-CA-C	-5.48	99.45	110.40
1	B	174	ARG	NE-CZ-NH2	-5.32	117.64	120.30
1	B	316	ASP	CB-CG-OD2	-5.32	113.51	118.30
1	B	392	ARG	NE-CZ-NH1	5.31	122.95	120.30
1	B	447	ARG	NE-CZ-NH2	5.15	122.88	120.30
1	A	316	ASP	CB-CG-OD1	5.13	122.92	118.30
1	B	125	ARG	NE-CZ-NH2	-5.06	117.77	120.30
1	B	217	GLN	CA-CB-CG	5.05	124.52	113.40
1	B	130	ARG	NE-CZ-NH1	5.05	122.83	120.30
1	A	229	ARG	NE-CZ-NH1	5.03	122.82	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	404	CYS	Peptide
1	A	406[A]	CYS	Peptide
1	A	428	ASP	Peptide
1	B	406[B]	CYS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5187	0	5026	36	0
1	B	5178	0	5016	48	0
2	A	24	0	18	2	0
2	B	20	0	15	0	0
3	A	324	0	0	2	0
3	B	280	0	0	2	0
All	All	11013	0	10075	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:571:ARG:HD3	1:B:642:LYS:HG3	1.45	0.94
1:B:74:ALA:HB2	1:B:141:LEU:HD13	1.55	0.86
1:A:343:GLU:OE2	2:A:706:ACT:H1	1.84	0.77
1:A:490:VAL:HG22	1:A:522:LEU:HD23	1.67	0.76
1:A:447:ARG:HD3	1:A:471:SER:O	1.86	0.75
1:A:404:CYS:SG	1:A:407[B]:CYS:HA	2.28	0.74
1:B:357:ALA:O	1:B:362:THR:HB	1.89	0.73
1:B:571:ARG:HB2	1:B:642:LYS:HE2	1.71	0.70
1:B:351:ALA:HB3	1:B:480:ARG:HD3	1.72	0.70
1:B:384:LYS:O	1:B:387:GLU:HG2	1.93	0.68
1:A:485[A]:CYS:SG	1:A:487:GLY:O	2.51	0.67
1:B:498:ASP:OD1	1:B:500:VAL:HG12	1.95	0.66
1:A:37:ASN:HD22	1:A:39:ARG:NH1	1.95	0.65
1:B:42:ASP:N	1:B:42:ASP:OD1	2.30	0.65
1:A:351:ALA:HB3	1:A:480:ARG:HD3	1.80	0.62
1:B:203:ASN:OD1	3:B:1049:HOH:O	2.17	0.59
1:B:425:GLN:NE2	1:B:426:THR:O	2.36	0.59
1:A:426:THR:HG22	1:A:427:SER:H	1.69	0.58
1:B:570:PRO:HA	1:B:642:LYS:HG2	1.85	0.57
1:A:308:ALA:HA	1:A:325:TYR:CD1	2.41	0.56
1:A:426:THR:HG22	1:A:427:SER:N	2.21	0.55
1:B:407[B]:CYS:HB3	1:B:408[B]:PRO:HD3	1.89	0.54
1:B:406[B]:CYS:O	1:B:407[B]:CYS:HB2	2.08	0.54
1:B:218:GLN:HG3	1:B:219:PRO:HA	1.89	0.53
1:A:264:LEU:HD11	1:A:304:VAL:HG21	1.90	0.53
1:B:576:GLU:OE2	1:B:578:HIS:ND1	2.38	0.53
1:A:595:GLU:OE1	1:A:642:LYS:HE3	2.09	0.53
1:B:233:GLU:HA	1:B:233:GLU:OE1	2.08	0.52
1:B:129:LEU:HD12	1:B:221:TYR:CE2	2.44	0.52
1:B:308:ALA:HA	1:B:325:TYR:CD1	2.45	0.51
1:A:566:ASN:HD21	1:A:597:VAL:H	1.58	0.51
1:A:135:LEU:HB2	1:A:183:ASP:HB2	1.93	0.51
1:B:486:ARG:HH21	1:B:529:GLU:CD	2.14	0.50
1:A:47:HIS:HD2	1:A:51:ASN:OD1	1.94	0.49
1:B:500:VAL:HG13	1:B:501:PRO:HD3	1.94	0.49
1:A:304:VAL:HG23	1:A:330:ALA:HB1	1.95	0.49
1:B:117:TYR:O	1:B:122:PRO:HA	2.12	0.49
1:A:465:LEU:N	1:A:465:LEU:HD12	2.29	0.48
1:B:110:ASP:OD2	1:B:125:ARG:NH2	2.45	0.48
1:B:373:PHE:CE2	1:B:408[A]:PRO:HB3	2.49	0.48
1:A:467:VAL:HG12	1:A:469:PRO:HD3	1.95	0.48
1:A:452:MET:O	1:A:465:LEU:HA	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:ASN:ND2	1:A:39:ARG:NH1	2.62	0.47
1:B:362:THR:HG22	1:B:363:ILE:N	2.30	0.47
1:B:599:GLU:O	1:B:602:TRP:HB2	2.14	0.47
1:B:58:GLU:OE2	1:B:105:ARG:NE	2.30	0.47
1:B:121:GLU:OE1	1:B:124:LYS:HE2	2.15	0.47
1:A:544:VAL:HG11	1:A:569:LEU:HD23	1.97	0.47
1:A:402:PHE:HD1	2:A:704:ACT:H3	1.81	0.46
1:A:155:ARG:HG3	1:A:155:ARG:O	2.16	0.46
1:B:627:TRP:CE3	1:B:628:CYS:HB2	2.51	0.45
1:B:334:LEU:O	1:B:338:THR:HG23	2.17	0.45
1:B:408[B]:PRO:HB2	1:B:409:PRO:HD3	1.98	0.45
1:B:346:MET:HG2	1:B:431:PHE:CE1	2.52	0.45
1:B:206:LYS:HA	1:B:209:GLN:HE21	1.82	0.45
1:A:407[A]:CYS:HB3	1:A:408[A]:PRO:HD3	1.98	0.44
1:A:380:GLU:HG3	1:A:636:VAL:HG22	1.99	0.44
1:A:112:TYR:CE2	1:A:114:ASN:HB3	2.52	0.44
1:B:603:ASN:OD1	1:B:603:ASN:N	2.49	0.44
1:B:408[A]:PRO:N	1:B:409:PRO:CD	2.81	0.44
1:A:605:GLU:HG2	3:A:957:HOH:O	2.17	0.44
1:B:304:VAL:HG23	1:B:330:ALA:HB1	2.00	0.44
1:A:640:ASN:ND2	3:A:869:HOH:O	2.51	0.44
1:B:479:LEU:HD13	1:B:522:LEU:HD21	1.99	0.43
1:B:610:ILE:O	1:B:610:ILE:HG13	2.18	0.43
1:A:237:TYR:CD2	1:A:243:TYR:HB2	2.54	0.43
1:B:571:ARG:H	1:B:642:LYS:HE2	1.84	0.43
1:B:555:LEU:O	1:B:635:MET:HA	2.18	0.43
1:B:227:GLU:HG3	1:B:228:ALA:N	2.34	0.42
1:B:460:ASP:OD2	3:B:1058:HOH:O	2.21	0.42
1:B:571:ARG:H	1:B:642:LYS:HG2	1.84	0.42
1:A:36:LEU:HD21	1:A:72:ASP:HB3	2.02	0.42
1:B:409:PRO:HA	1:B:412:ALA:HB3	2.02	0.42
1:A:58:GLU:OE2	1:A:105:ARG:NE	2.44	0.41
1:A:490:VAL:HG22	1:A:522:LEU:CD2	2.45	0.41
1:A:461:GLY:HA3	1:A:524:PHE:O	2.21	0.41
1:B:362:THR:HG22	1:B:363:ILE:H	1.85	0.41
1:A:602:TRP:CH2	1:A:610:ILE:HD11	2.56	0.41
1:B:95:LEU:O	1:B:99:VAL:HG23	2.21	0.41
1:A:214:GLN:O	1:A:217:GLN:HG2	2.21	0.40
1:B:274:ALA:HB2	1:B:282:LEU:HB3	2.03	0.40
1:A:89:ASP:C	1:A:89:ASP:OD1	2.57	0.40
1:B:192:LEU:HD23	1:B:192:LEU:HA	1.91	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:29:ILE:HA	1:B:29:ILE:HD13	1.90	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	640/648 (99%)	622 (97%)	16 (2%)	2 (0%)	46	57
1	B	638/648 (98%)	616 (97%)	16 (2%)	6 (1%)	21	24
All	All	1278/1296 (99%)	1238 (97%)	32 (2%)	8 (1%)	34	36

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	187	GLU
1	B	187	GLU
1	B	407[A]	CYS
1	B	407[B]	CYS
1	B	405	ALA
1	A	405	ALA
1	B	408[A]	PRO
1	B	408[B]	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	540/545 (99%)	517 (96%)	23 (4%)	35	47
1	B	538/545 (99%)	511 (95%)	27 (5%)	30	41
All	All	1078/1090 (99%)	1028 (95%)	50 (5%)	33	44

All (50) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	VAL
1	A	12	LEU
1	A	13	LYS
1	A	44	GLU
1	A	59	SER
1	A	91	GLU
1	A	95	LEU
1	A	113	LEU
1	A	155	ARG
1	A	157	PHE
1	A	176	GLU
1	A	282	LEU
1	A	407[A]	CYS
1	A	407[B]	CYS
1	A	410	ASN
1	A	411	LEU
1	A	430	LEU
1	A	444	ILE
1	A	516	GLN
1	A	571	ARG
1	A	574	LYS
1	A	605	GLU
1	A	639	VAL
1	B	8	THR
1	B	18	LYS
1	B	42	ASP
1	B	59	SER
1	B	60	ASP
1	B	91	GLU
1	B	113	LEU
1	B	141	LEU
1	B	164	TYR
1	B	217	GLN
1	B	218	GLN
1	B	238	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	282	LEU
1	B	304	VAL
1	B	362	THR
1	B	387	GLU
1	B	410	ASN
1	B	411	LEU
1	B	428	ASP
1	B	430	LEU
1	B	441	GLN
1	B	450	LYS
1	B	460	ASP
1	B	486	ARG
1	B	516	GLN
1	B	572	ASP
1	B	614	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	47	HIS
1	A	140	HIS
1	A	377	ASN
1	A	410	ASN
1	A	456	ASN
1	A	515	GLN
1	A	566	ASN
1	A	640	ASN
1	B	140	HIS
1	B	209	GLN
1	B	217	GLN
1	B	218	GLN
1	B	255	GLN
1	B	284	GLN
1	B	377	ASN
1	B	640	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	A	701	-	0,3,3	0.00	-	0,3,3	0.00	-
2	ACT	A	702	-	0,3,3	0.00	-	0,3,3	0.00	-
2	ACT	A	703	-	0,3,3	0.00	-	0,3,3	0.00	-
2	ACT	A	704	-	0,3,3	0.00	-	0,3,3	0.00	-
2	ACT	A	705	-	0,3,3	0.00	-	0,3,3	0.00	-
2	ACT	A	706	-	0,3,3	0.00	-	0,3,3	0.00	-
2	ACT	B	701	-	0,3,3	0.00	-	0,3,3	0.00	-
2	ACT	B	702	-	0,3,3	0.00	-	0,3,3	0.00	-
2	ACT	B	703	-	0,3,3	0.00	-	0,3,3	0.00	-
2	ACT	B	704	-	0,3,3	0.00	-	0,3,3	0.00	-
2	ACT	B	705	-	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACT	A	701	-	-	0/0/0/0	0/0/0/0
2	ACT	A	702	-	-	0/0/0/0	0/0/0/0
2	ACT	A	703	-	-	0/0/0/0	0/0/0/0
2	ACT	A	704	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACT	A	705	-	-	0/0/0/0	0/0/0/0
2	ACT	A	706	-	-	0/0/0/0	0/0/0/0
2	ACT	B	701	-	-	0/0/0/0	0/0/0/0
2	ACT	B	702	-	-	0/0/0/0	0/0/0/0
2	ACT	B	703	-	-	0/0/0/0	0/0/0/0
2	ACT	B	704	-	-	0/0/0/0	0/0/0/0
2	ACT	B	705	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	704	ACT	1	0
2	A	706	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	637/648 (98%)	-0.33	22 (3%)	48 56	18, 28, 59, 106	0
1	B	637/648 (98%)	-0.13	23 (3%)	46 55	20, 32, 64, 118	0
All	All	1274/1296 (98%)	-0.23	45 (3%)	48 56	18, 30, 62, 118	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	6	VAL	9.8
1	A	6	VAL	8.5
1	A	446	GLY	7.0
1	B	7	ALA	6.9
1	A	7	ALA	6.3
1	B	402	PHE	5.9
1	B	232	THR	5.8
1	A	407[A]	CYS	5.4
1	B	43	ALA	5.3
1	B	407[A]	CYS	4.4
1	B	445	ASP	4.4
1	B	427	SER	4.3
1	A	234	PRO	4.2
1	B	233	GLU	4.0
1	B	401	TRP	4.0
1	B	8	THR	3.9
1	A	402	PHE	3.8
1	A	445	ASP	3.8
1	A	233	GLU	3.8
1	A	177	GLY	3.8
1	B	44	GLU	3.6
1	A	42	ASP	3.5
1	B	403	SER	3.5
1	B	234	PRO	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	232	THR	3.3
1	B	603	ASN	3.2
1	B	446	GLY	3.2
1	A	176	GLU	3.1
1	B	238	ASP	3.1
1	B	41	PRO	3.1
1	B	42	ASP	2.8
1	A	403	SER	2.7
1	A	428	ASP	2.6
1	B	572	ASP	2.6
1	A	41	PRO	2.5
1	A	604	ASP	2.5
1	A	603	ASN	2.4
1	A	238	ASP	2.4
1	A	44	GLU	2.4
1	B	64	TYR	2.3
1	A	43	ALA	2.3
1	B	227	GLU	2.3
1	B	230	GLY	2.3
1	A	611	GLU	2.1
1	A	231	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ACT	B	702	4/4	0.66	0.36	5.80	50,52,57,58	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ACT	A	703	4/4	0.85	0.23	4.94	61,67,67,69	0
2	ACT	B	705	4/4	0.89	0.20	3.10	48,49,50,51	0
2	ACT	B	704	4/4	0.68	0.23	2.55	64,67,70,72	0
2	ACT	B	701	4/4	0.82	0.20	2.05	60,61,63,65	0
2	ACT	A	704	4/4	0.87	0.27	0.83	57,57,62,63	0
2	ACT	B	703	4/4	0.96	0.19	0.25	40,45,46,46	0
2	ACT	A	706	4/4	0.95	0.11	-0.67	47,48,49,59	0
2	ACT	A	701	4/4	0.97	0.09	-1.74	29,33,33,34	0
2	ACT	A	702	4/4	0.88	0.16	-	61,62,62,64	0
2	ACT	A	705	4/4	0.61	0.26	-	65,67,67,71	0

6.5 Other polymers [i](#)

There are no such residues in this entry.