



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:23 AM GMT

PDB ID : 2QKN
Title : Crystal structure of Maize cytokinin oxidase/dehydrogenase complexed with phenylurea inhibitor CPPU
Authors : Briozzo, P.
Deposited on : 2007-07-11
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	516	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	1602	-	-	-	X
5	245	A	1537[B]	-	-	X	-
6	GOL	A	718	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 4284 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cytokinin dehydrogenase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	496	Total	C	N	O	S	0	0	0
			3784	2402	665	706	11			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	79	ALA	GLY	VARIANT	UNP Q9T0N8
A	168	THR	ASN	VARIANT	UNP Q9T0N8
A	254	LEU	PHE	VARIANT	UNP Q9T0N8

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 5 is 1-(2-CHLOROPYRIDIN-4-YL)-3-PHENYLUREA (three-letter code: 245) (formula: $C_{12}H_{10}ClN_3O$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	Cl	N	O	0	1
			17	12	1	3	1		
5	A	1	Total	C	Cl	N	O	0	1
			17	12	1	3	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	295	Total 295	O 295	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cytokinin dehydrogenase 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	249.54Å 50.52Å 51.26Å 90.00° 94.01° 90.00°	Depositor
Resolution (Å)	30.00 – 2.15 29.84 – 2.09	Depositor EDS
% Data completeness (in resolution range)	10.0 (30.00-2.15) 89.7 (29.84-2.09)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.54 (at 2.08Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.191 , 0.231 0.210 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.746	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 56.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 34424 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4284	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 245, GOL, NAG, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	1/3873 (0.0%)	0.59	0/5277

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	462	VAL	C-N	6.96	1.50	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3784	0	3702	51	0
2	A	28	0	26	0	0
3	A	84	0	75	5	0
4	A	53	0	29	1	0
5	A	34	0	20	12	0
6	A	6	0	8	0	0
7	A	295	0	0	4	0
All	All	4284	0	3860	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 8.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1602:NAG:H62	3:A:1603:NAG:H82	1.36	1.05
5:A:1537[B]:245:H8	5:A:1537[B]:245:O1	1.79	0.80
5:A:1536[A]:245:H11	5:A:1536[A]:245:O1	1.81	0.79
5:A:1537[B]:245:H11	5:A:1537[B]:245:O1	1.81	0.79
1:A:141:TYR:CE1	1:A:239:ARG:HD2	2.17	0.79
1:A:143:ASP:OD1	1:A:239:ARG:HD3	1.83	0.79
5:A:1536[A]:245:H8	5:A:1536[A]:245:O1	1.85	0.74
1:A:463:ALA:HB3	1:A:466:ASP:HB2	1.73	0.69
1:A:141:TYR:CD1	1:A:239:ARG:HD2	2.29	0.66
1:A:524:ARG:HG3	1:A:524:ARG:HH11	1.61	0.65
1:A:47:LEU:HD22	1:A:73:LEU:HD13	1.78	0.64
1:A:492:LEU:CD1	5:A:1537[B]:245:CL7	2.83	0.64
1:A:136:SER:HB3	1:A:141:TYR:CE1	2.33	0.63
1:A:253:ARG:HD2	1:A:306:PHE:CD2	2.33	0.63
1:A:90:SER:O	1:A:92:PRO:HD3	1.98	0.63
5:A:1537[B]:245:H4	7:A:1611:HOH:O	2.00	0.62
1:A:463:ALA:HB3	1:A:466:ASP:CB	2.32	0.59
1:A:392:ARG:HH11	1:A:392:ARG:HG3	1.68	0.58
1:A:48:ARG:HD3	1:A:74:TYR:CE2	2.39	0.57
5:A:1536[A]:245:H15	7:A:1611:HOH:O	2.07	0.55
5:A:1537[B]:245:O1	5:A:1537[B]:245:C11	2.50	0.53
3:A:1602:NAG:C6	3:A:1603:NAG:H82	2.25	0.53
1:A:479:ARG:HD2	7:A:1881:HOH:O	2.11	0.51
1:A:428:LEU:HD22	1:A:457:LEU:HD22	1.94	0.49
1:A:41:LEU:CB	1:A:47:LEU:HD23	2.42	0.49
1:A:495:HIS:HD2	7:A:1811:HOH:O	1.95	0.49
1:A:41:LEU:HB2	1:A:47:LEU:HD23	1.94	0.48
1:A:386:ASN:OD1	1:A:392:ARG:NH1	2.47	0.48
1:A:398:LEU:HG	1:A:400:MET:HE2	1.96	0.48
3:A:1606:NAG:O3	3:A:1606:NAG:H83	2.13	0.47
1:A:392:ARG:NH1	1:A:392:ARG:HG3	2.30	0.47
1:A:47:LEU:CD2	1:A:73:LEU:HD13	2.44	0.46
1:A:66:SER:HA	1:A:112:PHE:CE2	2.50	0.46
1:A:73:LEU:HG	1:A:75:PRO:HD3	1.98	0.46
1:A:164:PRO:HD2	1:A:167:TRP:CH2	2.51	0.46
3:A:1602:NAG:H62	3:A:1603:NAG:C8	2.26	0.45
5:A:1536[A]:245:O1	5:A:1536[A]:245:C8	2.51	0.45
1:A:88:ALA:HA	1:A:94:TRP:CD2	2.51	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:401:PHE:HA	1:A:453:TYR:O	2.16	0.45
3:A:1606:NAG:H61	3:A:1607:NAG:HN2	1.82	0.45
1:A:336:TYR:CD2	1:A:346:VAL:HG21	2.52	0.44
1:A:136:SER:HB3	1:A:141:TYR:CD1	2.53	0.44
1:A:254:LEU:O	1:A:329:SER:HA	2.17	0.44
1:A:254:LEU:HD12	1:A:364:ALA:HA	1.99	0.44
1:A:48:ARG:HB3	1:A:50:ASP:OD1	2.17	0.44
1:A:382:GLU:HG3	1:A:391:TRP:CZ2	2.53	0.43
1:A:462:VAL:HG23	1:A:463:ALA:N	2.34	0.43
1:A:252:VAL:HG22	1:A:253:ARG:N	2.34	0.43
1:A:492:LEU:HD11	5:A:1536[A]:245:H13	2.01	0.42
1:A:397:TRP:CZ3	5:A:1537[B]:245:CL7	3.10	0.42
1:A:427:PRO:HG2	1:A:458:LEU:HD12	2.00	0.42
5:A:1537[B]:245:O1	5:A:1537[B]:245:C8	2.47	0.42
1:A:463:ALA:HB1	1:A:464:PRO:HD2	2.02	0.42
1:A:524:ARG:CG	1:A:524:ARG:HH11	2.30	0.41
1:A:398:LEU:HG	1:A:400:MET:CE	2.50	0.41
1:A:142:VAL:HG13	1:A:142:VAL:O	2.20	0.41
1:A:214:SER:HA	1:A:222:PHE:CD1	2.55	0.41
1:A:103:ARG:HA	1:A:103:ARG:HD3	1.90	0.41
1:A:47:LEU:HD22	1:A:73:LEU:CD1	2.49	0.41
1:A:61:PHE:CD2	4:A:1535:FAD:HM82	2.56	0.41
1:A:102:GLY:O	1:A:103:ARG:NH1	2.53	0.41
1:A:318:LEU:HD21	1:A:360:VAL:HG21	2.03	0.41
1:A:338:ASN:O	1:A:339:ALA:C	2.59	0.41
1:A:202:ASP:HB2	1:A:237:ARG:HB3	2.03	0.40
1:A:411:ASP:OD2	1:A:416:LYS:NZ	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	492/516 (95%)	473 (96%)	16 (3%)	3 (1%)	30 21

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	464	PRO
1	A	182	ALA
1	A	282	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	386/395 (98%)	382 (99%)	4 (1%)	82 87

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	LEU
1	A	95	PRO
1	A	254	LEU
1	A	494	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	335	ASN
1	A	399	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	1602	1,3	14,14,15	0.54	0	15,19,21	0.77	1 (6%)
3	NAG	A	1603	3	14,14,15	0.48	0	15,19,21	0.85	1 (6%)
3	NAG	A	1604	1,3	14,14,15	0.48	0	15,19,21	0.70	1 (6%)
3	NAG	A	1605	3	14,14,15	0.50	0	15,19,21	0.77	1 (6%)
3	NAG	A	1606	1,3	14,14,15	0.56	0	15,19,21	0.73	1 (6%)
3	NAG	A	1607	3	14,14,15	0.45	0	15,19,21	0.67	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1602	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1603	3	-	0/6/23/26	0/1/1/1
3	NAG	A	1604	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1605	3	-	0/6/23/26	0/1/1/1
3	NAG	A	1606	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1607	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1603	NAG	C2-N2-C7	-2.61	119.69	123.04

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1602	NAG	C2-N2-C7	-2.37	119.99	123.04
3	A	1605	NAG	C2-N2-C7	-2.28	120.11	123.04
3	A	1606	NAG	C2-N2-C7	-2.20	120.21	123.04
3	A	1604	NAG	C2-N2-C7	-2.12	120.31	123.04
3	A	1607	NAG	C2-N2-C7	-2.07	120.38	123.04

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1602	NAG	3	0
3	A	1603	NAG	3	0
3	A	1606	NAG	2	0
3	A	1607	NAG	1	0

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	FAD	A	1535	1	48,58,58	1.99	10 (20%)	54,89,89	1.86	2 (3%)
5	245	A	1536[A]	-	18,18,18	1.67	4 (22%)	22,23,23	1.82	4 (18%)
5	245	A	1537[B]	-	18,18,18	1.66	4 (22%)	22,23,23	1.83	5 (22%)
2	NAG	A	1601	1	14,14,15	0.46	0	15,19,21	0.64	0
2	NAG	A	1608	1	14,14,15	0.47	0	15,19,21	0.71	1 (6%)
6	GOL	A	718	-	5,5,5	0.31	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FAD	A	1535	1	-	0/30/50/50	0/6/6/6
5	245	A	1536[A]	-	-	0/8/8/8	0/2/2/2
5	245	A	1537[B]	-	-	0/8/8/8	0/2/2/2
2	NAG	A	1601	1	-	0/6/23/26	0/1/1/1
2	NAG	A	1608	1	-	0/6/23/26	0/1/1/1
6	GOL	A	718	-	-	0/4/4/4	0/0/0/0

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1536[A]	245	C3-N2	-4.36	1.33	1.41
5	A	1537[B]	245	C10-N9	-4.26	1.33	1.41
5	A	1537[B]	245	C3-N2	-4.26	1.33	1.41
5	A	1536[A]	245	C10-N9	-4.23	1.33	1.41
5	A	1536[A]	245	C1-N9	-2.26	1.33	1.37
5	A	1537[B]	245	C1-N9	-2.23	1.33	1.37
5	A	1537[B]	245	C1-N2	-2.23	1.33	1.37
5	A	1536[A]	245	C1-N2	-2.13	1.33	1.37
4	A	1535	FAD	C5A-C4A	-2.09	1.35	1.40
4	A	1535	FAD	P-O5'	2.03	1.68	1.59
4	A	1535	FAD	O4B-C1B	2.39	1.44	1.41
4	A	1535	FAD	C9-C8	3.20	1.46	1.37
4	A	1535	FAD	C4A-N3A	3.70	1.41	1.35
4	A	1535	FAD	C5X-N5	3.82	1.41	1.35
4	A	1535	FAD	C8M-C8	3.86	1.58	1.51
4	A	1535	FAD	C4-N3	4.40	1.41	1.33
4	A	1535	FAD	C9A-N10	5.47	1.46	1.38
4	A	1535	FAD	C4X-N5	5.98	1.42	1.33

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1535	FAD	C4X-C4-N3	-5.91	115.51	123.59
5	A	1537[B]	245	C8-C7-N6	-4.85	120.41	125.46
5	A	1536[A]	245	C8-C7-N6	-4.83	120.43	125.46
5	A	1536[A]	245	C4-C5-N6	-2.87	120.62	123.90
5	A	1537[B]	245	C4-C5-N6	-2.79	120.72	123.90
2	A	1608	NAG	C2-N2-C7	-2.26	120.14	123.04
5	A	1537[B]	245	C3-N2-C1	-2.16	122.48	126.65
5	A	1536[A]	245	N2-C1-N9	2.08	115.79	112.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
5	A	1537[B]	245	N2-C1-N9	2.18	115.96	112.53
5	A	1537[B]	245	C5-N6-C7	4.78	121.65	115.36
5	A	1536[A]	245	C5-N6-C7	4.80	121.67	115.36
4	A	1535	FAD	C4-N3-C2	10.76	124.55	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1535	FAD	1	0
5	A	1536[A]	245	5	0
5	A	1537[B]	245	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	496/516 (96%)	0.40	51 (10%) 9 14	18, 30, 50, 60	14 (2%)

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	128	ALA	9.2
1	A	464	PRO	9.0
1	A	129	ALA	9.0
1	A	127	ALA	7.7
1	A	390	LEU	4.3
1	A	389	GLY	4.2
1	A	92	PRO	4.2
1	A	388	LEU	4.1
1	A	462	VAL	4.1
1	A	234	VAL	3.6
1	A	229	LEU	3.5
1	A	344	ALA	3.4
1	A	273	ARG	3.4
1	A	130	PRO	3.3
1	A	100	PHE	3.2
1	A	180	SER	3.2
1	A	352	SER	3.1
1	A	179	LEU	2.9
1	A	217	LEU	2.9
1	A	131	PRO	2.9
1	A	182	ALA	2.8
1	A	348	GLN	2.8
1	A	126	ASP	2.7
1	A	383	VAL	2.6
1	A	384	ALA	2.6
1	A	225	VAL	2.5
1	A	463	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	385	LEU	2.5
1	A	204	ILE	2.5
1	A	430	VAL	2.5
1	A	228	GLY	2.4
1	A	216	GLN	2.4
1	A	230	GLY	2.4
1	A	36	ALA	2.4
1	A	465	ASN	2.3
1	A	386	ASN	2.3
1	A	391	TRP	2.3
1	A	91	THR	2.3
1	A	351	ALA	2.3
1	A	235	ILE	2.2
1	A	343	ALA	2.2
1	A	51	SER	2.1
1	A	99	ALA	2.1
1	A	177	GLY	2.1
1	A	210	MET	2.1
1	A	119	VAL	2.1
1	A	339	ALA	2.1
1	A	380	GLY	2.0
1	A	338	ASN	2.0
1	A	137	ALA	2.0
1	A	347	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	A	1602	14/15	0.60	0.34	2.54	46,48,51,52	14
3	NAG	A	1605	14/15	0.81	0.17	-	45,47,49,49	14

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAG	A	1606	14/15	0.83	0.23	-	39,42,44,47	14
3	NAG	A	1607	14/15	0.64	0.41	-	50,53,54,55	14
3	NAG	A	1603	14/15	0.42	0.46	-	53,54,55,56	14
3	NAG	A	1604	14/15	0.90	0.14	-	32,37,42,42	14

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	GOL	A	718	6/6	0.85	0.19	5.56	33,37,41,46	0
5	245	A	1537[B]	17/17	0.85	0.19	0.85	15,20,28,29	17
5	245	A	1536[A]	17/17	0.84	0.18	0.43	14,21,29,30	17
4	FAD	A	1535	53/53	0.97	0.17	-0.31	18,23,26,27	0
2	NAG	A	1601	14/15	0.93	0.11	-0.66	29,33,37,37	14
2	NAG	A	1608	14/15	0.48	0.34	-	54,55,57,58	14

6.5 Other polymers

There are no such residues in this entry.