



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:58 AM GMT

PDB ID : 3QN1
Title : Crystal structure of the PYR1 Absciscic Acid receptor in complex with the
HAB1 type 2C phosphatase catalytic domain
Authors : Betz, K.; Dupeux, F.; Santiago, J.; Marquez, J.A.
Deposited on : 2011-02-07
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	193	<div> <div>4%</div> <div>86%</div> <div>7%</div> <div>7%</div> </div>
2	B	337	<div> <div>6%</div> <div>78%</div> <div>9%</div> <div>12%</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4161 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Absciscic acid receptor PYR1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	179	Total	C	N	O	S	0	0	0
			1414	885	247	274	8			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	EXPRESSION TAG	UNP O49686
A	0	ALA	-	EXPRESSION TAG	UNP O49686
A	1	MET	-	EXPRESSION TAG	UNP O49686
A	2	ALA	-	EXPRESSION TAG	UNP O49686


- Molecule 2 is a protein called Protein phosphatase 2C 16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	295	Total	C	N	O	S	0	0	0
			2260	1423	398	422	17			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	175	GLY	-	EXPRESSION TAG	UNP Q9CAJ0
B	176	ALA	-	EXPRESSION TAG	UNP Q9CAJ0
B	177	MET	-	EXPRESSION TAG	UNP Q9CAJ0

- Molecule 3 is (2Z,4E)-5-[(1S)-1-HYDROXY-2,6,6-TRIMETHYL-4-OXOCYCLOHEX-2-EN-1-YL]-3-METHYLPENTA-2,4-DIENOIC ACID (three-letter code: A8S) (formula: C₁₅H₂₀O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			19	15	4		

- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	3	Total	Mn	0	0
			3	3		



- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	218	Total	O	0	0
			218	218		
5	B	247	Total	O	0	0
			247	247		




3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Absciscic acid receptor PYR1



- Molecule 2: Protein phosphatase 2C 16



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	45.85Å 65.86Å 170.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.24 – 1.80 28.24 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (28.24-1.80) 100.0 (28.24-1.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.81 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.174 , 0.218 0.175 , 0.218	Depositor DCC
R_{free} test set	2312 reflections (4.96%)	DCC
Wilson B-factor (Å ²)	21.4	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 48943 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4161	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A8S, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	0/1440	0.84	0/1950
2	B	0.91	0/2301	0.85	3/3109 (0.1%)
All	All	0.94	0/3741	0.85	3/5059 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	199	ARG	NE-CZ-NH2	-5.88	117.36	120.30
2	B	257	ARG	NE-CZ-NH2	-5.69	117.45	120.30
2	B	353	ARG	NE-CZ-NH1	-5.50	117.55	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1414	0	1384	30	0
2	B	2260	0	2219	27	0
3	A	19	0	19	2	0
4	B	3	0	0	0	0
5	A	218	0	0	4	0
5	B	247	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4161	0	3622	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:158:MET:CE	2:B:383:ILE:HD11	1.64	1.26
1:A:58:TYR:HA	1:A:170:LYS:HE2	1.34	1.09
1:A:58:TYR:O	1:A:170:LYS:HD3	1.53	1.07
1:A:158:MET:HE2	2:B:383:ILE:CD1	1.89	1.01
1:A:67:VAL:HG11	1:A:73:MET:CE	1.96	0.95
1:A:60:HIS:CD2	1:A:170:LYS:HD2	2.02	0.95
1:A:158:MET:HE2	2:B:383:ILE:HD11	0.93	0.92
2:B:345:GLY:H	2:B:364:HIS:HD2	1.19	0.89
1:A:67:VAL:HG13	5:A:252:HOH:O	1.77	0.84
2:B:245:HIS:HE1	2:B:402:ASP:OD2	1.60	0.84
1:A:67:VAL:HG11	1:A:73:MET:HE3	1.56	0.83
1:A:58:TYR:HA	1:A:170:LYS:CE	2.13	0.76
2:B:185:ASP:N	2:B:185:ASP:OD2	2.18	0.76
1:A:158:MET:CE	2:B:383:ILE:CD1	2.57	0.75
5:A:231:HOH:O	2:B:323:GLU:HG2	1.87	0.73
2:B:345:GLY:H	2:B:364:HIS:CD2	2.06	0.71
1:A:60:HIS:CE1	1:A:170:LYS:NZ	2.59	0.70
1:A:58:TYR:C	1:A:170:LYS:HD3	2.13	0.69
1:A:67:VAL:HG12	1:A:68:GLU:N	2.10	0.66
2:B:203:GLU:H	2:B:493:ASN:ND2	1.94	0.66
2:B:245:HIS:HD2	5:B:9:HOH:O	1.77	0.66
1:A:158:MET:HE1	2:B:381:LYS:HB2	1.77	0.65
1:A:67:VAL:CG1	1:A:73:MET:CE	2.73	0.65
1:A:67:VAL:HG11	1:A:73:MET:SD	2.37	0.64
2:B:364:HIS:HE1	2:B:414:GLU:OE2	1.84	0.61
1:A:158:MET:SD	2:B:381:LYS:HG3	2.44	0.58
1:A:60:HIS:CE1	1:A:170:LYS:HZ2	2.19	0.57
2:B:185:ASP:N	2:B:420:ARG:HH12	2.03	0.57
1:A:67:VAL:CG1	1:A:73:MET:HE3	2.31	0.56
1:A:67:VAL:CG1	1:A:68:GLU:N	2.73	0.51
2:B:265:GLU:OE2	2:B:268:ARG:NH1	2.44	0.51
2:B:480:TYR:CE2	2:B:484:LEU:HD11	2.46	0.50
2:B:219:MET:HG2	2:B:285:GLN:NE2	2.26	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:203:GLU:H	2:B:493:ASN:HD21	1.61	0.48
2:B:245:HIS:CE1	2:B:402:ASP:OD2	2.52	0.47
3:A:192:A8S:O11	3:A:192:A8S:H4	2.15	0.47
1:A:60:HIS:CG	1:A:170:LYS:HZ2	2.32	0.47
2:B:364:HIS:CE1	2:B:414:GLU:OE2	2.66	0.46
2:B:352:PHE:CG	2:B:444:CYS:HB3	2.50	0.46
1:A:147:MET:SD	1:A:153:GLU:HG2	2.58	0.44
3:A:192:A8S:H6	3:A:192:A8S:H5	1.87	0.44
1:A:157:ARG:HD3	5:A:394:HOH:O	2.16	0.44
1:A:60:HIS:ND1	1:A:170:LYS:NZ	2.57	0.43
2:B:193:VAL:HG11	2:B:482:SER:HB3	2.01	0.43
1:A:158:MET:HB3	1:A:158:MET:HE3	1.80	0.42
1:A:60:HIS:HD2	5:A:299:HOH:O	2.02	0.42
1:A:60:HIS:CG	1:A:170:LYS:HD2	2.50	0.42
2:B:301:ILE:HA	2:B:320:VAL:HG23	2.02	0.42
2:B:323:GLU:HB3	2:B:404:TYR:HB3	2.01	0.41
1:A:158:MET:HE3	2:B:391:PHE:HD1	1.85	0.41
1:A:60:HIS:CE1	1:A:170:LYS:HZ1	2.39	0.41
2:B:345:GLY:N	2:B:364:HIS:HD2	2.01	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	175/193 (91%)	170 (97%)	4 (2%)	1 (1%)	30	14
2	B	287/337 (85%)	284 (99%)	3 (1%)	0	100	100
All	All	462/530 (87%)	454 (98%)	7 (2%)	1 (0%)	52	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	132	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/171 (92%)	156 (99%)	2 (1%)	76	68
2	B	239/280 (85%)	234 (98%)	5 (2%)	61	47
All	All	397/451 (88%)	390 (98%)	7 (2%)	66	54

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	134	ARG
1	A	181	ASN
2	B	185	ASP
2	B	269	ILE
2	B	270	LYS
2	B	421	SER
2	B	466	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	HIS
1	A	90	ASN
2	B	245	HIS
2	B	364	HIS
2	B	487	GLN
2	B	493	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	A8S	A	192	-	13,19,19	5.56	6 (46%)	12,29,29	1.48	4 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A8S	A	192	-	-	0/8/34/34	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	192	A8S	C4-C3	-4.94	1.35	1.45
3	A	192	A8S	C9-C10	-3.14	1.38	1.45
3	A	192	A8S	C13-C8	2.03	1.55	1.50
3	A	192	A8S	O10-C10	7.10	1.34	1.23
3	A	192	A8S	C4-C5	8.86	1.49	1.32
3	A	192	A8S	C9-C8	15.19	1.50	1.33

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	A	192	A8S	C8-C9-C10	-2.53	121.18	123.87
3	A	192	A8S	C6-C3-C2	-2.18	120.35	125.17
3	A	192	A8S	C15-C12-C11	-2.05	103.71	108.16
3	A	192	A8S	C11-C10-C9	2.33	120.38	117.60

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	192	A8S	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	179/193 (92%)	-0.20	7 (3%) 43 37	14, 21, 39, 52	0
2	B	295/337 (87%)	-0.03	20 (6%) 20 16	14, 21, 50, 59	0
All	All	474/530 (89%)	-0.10	27 (5%) 27 22	14, 21, 46, 59	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	71	PHE	6.8
2	B	232	THR	5.8
2	B	217	ILE	4.7
2	B	284	VAL	4.5
2	B	216	PRO	4.4
2	B	310	VAL	4.2
1	A	132	GLU	3.8
1	A	133	ASN	3.6
2	B	220	LEU	3.6
2	B	186	CYS	3.6
2	B	460	ALA	3.5
1	A	181	ASN	2.9
2	B	215	LEU	2.7
2	B	185	ASP	2.6
1	A	73	MET	2.5
2	B	233	HIS	2.5
2	B	332	ALA	2.3
1	A	67	VAL	2.3
2	B	459	GLY	2.3
2	B	331	VAL	2.3
2	B	221	MET	2.3
2	B	466	ARG	2.2
2	B	219	MET	2.2
2	B	269	ILE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	468	LYS	2.2
2	B	467	GLY	2.1
1	A	68	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	A8S	A	192	19/19	0.97	0.11	1.51	13,16,19,21	0
4	MN	B	1	1/1	1.00	0.06	-0.92	16,16,16,16	0
4	MN	B	2	1/1	1.00	0.06	-1.55	27,27,27,27	0
4	MN	B	3	1/1	0.94	0.47	-	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.