



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:00 PM GMT

PDB ID : 4QNE
Title : Inosine 5'-monophosphate dehydrogenase from *Vibrio cholerae*, deletion mutant, in complex with NAD and IMP
Authors : Osipiuk, J.; Maltseva, N.; Makowska-Grzyska, M.; Gu, M.; Anderson, W.F.; Joachimiak, A.; Center for Structural Genomics of Infectious Diseases (CS-GID)
Deposited on : 2014-06-17
Resolution : 2.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4425 (2.34-2.30)
Clashscore	102246	5057 (2.34-2.30)
Ramachandran outliers	100387	5008 (2.34-2.30)
Sidechain outliers	100360	5007 (2.34-2.30)
RSRZ outliers	91569	4432 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	
1	B	366	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAJ	A	502	-	-	-	X
3	NAJ	B	502	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5409 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Inosine 5'-monophosphate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	342	Total	C	N	O	S	0	7	0
			2530	1582	454	480	14			
1	B	342	Total	C	N	O	S	0	9	0
			2545	1590	457	484	14			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	EXPRESSION TAG	UNP Q9KTW3
A	-1	ASN	-	EXPRESSION TAG	UNP Q9KTW3
A	0	ALA	-	EXPRESSION TAG	UNP Q9KTW3
A	91	SER	-	LINKER	UNP Q9KTW3
A	92	GLY	-	LINKER	UNP Q9KTW3
A	93	GLY	-	LINKER	UNP Q9KTW3
B	-2	SER	-	EXPRESSION TAG	UNP Q9KTW3
B	-1	ASN	-	EXPRESSION TAG	UNP Q9KTW3
B	0	ALA	-	EXPRESSION TAG	UNP Q9KTW3
B	91	SER	-	LINKER	UNP Q9KTW3
B	92	GLY	-	LINKER	UNP Q9KTW3
B	93	GLY	-	LINKER	UNP Q9KTW3

- Molecule 2 is INOSINIC ACID (three-letter code: IMP) (formula: C₁₀H₁₃N₄O₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	1
			33	15	8	9	1		
2	B	1	Total	C	N	O	P	0	1
			33	15	8	9	1		

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (ACIDIC FORM) (three-letter code: NAJ) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	K	0	0
			1	1		
4	A	1	Total	K	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	103	Total	O	0	0
			103	103		
5	B	75	Total	O	0	0
			75	75		

4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	121.33Å 121.33Å 94.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.50 – 2.32 47.05 – 2.32	Depositor EDS
% Data completeness (in resolution range)	99.0 (47.50-2.32) 99.1 (47.05-2.32)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 2.32Å)	Xtriage
Refinement program	REFMAC 5.8.0071	Depositor
R, R_{free}	0.196 , 0.270 0.202 , 0.266	Depositor DCC
R_{free} test set	1568 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	40.4	Xtriage
Anisotropy	0.248	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	2 of 30891 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5409	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 48.58 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 8.4342e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAJ, IMP, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	0/2583	0.93	4/3490 (0.1%)
1	B	0.76	0/2604	0.91	2/3517 (0.1%)
All	All	0.80	0/5187	0.92	6/7007 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	476	ASP	CB-CG-OD2	6.11	123.80	118.30
1	A	441	ARG	NE-CZ-NH1	6.11	123.35	120.30
1	B	476	ASP	CB-CG-OD1	-6.04	112.87	118.30
1	B	441	ARG	NE-CZ-NH2	-5.74	117.43	120.30
1	A	456	ARG	NE-CZ-NH2	5.56	123.08	120.30
1	A	31	ASP	CB-CG-OD1	5.24	123.01	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	7	ALA	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2530	0	2600	24	0
1	B	2545	0	2614	36	0
2	A	33	0	7	3	0
2	B	33	0	7	6	0
3	A	44	0	27	3	0
3	B	44	0	27	8	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	103	0	0	0	0
5	B	75	0	0	3	0
All	All	5409	0	5282	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:306[B]:ILE:HD12	1:B:389:GLY:N	1.88	0.88
1:B:302:GLY:HA2	1:B:307[A]:CYS:SG	2.17	0.84
1:A:306[B]:ILE:HD13	1:A:387:TYR:CE2	2.13	0.84
1:B:307[A]:CYS:SG	2:B:501[A]:IMP:H2	2.18	0.83
1:A:47:VAL:HG12	1:A:69[B]:ILE:HG22	1.69	0.75
1:B:307[A]:CYS:SG	2:B:501[A]:IMP:C2	2.76	0.74
1:A:6:ILE:HG23	1:A:6:ILE:O	1.93	0.68
1:B:474:VAL:HG11	1:B:479:ILE:HD11	1.76	0.68
1:A:7:ALA:O	1:A:8:LYS:HB3	1.92	0.67
3:B:502:NAJ:H2A	5:B:629:HOH:O	1.95	0.66
1:B:306[B]:ILE:HD12	1:B:388:ARG:C	2.16	0.66
1:A:307[A]:CYS:SG	2:A:501[A]:IMP:H2	2.38	0.64
1:A:474:VAL:HG11	1:A:479:ILE:HD11	1.81	0.60
1:B:388:ARG:HB3	1:B:420[A]:GLU:HG3	1.83	0.60
3:B:502:NAJ:C2A	5:B:629:HOH:O	2.49	0.60
1:A:302:GLY:HA2	1:A:307[A]:CYS:SG	2.41	0.60
1:B:6:ILE:HG23	1:B:6:ILE:O	2.02	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:42:LEU:CD2	1:B:69:ILE:HG22	2.34	0.58
1:B:42:LEU:HD21	1:B:69:ILE:CG2	2.35	0.56
1:B:287:ARG:NH2	1:B:332[A]:GLU:OE1	2.40	0.54
2:B:501[A]:IMP:C2	3:B:502:NAJ:C4N	2.86	0.54
1:B:252:SER:OG	3:B:502:NAJ:H2A	2.07	0.53
1:B:426:LYS:HE2	5:B:624:HOH:O	2.07	0.53
1:B:306[B]:ILE:HD13	1:B:387:TYR:CE2	2.44	0.53
1:A:6:ILE:CG2	1:A:6:ILE:O	2.57	0.53
1:A:306[B]:ILE:HB	1:A:419:ILE:HD12	1.90	0.53
1:B:8:LYS:HG2	1:B:9:GLU:O	2.10	0.52
1:B:42:LEU:HD21	1:B:69:ILE:HG22	1.93	0.51
1:A:307[A]:CYS:SG	2:A:501[A]:IMP:C2	2.99	0.49
3:A:502:NAJ:H6N	3:A:502:NAJ:H51N	1.94	0.49
1:A:252:SER:HB3	3:A:502:NAJ:O1N	2.13	0.49
1:B:454:ASP:OD1	1:B:458:LYS:NZ	2.46	0.49
1:A:382:ARG:HG2	1:B:382:ARG:HG2	1.94	0.48
1:B:302:GLY:N	1:B:303:PRO:CD	2.76	0.48
1:B:42:LEU:HD22	1:B:69:ILE:HG22	1.97	0.47
1:B:252:SER:HG	3:B:502:NAJ:PN	2.38	0.46
1:A:257:GLU:OE2	1:A:264:ARG:NH1	2.48	0.46
1:B:249:ILE:HD12	1:B:275:ILE:HG23	1.98	0.46
1:B:488:LEU:HD13	1:B:488:LEU:C	2.36	0.46
1:A:302:GLY:N	1:A:303:PRO:CD	2.80	0.45
1:A:8:LYS:HG3	1:A:9:GLU:O	2.17	0.44
1:A:290:ILE:HG21	1:A:333:TYR:CG	2.53	0.44
2:B:501[A]:IMP:H2	3:B:502:NAJ:C4N	2.47	0.44
1:A:302:GLY:HA3	1:A:310:ARG:HG3	1.99	0.44
1:B:52:ASP:HA	1:B:73:HIS:CD2	2.52	0.44
1:B:232:ASN:O	1:B:236:VAL:HG23	2.18	0.44
2:A:501[A]:IMP:N3	3:A:502:NAJ:C2N	2.82	0.43
1:B:419:ILE:C	1:B:419:ILE:HD12	2.39	0.43
1:B:85:VAL:O	1:B:89:LYS:HG2	2.19	0.43
1:B:252:SER:N	3:B:502:NAJ:O1N	2.53	0.42
1:A:276:ILE:HG12	1:A:296:ALA:HB3	2.00	0.42
1:B:466:GLY:O	1:B:470:LYS:HD3	2.19	0.42
1:B:349:ILE:HG23	1:B:360:VAL:HG11	2.02	0.42
1:A:47:VAL:CG1	1:A:69[B]:ILE:HG22	2.45	0.42
1:B:36:LEU:HG	1:B:37:THR:HG23	2.02	0.41
1:B:426:LYS:HB3	1:B:432:ILE:HD11	2.02	0.41
1:B:11:LEU:HD22	1:B:15:ASP:HB3	2.03	0.41
1:A:475:HIS:ND1	1:A:476:ASP:OD2	2.53	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:307[A]:CYS:HG	2:B:501[A]:IMP:H2	1.85	0.41
1:B:26:LEU:HB2	1:B:29:THR:HG23	2.02	0.40
1:A:424:ALA:O	1:A:426:LYS:HG3	2.20	0.40
2:B:501[A]:IMP:C2	3:B:502:NAJ:C3N	2.99	0.40
1:A:7:ALA:O	1:A:8:LYS:CB	2.63	0.40
1:B:301:ILE:C	1:B:303:PRO:HD3	2.41	0.40
1:A:281:ALA:HB1	1:A:317:VAL:HB	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/366 (94%)	335 (97%)	9 (3%)	1 (0%)	46	56
1	B	347/366 (95%)	335 (96%)	12 (4%)	0	100	100
All	All	692/732 (94%)	670 (97%)	21 (3%)	1 (0%)	56	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	8	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/274 (96%)	254 (97%)	8 (3%)	47	63
1	B	264/274 (96%)	247 (94%)	17 (6%)	22	28
All	All	526/548 (96%)	501 (95%)	25 (5%)	37	42

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	LEU
1	A	262	ARG
1	A	306[A]	ILE
1	A	306[B]	ILE
1	A	393	LEU
1	A	414	LEU
1	A	482	GLU
1	A	487	ARG
1	B	8	LYS
1	B	69	ILE
1	B	234[A]	GLU
1	B	234[B]	GLU
1	B	262	ARG
1	B	291	GLU
1	B	295	SER
1	B	306[A]	ILE
1	B	306[B]	ILE
1	B	393	LEU
1	B	414	LEU
1	B	431[A]	GLU
1	B	431[B]	GLU
1	B	458	LYS
1	B	460	GLN
1	B	462	VAL
1	B	470	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	80	GLN
1	B	84	GLN
1	B	436	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	IMP	A	501[A]	-	20,25,25	1.15	1 (5%)	22,38,38	2.55	5 (22%)
2	IMP	A	501[B]	-	20,25,25	1.13	1 (5%)	22,38,38	2.79	5 (22%)
3	NAJ	A	502	-	38,48,48	0.93	3 (7%)	47,73,73	2.26	9 (19%)
2	IMP	B	501[A]	-	20,25,25	1.78	2 (10%)	22,38,38	2.68	6 (27%)
2	IMP	B	501[B]	-	20,25,25	1.77	2 (10%)	22,38,38	2.89	7 (31%)
3	NAJ	B	502	-	38,48,48	0.90	2 (5%)	47,73,73	1.74	8 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IMP	A	501[A]	-	-	0/6/26/26	0/3/3/3
2	IMP	A	501[B]	-	-	0/6/26/26	0/3/3/3
3	NAJ	A	502	-	-	0/22/62/62	0/5/5/5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IMP	B	501[A]	-	-	0/6/26/26	0/3/3/3
2	IMP	B	501[B]	-	-	0/6/26/26	0/3/3/3
3	NAJ	B	502	-	-	0/22/62/62	0/5/5/5

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501[A]	IMP	P-O2P	2.12	1.62	1.54
2	B	501[B]	IMP	P-O2P	2.12	1.62	1.54
3	A	502	NAJ	C2A-N3A	2.21	1.36	1.32
3	A	502	NAJ	O4D-C1D	2.46	1.44	1.41
3	B	502	NAJ	O4B-C1B	2.49	1.44	1.41
3	B	502	NAJ	O4D-C1D	2.98	1.45	1.41
3	A	502	NAJ	O4B-C1B	3.13	1.45	1.41
2	A	501[B]	IMP	O4'-C1'	3.73	1.45	1.41
2	A	501[A]	IMP	O4'-C1'	3.73	1.45	1.41
2	B	501[A]	IMP	O4'-C1'	6.35	1.49	1.41
2	B	501[B]	IMP	O4'-C1'	6.35	1.49	1.41

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501[B]	IMP	N3-C2-N1	-11.32	120.23	128.89
2	A	501[B]	IMP	N3-C2-N1	-10.82	120.61	128.89
2	B	501[A]	IMP	N3-C2-N1	-10.35	120.97	128.89
3	A	502	NAJ	N3A-C2A-N1A	-10.00	121.23	128.89
2	A	501[A]	IMP	N3-C2-N1	-9.51	121.61	128.89
3	B	502	NAJ	N3A-C2A-N1A	-7.67	123.02	128.89
3	A	502	NAJ	PN-O3-PA	-4.26	120.78	132.73
2	B	501[B]	IMP	C1'-N9-C4	-3.47	121.70	126.94
2	B	501[A]	IMP	C1'-N9-C4	-3.11	122.25	126.94
2	A	501[A]	IMP	C4-C5-N7	-3.03	106.69	109.48
2	A	501[B]	IMP	C1'-N9-C4	-2.98	122.44	126.94
2	B	501[A]	IMP	C4'-O4'-C1'	-2.96	106.46	109.72
2	B	501[B]	IMP	C4'-O4'-C1'	-2.96	106.46	109.72
2	A	501[B]	IMP	C4-C5-N7	-2.76	106.94	109.48
3	B	502	NAJ	PN-O3-PA	-2.62	125.38	132.73
2	A	501[A]	IMP	C1'-N9-C4	-2.60	123.02	126.94
3	A	502	NAJ	C4B-O4B-C1B	-2.42	107.06	109.72
2	B	501[B]	IMP	C4-C5-N7	-2.26	107.40	109.48
3	B	502	NAJ	C4B-O4B-C1B	-2.25	107.25	109.72
2	B	501[B]	IMP	C6-C5-C4	-2.21	118.26	120.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	NAJ	O7N-C7N-C3N	-2.15	117.24	119.59
2	B	501[A]	IMP	C6-C5-C4	-2.10	118.39	120.90
3	B	502	NAJ	C4A-C5A-N7A	-2.09	107.56	109.48
2	A	501[B]	IMP	C4'-O4'-C1'	-2.05	107.47	109.72
2	A	501[A]	IMP	C4'-O4'-C1'	-2.05	107.47	109.72
3	B	502	NAJ	C2D-C3D-C4D	2.00	106.73	102.61
3	A	502	NAJ	O5D-C5D-C4D	2.19	117.19	109.12
2	B	501[A]	IMP	O3P-P-O1P	2.35	118.14	110.58
2	B	501[B]	IMP	O3P-P-O1P	2.35	118.14	110.58
3	B	502	NAJ	C4D-O4D-C1D	2.37	112.33	109.72
3	B	502	NAJ	O4D-C1D-N1N	2.45	110.82	108.13
3	A	502	NAJ	C2D-C3D-C4D	3.10	108.98	102.61
3	A	502	NAJ	C4D-O4D-C1D	3.34	113.39	109.72
3	B	502	NAJ	O4B-C1B-N9A	3.47	115.36	108.10
2	A	501[A]	IMP	C2-N1-C6	3.67	121.60	116.04
2	B	501[A]	IMP	C2-N1-C6	3.77	121.75	116.04
2	A	501[B]	IMP	C2-N1-C6	3.84	121.85	116.04
2	B	501[B]	IMP	C2-N1-C6	3.89	121.94	116.04
3	A	502	NAJ	O4D-C1D-N1N	4.52	113.10	108.13
3	A	502	NAJ	O4B-C1B-N9A	4.77	118.08	108.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501[A]	IMP	3	0
3	A	502	NAJ	3	0
2	B	501[A]	IMP	6	0
3	B	502	NAJ	8	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	342/366 (93%)	-0.03	7 (2%) 68 76	26, 38, 59, 79	1 (0%)
1	B	342/366 (93%)	0.13	8 (2%) 64 72	30, 44, 68, 85	1 (0%)
All	All	684/732 (93%)	0.05	15 (2%) 65 73	26, 41, 65, 85	2 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	7	ALA	6.9
1	B	327	ALA	3.8
1	A	79[A]	GLU	3.6
1	A	414	LEU	2.7
1	B	234[A]	GLU	2.7
1	B	434[A]	HIS	2.4
1	A	396	MET	2.3
1	B	24	THR	2.2
1	A	7	ALA	2.2
1	A	395	ALA	2.2
1	B	228	ALA	2.1
1	B	306[A]	ILE	2.1
1	B	273	LEU	2.1
1	A	228	ALA	2.0
1	A	488	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAJ	A	502	44/44	0.83	0.26	3.93	51,72,83,96	0
3	NAJ	B	502	44/44	0.83	0.23	2.09	48,73,94,104	0
4	K	B	503	1/1	0.99	0.12	-0.49	34,34,34,34	0
2	IMP	A	501[B]	23/23	0.97	0.13	-0.51	26,33,42,44	10
2	IMP	B	501[A]	23/23	0.96	0.12	-0.57	29,36,47,48	10
2	IMP	B	501[B]	23/23	0.96	0.12	-0.57	29,36,44,45	10
2	IMP	A	501[A]	23/23	0.97	0.13	-0.58	26,33,45,49	10
4	K	A	503	1/1	0.98	0.09	-3.68	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.