



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:04 PM GMT

PDB ID : 4QRR
Title : Crystal Structure of HLA B*3501-IPS in complex with a Delta-Beta TCR, clone 12 TCR
Authors : Gras, S.; Chabrol, E.; Rossjohn, J.
Deposited on : 2014-07-02
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	<div> <div>91%</div> <div>9%</div> </div>
2	B	99	<div> <div>88%</div> <div>12%</div> </div>
3	D	206	<div> <div>4%</div> <div>85%</div> <div>13%</div> <div>•</div> </div>
4	E	241	<div> <div>87%</div> <div>11%</div> <div>•</div> </div>
5	P	9	<div> <div>78%</div> <div>11%</div> <div>11%</div> </div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6696 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, B-35 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	276	Total	C	N	O	S	0	0	0
			2254	1405	411	431	7			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	0	0
			829	528	140	158	3			

- Molecule 3 is a protein called IPS peptide from CMV, IPSINVHHY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	203	Total	C	N	O	S	0	0	0
			1591	1010	261	312	8			

- Molecule 4 is a protein called clone12 TCR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	241	Total	C	N	O	S	0	0	0
			1922	1210	338	369	5			

- Molecule 5 is a protein called clone12 TCR beta chain.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	P	9	Total	C	N	O	0	0	0
			77	50	14	13			

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	11	Total	O	0	0
			11	11		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	2	Total	O	0	0
			2	2		
6	D	4	Total	O	0	0
			4	4		
6	E	6	Total	O	0	0
			6	6		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HLA class I histocompatibility antigen, B-35 alpha chain

Chain A: 





- Molecule 2: Beta-2-microglobulin

Chain B: 




- Molecule 3: IPS peptide from CMV, IPSINVHHY

Chain D: 




- Molecule 4: clone12 TCR alpha chain

Chain E: 



- Molecule 5: clone12 TCR beta chain

Chain P: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	65.96Å 192.15Å 162.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.95 – 3.00 45.95 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.95-3.00) 100.0 (45.95-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.94 (at 3.01Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.210 , 0.258 0.235 , 0.280	Depositor DCC
R_{free} test set	1085 reflections (5.41%)	DCC
Wilson B-factor (Å ²)	49.0	Xtriage
Anisotropy	0.570	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 33.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 21136 reflections	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	6696	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/2317	0.55	0/3150
2	B	0.37	0/852	0.54	0/1152
3	D	0.39	0/1626	0.58	0/2202
4	E	0.35	0/1975	0.54	0/2685
5	P	0.47	0/80	0.53	0/108
All	All	0.37	0/6850	0.55	0/9297

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2254	0	2121	16	0
2	B	829	0	796	9	0
3	D	1591	0	1543	16	0
4	E	1922	0	1823	16	0
5	P	77	0	74	1	0
6	A	11	0	0	0	0
6	B	2	0	0	0	0
6	D	4	0	0	0	0
6	E	6	0	0	0	0
All	All	6696	0	6357	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:63:TYR:O	2:B:64:LEU:HD12	1.48	1.12
3:D:202:CYS:SG	3:D:203:ALA:HA	2.01	1.00
4:E:45:THR:OG1	4:E:48:GLN:HB2	1.66	0.95
3:D:143:GLN:NE2	3:D:202:CYS:SG	2.48	0.87
1:A:228:THR:HG22	1:A:230:LEU:HD12	1.59	0.84
1:A:228:THR:HG22	1:A:230:LEU:CD1	2.10	0.80
2:B:63:TYR:C	2:B:64:LEU:HD12	2.05	0.76
1:A:215:LEU:HD12	1:A:215:LEU:O	1.88	0.73
3:D:202:CYS:HA	3:D:204:ASN:H	1.57	0.69
4:E:170:LEU:HG	4:E:225:VAL:HG22	1.75	0.68
4:E:45:THR:HG22	4:E:100:ALA:HB2	1.74	0.67
4:E:45:THR:CG2	4:E:100:ALA:HB2	2.25	0.66
3:D:13:MET:SD	3:D:14:PRO:HD2	2.37	0.64
3:D:202:CYS:SG	3:D:203:ALA:CA	2.85	0.63
5:P:1:ILE:HG13	5:P:2:PRO:HD2	1.84	0.59
1:A:228:THR:CG2	1:A:230:LEU:CD1	2.80	0.59
4:E:109:GLU:HG2	4:E:114:ASN:HD21	1.68	0.57
2:B:54:LEU:HD12	2:B:64:LEU:HD11	1.85	0.57
1:A:215:LEU:CD1	1:A:245:ALA:HB2	2.35	0.57
4:E:45:THR:HG22	4:E:100:ALA:CB	2.36	0.56
3:D:14:PRO:HG2	3:D:17:LYS:HG3	1.88	0.56
4:E:21:LEU:HD22	4:E:122:THR:HG21	1.88	0.55
2:B:24:ASN:HB3	2:B:65:LEU:HD11	1.87	0.55
1:A:5:MET:HB2	1:A:168:LEU:HD13	1.90	0.54
4:E:45:THR:OG1	4:E:48:GLN:CB	2.50	0.54
3:D:118:GLY:HA3	4:E:55:GLU:HG3	1.90	0.54
3:D:169:SER:N	3:D:170:ASP:HA	2.22	0.54
2:B:30:PHE:HZ	2:B:64:LEU:HD13	1.74	0.52
4:E:45:THR:CG2	4:E:100:ALA:CB	2.88	0.52
1:A:215:LEU:HD11	1:A:245:ALA:HB2	1.91	0.51
1:A:22:PHE:HB3	1:A:71:THR:HG22	1.92	0.50
3:D:165:GLN:HG2	3:D:165:GLN:O	2.11	0.50
1:A:228:THR:CG2	1:A:230:LEU:HD11	2.43	0.49
4:E:175:ASN:HA	4:E:220:HIS:HB3	1.95	0.49
1:A:14:ARG:HB3	1:A:17:ARG:HB2	1.95	0.48
1:A:69:THR:HG21	3:D:109:LEU:HD11	1.96	0.48
2:B:29:GLY:HA2	2:B:61:SER:HB2	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:103:LEU:HD13	1:A:168:LEU:HD23	1.95	0.47
1:A:189:VAL:HG21	1:A:274:TRP:HA	1.96	0.47
4:E:222:ARG:HG3	4:E:251:GLU:HB3	1.97	0.47
2:B:39:LEU:HD13	2:B:49:VAL:HG21	2.00	0.44
3:D:168:ASP:HA	3:D:171:VAL:H	1.82	0.44
1:A:189:VAL:HG13	1:A:272:LEU:HD23	2.00	0.44
4:E:23:CYS:SG	4:E:104:CYS:SG	3.11	0.43
2:B:39:LEU:HD22	2:B:68:THR:HG22	2.00	0.43
4:E:158:CYS:SG	4:E:223:CYS:SG	3.08	0.43
3:D:112:ALA:HA	3:D:113:GLY:HA2	1.69	0.43
1:A:123:TYR:HD2	1:A:124:ILE:HG22	1.84	0.42
4:E:228:TYR:HA	4:E:245:THR:HG23	2.00	0.42
2:B:23:LEU:HD23	2:B:39:LEU:HD23	2.01	0.42
3:D:105:ALA:HB1	3:D:120:LEU:HB3	2.02	0.41
4:E:6:GLN:HG3	4:E:23:CYS:HB2	2.02	0.41
3:D:23:CYS:HG	3:D:104:CYS:HG	1.68	0.41
1:A:123:TYR:CZ	1:A:140:ALA:HA	2.55	0.41
3:D:43:LYS:HB2	3:D:53:LEU:HD11	2.02	0.40
3:D:19:VAL:HB	3:D:94:LEU:HD11	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/276 (99%)	263 (96%)	8 (3%)	3 (1%)	17	58
2	B	97/99 (98%)	95 (98%)	2 (2%)	0	100	100
3	D	199/206 (97%)	177 (89%)	20 (10%)	2 (1%)	19	61
4	E	239/241 (99%)	224 (94%)	12 (5%)	3 (1%)	15	53
5	P	7/9 (78%)	5 (71%)	2 (29%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	816/831 (98%)	764 (94%)	44 (5%)	8 (1%)	19	61

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	15	PRO
1	A	195	SER
3	D	197	LYS
4	E	175	ASN
1	A	29	ASP
3	D	149	LYS
4	E	240	ARG
4	E	241	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/234 (100%)	229 (98%)	5 (2%)	61	89
2	B	94/94 (100%)	94 (100%)	0	100	100
3	D	177/180 (98%)	175 (99%)	2 (1%)	80	94
4	E	210/210 (100%)	199 (95%)	11 (5%)	29	68
5	P	9/9 (100%)	8 (89%)	1 (11%)	8	29
All	All	724/727 (100%)	705 (97%)	19 (3%)	54	85

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	ASP
1	A	155	GLN
1	A	164	CYS
1	A	166	GLU
1	A	189	VAL
3	D	159	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	D	170	ASP
4	E	11	LEU
4	E	23	CYS
4	E	54	PHE
4	E	108	LEU
4	E	116	GLN
4	E	158	CYS
4	E	170	LEU
4	E	196	LEU
4	E	206	ARG
4	E	233	ASN
4	E	234	ASP
5	P	1	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	197	HIS
2	B	31	HIS
3	D	143	GLN
3	D	208	ASN
4	E	48	GLN
4	E	61	GLN
4	E	114	ASN
4	E	238	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	276/276 (100%)	-0.20	0	100	100	22, 45, 92, 109	1 (0%)
2	B	99/99 (100%)	-0.27	0	100	100	35, 50, 71, 81	0
3	D	203/206 (98%)	-0.07	8 (3%)	43	18	21, 46, 103, 114	6 (2%)
4	E	241/241 (100%)	-0.21	0	100	100	25, 47, 73, 84	0
5	P	9/9 (100%)	-0.38	0	100	100	31, 33, 40, 40	0
All	All	828/831 (99%)	-0.18	8 (0%)	84	60	21, 47, 91, 114	7 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	147	SER	3.9
3	D	199	ASP	3.2
3	D	148	ASP	2.9
3	D	216	PHE	2.9
3	D	169	SER	2.8
3	D	214	ASP	2.8
3	D	165	GLN	2.4
3	D	163	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.