



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:20 PM GMT

PDB ID : 3QUZ  
Title : Structure of the mouse CD1d-NU-alpha-GalCer-iNKT TCR complex  
Authors : Li, Y.; Girardi, E.; Yu, E.D.; Zajonc, D.M.  
Deposited on : 2011-02-24  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	285	<div> <div>5%</div> <div>82%</div> <div>10%</div> <div>7%</div> </div>
2	B	99	<div> <div>%</div> <div>87%</div> <div>8%</div> <div>.</div> </div>
3	C	209	<div> <div>7%</div> <div>90%</div> <div>7%</div> <div>.</div> </div>
4	D	241	<div> <div>5%</div> <div>91%</div> <div>7%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	NAG	A	511	-	-	-	X
6	FUC	A	513	-	-	-	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6718 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Antigen-presenting glycoprotein CD1d1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	266	Total	C	N	O	S	0	0	0
			2132	1363	364	392	13			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	201	HIS	ASP	SEE REMARK 999	UNP P11609
A	280	HIS	-	EXPRESSION TAG	UNP P11609
A	281	HIS	-	EXPRESSION TAG	UNP P11609
A	282	HIS	-	EXPRESSION TAG	UNP P11609
A	283	HIS	-	EXPRESSION TAG	UNP P11609
A	284	HIS	-	EXPRESSION TAG	UNP P11609
A	285	HIS	-	EXPRESSION TAG	UNP P11609

- Molecule 2 is a protein called Beta-2 microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	96	Total	C	N	O	S	0	1	0
			784	503	135	140	6			

- Molecule 3 is a protein called Valpha14 (mouse variable domain, human constant domain).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	203	Total	C	N	O	S	0	1	0
			1561	968	269	316	8			

- Molecule 4 is a protein called Vbeta8.2 (mouse variable domain, human constant domain).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	239	Total	C	N	O	S	0	0	0
			1876	1177	334	359	6			

- Molecule 5 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	3	Total	C	N	O	0	0
			38	22	2	14		

- Molecule 7 is N-[(2S,3S,4R)-1-({6-DEOXY-6-[(NAPHTHALEN-1-YLCARBAMOYL)AMINO]-ALPHA-D-GALACTOPYRANOSYL}OXY)-3,4-DIHYDROXYOCTADECAN-2-YL]HEXACOSANAMIDE (three-letter code: QUV) (formula:  $C_{61}H_{107}N_3O_9$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			73	61	3	9		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	85	Total	O	0	0
			85	85		
8	B	28	Total	O	0	0
			28	28		
8	C	50	Total	O	0	0
			50	50		
8	D	63	Total	O	0	0
			63	63		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Antigen-presenting glycoprotein CD1d1




- Molecule 2: Beta-2 microglobulin



- Molecule 3: Valpha14 (mouse variable domain, human constant domain)



- Molecule 4: Vbeta8.2 (mouse variable domain, human constant domain)



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.29Å 190.98Å 151.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.38 – 2.30 35.38 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.5 (35.38-2.30) 99.5 (35.38-2.30)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.20 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.192 , 0.233 0.188 , 0.228	Depositor DCC
$R_{free}$ test set	2598 reflections (5.36%)	DCC
Wilson B-factor (Å <sup>2</sup> )	38.3	Xtriage
Anisotropy	0.029	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 38.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 51098 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6718	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, NAG, QUV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.76	0/2194	0.74	3/2984 (0.1%)
2	B	0.68	0/814	0.72	0/1110
3	C	0.72	1/1593 (0.1%)	0.76	2/2167 (0.1%)
4	D	0.75	1/1927 (0.1%)	0.71	1/2627 (0.0%)
All	All	0.74	2/6528 (0.0%)	0.73	6/8888 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	54	SER	CB-OG	6.33	1.50	1.42
3	C	16	GLU	CG-CD	5.13	1.59	1.51

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	79	ARG	NE-CZ-NH2	-7.95	116.32	120.30
1	A	79	ARG	NE-CZ-NH1	6.93	123.77	120.30
3	C	60	ARG	NE-CZ-NH1	6.88	123.74	120.30
3	C	60	ARG	NE-CZ-NH2	-6.69	116.95	120.30
4	D	207	CYS	CA-CB-SG	-5.32	104.43	114.00
1	A	80	ASP	CB-CG-OD1	5.26	123.03	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2132	0	2042	20	0
2	B	784	0	747	9	0
3	C	1561	0	1493	9	0
4	D	1876	0	1780	11	0
5	A	28	0	26	0	0
6	A	38	0	34	0	0
7	A	73	0	107	1	0
8	A	85	0	0	2	0
8	B	28	0	0	1	0
8	C	50	0	0	3	0
8	D	63	0	0	0	0
All	All	6718	0	6229	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:88:MET:CE	1:A:145:LEU:HD23	2.08	0.83
1:A:88:MET:HE1	1:A:145:LEU:HD23	1.68	0.75
2:B:67:HIS:ND1	8:B:112:HOH:O	2.20	0.73
2:B:96:ASP:O	2:B:97:ARG:HG3	1.90	0.71
1:A:88:MET:HE3	1:A:145:LEU:HD23	1.75	0.66
3:C:82:LEU:HD22	3:C:174:LYS:HD3	1.79	0.63
4:D:204:HIS:NE2	4:D:235:GLU:CG	2.62	0.62
3:C:60:ARG:NH2	3:C:84:ASP:OD2	2.32	0.62
2:B:96:ASP:O	2:B:97:ARG:CB	2.49	0.61
4:D:204:HIS:NE2	4:D:235:GLU:HG3	2.16	0.61
2:B:96:ASP:O	2:B:97:ARG:CG	2.54	0.56
1:A:258:GLU:HB3	1:A:279:TRP:CD1	2.41	0.56
1:A:35:LEU:HD12	1:A:183:LEU:HD23	1.89	0.55
1:A:234:ARG:HD2	8:A:292:HOH:O	2.06	0.54
3:C:166:LEU:HB3	4:D:168:CYS:HB3	1.89	0.54
1:A:79:ARG:HD2	8:A:289:HOH:O	2.08	0.53
1:A:107:TYR:HB3	1:A:108:PRO:HD2	1.91	0.52
1:A:219:TRP:CZ2	1:A:221:MET:HG3	2.45	0.51
3:C:33:ARG:HD2	3:C:35:PHE:CE2	2.45	0.51
2:B:21:ASN:HB3	2:B:70:PHE:CE1	2.47	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:69:MET:HB3	7:A:286:QUV:CCL	2.42	0.50
4:D:204:HIS:NE2	4:D:235:GLU:HG2	2.27	0.49
4:D:140:LEU:HD12	4:D:140:LEU:N	2.28	0.48
1:A:191:ALA:HA	1:A:209:HIS:O	2.15	0.47
4:D:206:ARG:HD3	4:D:233:SER:OG	2.14	0.47
1:A:127:ARG:NE	1:A:134:GLN:OE1	2.46	0.46
2:B:96:ASP:O	2:B:97:ARG:HB2	2.16	0.46
3:C:1:THR:N	8:C:231:HOH:O	2.49	0.46
2:B:14:PRO:HA	2:B:15:PRO:HD3	1.71	0.45
4:D:113:ASP:OD1	4:D:115:ARG:HG2	2.16	0.45
1:A:221:MET:HE2	1:A:228:GLU:HB3	1.99	0.45
4:D:117:VAL:HG12	4:D:227:PRO:HB2	1.98	0.45
1:A:196:VAL:O	1:A:205:GLN:N	2.50	0.45
1:A:110:ASN:OD1	1:A:110:ASN:N	2.38	0.45
1:A:124:TYR:CZ	1:A:136:VAL:HG11	2.52	0.45
4:D:7:SER:HA	4:D:8:PRO:C	2.38	0.44
3:C:103:ARG:HD3	8:C:228:HOH:O	2.18	0.44
4:D:2:ALA:HA	4:D:27:ASN:OD1	2.18	0.44
2:B:46:ILE:HA	2:B:47:PRO:HD3	1.89	0.44
3:C:103:ARG:NH2	8:C:242:HOH:O	2.51	0.42
1:A:15:MET:HG2	2:B:62:PHE:HE2	1.84	0.42
1:A:106:MET:CE	1:A:173:ARG:HG3	2.51	0.41
1:A:59:SER:OG	1:A:62:GLN:HG3	2.21	0.41
3:C:85:THR:O	3:C:86:ALA:HB2	2.20	0.41
3:C:159:TYR:O	3:C:180:ALA:HA	2.20	0.41
4:D:121:LYS:O	4:D:144:ALA:HA	2.20	0.41
1:A:219:TRP:HB3	1:A:266:LYS:HB2	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/285 (92%)	257 (98%)	5 (2%)	0	100	100
2	B	95/99 (96%)	93 (98%)	2 (2%)	0	100	100
3	C	202/209 (97%)	194 (96%)	8 (4%)	0	100	100
4	D	237/241 (98%)	231 (98%)	5 (2%)	1 (0%)	39	48
All	All	796/834 (95%)	775 (97%)	20 (2%)	1 (0%)	56	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	127	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/249 (93%)	227 (98%)	4 (2%)	68	83
2	B	87/93 (94%)	85 (98%)	2 (2%)	58	75
3	C	179/188 (95%)	178 (99%)	1 (1%)	90	96
4	D	203/208 (98%)	199 (98%)	4 (2%)	63	79
All	All	700/738 (95%)	689 (98%)	11 (2%)	70	84

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	SER
1	A	110	ASN
1	A	196	VAL
1	A	226	ASP
2	B	70	PHE
2	B	97	ARG
3	C	165	VAL
4	D	54	SER
4	D	168	CYS
4	D	235	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
4	D	239	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

3 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
6	NAG	A	511	1,6	14,14,15	0.69	0	15,19,21	1.36	2 (13%)
6	NAG	A	512	6	14,14,15	0.57	0	15,19,21	1.10	0
6	FUC	A	513	6	10,10,11	0.90	0	14,14,16	1.41	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	511	1,6	-	0/6/23/26	0/1/1/1
6	NAG	A	512	6	-	0/6/23/26	0/1/1/1
6	FUC	A	513	6	-	0/0/17/20	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	513	FUC	O5-C1-C2	-2.47	106.85	110.86
6	A	511	NAG	O5-C5-C6	2.17	112.04	107.35
6	A	513	FUC	O5-C5-C6	2.35	110.02	106.13
6	A	513	FUC	O3-C3-C2	2.59	114.68	110.00
6	A	511	NAG	C1-O5-C5	3.58	116.79	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
7	QUV	A	286	-	74,75,75	0.88	5 (6%)	86,90,90	1.04	4 (4%)
5	NAG	A	500	1	14,14,15	0.90	1 (7%)	15,19,21	1.96	1 (6%)
5	NAG	A	501	1	14,14,15	0.75	0	15,19,21	1.78	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	QUV	A	286	-	-	0/65/85/85	0/3/3/3
5	NAG	A	500	1	-	0/6/23/26	0/1/1/1
5	NAG	A	501	1	-	0/6/23/26	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	286	QUV	CCI-NCJ	-2.12	1.33	1.37
7	A	286	QUV	CCU-NCJ	-2.09	1.35	1.41
7	A	286	QUV	CCS-CCR	2.06	1.41	1.36
7	A	286	QUV	O4-C4	2.09	1.48	1.43
7	A	286	QUV	CCT-CCU	2.13	1.43	1.38
5	A	500	NAG	C1-C2	3.00	1.56	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	501	NAG	C3-C4-C5	-2.72	105.45	110.20
7	A	286	QUV	C2-N2-CAA	-2.48	119.02	123.61
7	A	286	QUV	CCT-CCU-NCJ	-2.21	116.86	123.10
7	A	286	QUV	CCR-CCQ-CCP	-2.20	118.02	123.22
5	A	501	NAG	O5-C5-C6	2.44	112.64	107.35
5	A	501	NAG	O4-C4-C5	2.72	116.44	109.24
7	A	286	QUV	CCL-CCU-NCJ	3.55	124.79	118.42
5	A	501	NAG	C1-O5-C5	4.03	117.36	112.25
5	A	500	NAG	C1-O5-C5	6.76	120.83	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	286	QUV	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	266/285 (93%)	0.16	13 (4%) 33 42	19, 35, 97, 129	0
2	B	96/99 (96%)	-0.04	1 (1%) 84 88	25, 43, 65, 77	0
3	C	203/209 (97%)	0.30	15 (7%) 17 25	19, 38, 99, 122	0
4	D	239/241 (99%)	0.11	12 (5%) 32 41	21, 36, 63, 95	0
All	All	804/834 (96%)	0.16	41 (5%) 32 41	19, 37, 87, 129	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	185	SER	8.4
4	D	3	ALA	7.1
1	A	197	PRO	5.9
4	D	181	ASN	4.8
4	D	180	LEU	4.8
3	C	153	SER	4.7
3	C	183	ASN	4.7
3	C	186	ASP	4.4
4	D	179	ALA	4.4
3	C	184	LYS	4.4
3	C	134	SER	4.4
3	C	154	LYS	4.2
3	C	195	ASN	4.0
4	D	2	ALA	4.0
3	C	157	ASP	3.6
1	A	259	ALA	3.6
4	D	223	ASP	3.5
3	C	119	GLN	3.5
1	A	260	GLY	3.4
1	A	110	ASN	3.3
1	A	254	GLU	3.2

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	256	GLY	3.2
4	D	217	ASN	3.2
1	A	257	GLU	3.2
3	C	155	ASP	3.1
4	D	240	ALA	3.0
2	B	88	ALA	2.8
3	C	203	PHE	2.8
3	C	201	ASP	2.7
4	D	178	PRO	2.7
1	A	255	ALA	2.7
3	C	194	ASN	2.6
1	A	227	GLN	2.5
4	D	80	LEU	2.5
1	A	226	ASP	2.4
4	D	216	GLU	2.3
1	A	225	GLY	2.2
4	D	177	GLN	2.2
1	A	230	GLN	2.1
3	C	190	ALA	2.1
1	A	118	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	FUC	A	513	10/11	0.83	0.22	2.35	46,49,52,56	0
6	NAG	A	511	14/15	0.95	0.16	2.31	32,38,47,52	0
6	NAG	A	512	14/15	0.90	0.30	-	47,58,67,67	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
7	QUV	A	286	73/73	0.92	0.19	1.23	19,32,43,48	0
5	NAG	A	500	14/15	0.87	0.19	-	57,67,74,76	0
5	NAG	A	501	14/15	0.92	0.14	-	37,41,44,48	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.