



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:52 PM GMT

PDB ID : 1R0R
Title : 1.1 Angstrom Resolution Structure of the Complex Between the Protein Inhibitor, OMTKY3, and the Serine Protease, Subtilisin Carlsberg
Authors : Horn, J.R.; Ramaswamy, S.; Murphy, K.P.
Deposited on : 2003-09-22
Resolution : 1.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1006 (1.14-1.06)
Clashscore	102246	1055 (1.14-1.06)
Ramachandran outliers	100387	1016 (1.14-1.06)
Sidechain outliers	100360	1014 (1.14-1.06)
RSRZ outliers	91569	1009 (1.14-1.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	274	<div> <div>7%</div> <div>96%</div> <div>••</div> </div>
2	I	51	<div> <div>12%</div> <div>80%</div> <div>16%</div> <div>•</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CA	E	301	-	-	-	X
3	CA	E	302	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2773 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called subtilisin carlsberg.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	274	Total	C	N	O	S	8	14	0
			1962	1216	335	405	6			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	158	ASN	SER	CONFLICT	UNP P00780
E	161	SER	ASN	CONFLICT	UNP P00780
E	225	PRO	GLU	CONFLICT	UNP P00780

- Molecule 2 is a protein called Ovomucoid.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	51	Total	C	N	O	S	21	3	0
			400	249	67	78	6			

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	4	Total	Ca	0	2
			6	6		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	324	Total	O	0	0
			324	324		
4	I	81	Total	O	0	0
			81	81		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: subtilisin carlsberg

Chain E: 



- Molecule 2: Ovomucoid

Chain I: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	62.33Å 70.81Å 127.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.25 – 1.10 31.16 – 1.10	Depositor EDS
% Data completeness (in resolution range)	94.1 (15.25-1.10) 94.2 (31.16-1.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.02 (at 1.10Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.158 , 0.184 0.168 , 0.195	Depositor DCC
R_{free} test set	1090 reflections (1.01%)	DCC
Wilson B-factor (Å ²)	10.4	Xtriage
Anisotropy	0.712	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 66.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 108224 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2773	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	0.60	2/2053 (0.1%)	0.74	3/2797 (0.1%)
2	I	1.48	7/421 (1.7%)	1.37	6/566 (1.1%)
All	All	0.82	9/2474 (0.4%)	0.88	9/3363 (0.3%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	10	GLU	CB-CG	15.70	1.81	1.52
2	I	21	ARG	CZ-NH2	-12.72	1.16	1.33
2	I	21	ARG	CZ-NH1	10.84	1.47	1.33
1	E	158	ASN	CG-ND2	-7.38	1.14	1.32
2	I	34[C]	LYS	CE-NZ	7.30	1.67	1.49
2	I	34[D]	LYS	CE-NZ	7.30	1.67	1.49
1	E	158	ASN	CG-OD1	7.08	1.39	1.24
2	I	29[A]	LYS	CE-NZ	-5.75	1.34	1.49
2	I	29[B]	LYS	CE-NZ	-5.75	1.34	1.49

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	21	ARG	NE-CZ-NH1	-18.81	110.90	120.30
2	I	21	ARG	NE-CZ-NH2	13.34	126.97	120.30
2	I	10	GLU	CA-CB-CG	-7.56	96.77	113.40
2	I	7	ASP	CB-CG-OD1	7.00	124.60	118.30
1	E	158	ASN	CB-CG-OD1	-6.73	108.14	121.60
2	I	34[C]	LYS	CD-CE-NZ	-5.72	98.55	111.70
2	I	34[D]	LYS	CD-CE-NZ	-5.72	98.55	111.70
1	E	172	ASP	CB-CG-OD2	5.23	123.01	118.30
1	E	14	ASP	CB-CG-OD2	5.10	122.89	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1962	0	1911	10	0
2	I	400	0	383	8	0
3	E	6	0	0	0	0
4	E	324	0	0	3	0
4	I	81	0	0	0	0
All	All	2773	0	2294	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:263[B]:TYR:OH	4:E:4165:HOH:O	1.61	1.18
1:E:248[A]:ASN:OD1	4:E:4413:HOH:O	1.66	1.13
1:E:3[A]:THR:OG1	4:E:4566:HOH:O	1.66	1.09
1:E:104:TYR:CD1	1:E:135[B]:MET:CE	2.58	0.87
2:I:23[B]:LEU:CD2	2:I:53:PHE:CD1	2.59	0.86
1:E:104:TYR:CD1	1:E:135[B]:MET:HE3	2.11	0.84
2:I:23[B]:LEU:HD21	2:I:53:PHE:CD1	2.22	0.74
2:I:23[B]:LEU:HD22	2:I:53:PHE:CD1	2.24	0.72
2:I:23[B]:LEU:CD2	2:I:53:PHE:HD1	2.07	0.66
1:E:104:TYR:HD1	1:E:135[B]:MET:CE	2.10	0.61
2:I:11:TYR:HB3	2:I:12:PRO:HA	1.88	0.54
1:E:104:TYR:CE1	1:E:135[B]:MET:HE3	2.44	0.53
1:E:104:TYR:CD1	1:E:135[B]:MET:HE1	2.42	0.52
1:E:104:TYR:CE1	1:E:135[B]:MET:CE	2.94	0.50
2:I:23[B]:LEU:HD21	2:I:53:PHE:HD1	1.67	0.47
2:I:23[B]:LEU:HD21	2:I:53:PHE:CE1	2.50	0.46
2:I:6:VAL:CG1	2:I:34[C]:LYS:HE3	2.48	0.43
1:E:15:LYS:HA	1:E:15:LYS:HD2	1.91	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	286/274 (104%)	280 (98%)	6 (2%)	0	100	100
2	I	52/51 (102%)	51 (98%)	1 (2%)	0	100	100
All	All	338/325 (104%)	331 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	212/198 (107%)	208 (98%)	4 (2%)	65	24
2	I	48/45 (107%)	47 (98%)	1 (2%)	61	19
All	All	260/243 (107%)	255 (98%)	5 (2%)	70	24

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	3[A]	THR
1	E	3[B]	THR
1	E	158	ASN
1	E	222	MET
2	I	10	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	240	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	E	274/274 (100%)	0.48	18 (6%)	22 20	12, 14, 20, 24	4 (1%)
2	I	51/51 (100%)	0.65	6 (11%)	6 8	13, 16, 24, 27	7 (13%)
All	All	325/325 (100%)	0.51	24 (7%)	17 17	12, 15, 21, 27	11 (3%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	I	56	CYS	6.6
1	E	161[A]	SER	4.0
1	E	275	GLN	3.9
1	E	240	ASN	3.8
1	E	160	GLY	3.3
2	I	9	SER	3.2
2	I	6	VAL	3.2
1	E	18	ALA	3.2
2	I	53	PHE	3.0
1	E	130	SER	2.7
1	E	159	SER	2.6
2	I	10	GLU	2.5
1	E	162	THR	2.5
1	E	104	TYR	2.5
1	E	68	VAL	2.4
1	E	248[A]	ASN	2.4
1	E	21	PHE	2.3
1	E	256	TYR	2.3
1	E	238	HIS	2.2
1	E	241	LEU	2.1
1	E	260	SER	2.1
1	E	72	VAL	2.1
1	E	272	ALA	2.1
2	I	8	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	CA	E	302	1/1	1.00	0.19	5.52	19,19,19,19	0
3	CA	E	301	1/1	0.99	0.14	3.78	22,22,22,22	0
3	CA	E	305[A]	1/1	0.99	0.11	1.49	42,42,42,42	1
3	CA	E	305[B]	1/1	0.99	0.11	1.32	22,22,22,22	1
3	CA	E	303[B]	1/1	0.99	0.11	0.17	26,26,26,26	1
3	CA	E	303[A]	1/1	0.99	0.11	0.09	27,27,27,27	1

6.5 Other polymers [i](#)

There are no such residues in this entry.