



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:30 AM GMT

PDB ID : 2R16  
Title : Crystal Structure of bovine neuexin 1 alpha LNS/LG domain 4 (with no splice insert)  
Authors : Rudenko, G.  
Deposited on : 2007-08-22  
Resolution : 1.04 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1032 (1.10-0.98)
Clashscore	102246	1109 (1.10-0.98)
Ramachandran outliers	100387	1041 (1.10-0.98)
Sidechain outliers	100360	1039 (1.10-0.98)
RSRZ outliers	91569	1034 (1.10-0.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	182	<div> <div>19%</div> <div>79%</div> <div>18%</div> <div>.</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1607 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neurexin-1-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	175	Total	C	N	O	S	0	23	0
			1476	924	257	281	14			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	719	GLY	-	EXPRESSION TAG	UNP Q28146
A	720	SER	-	EXPRESSION TAG	UNP Q28146
A	?	-	ASP	DELETION	UNP Q28146
A	?	-	CYS	DELETION	UNP Q28146
A	?	-	ILE	DELETION	UNP Q28146
A	?	-	ARG	DELETION	UNP Q28146
A	?	-	ILE	DELETION	UNP Q28146
A	?	-	ASN	DELETION	UNP Q28146
A	?	-	CYS	DELETION	UNP Q28146
A	?	-	ASN	DELETION	UNP Q28146
A	?	-	SER	DELETION	UNP Q28146
A	790	GLY	SER	EXPRESSION TAG	UNP Q28146

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	130	Total	O	0	0
			130	130		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Neurexin-1-alpha



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.32Å 90.32Å 39.62Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	23.70 – 1.04 23.70 – 1.04	Depositor EDS
% Data completeness (in resolution range)	100.0 (23.70-1.04) 91.4 (23.70-1.04)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.17 (at 1.04Å)	Xtriage
Refinement program	REFMAC 5.3	Depositor
R, $R_{free}$	0.146 , 0.165 0.153 , 0.169	Depositor DCC
$R_{free}$ test set	8111 reflections (10.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	9.1	Xtriage
Anisotropy	0.141	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 41.0	EDS
Estimated twinning fraction	0.026 for h,-h-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 80874 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	1607	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.66	0/1558	0.90	2/2100 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	763[A]	MET	CG-SD-CE	-7.86	87.62	100.20
1	A	763[B]	MET	CG-SD-CE	-7.86	87.62	100.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1476	0	1455	38	0
2	A	1	0	0	0	0
3	A	130	0	0	6	0
All	All	1607	0	1455	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:803[B]:GLU:HG2	1:A:842[B]:MET:CE	1.65	1.26
1:A:803[B]:GLU:CG	1:A:842[B]:MET:HE1	1.66	1.25
1:A:803[B]:GLU:CD	1:A:842[B]:MET:HE3	1.71	1.11
1:A:890[B]:CYS:SG	1:A:900:LEU:CD2	2.54	0.95
1:A:803[B]:GLU:CD	1:A:842[B]:MET:CE	2.34	0.95
1:A:803[B]:GLU:CG	1:A:842[B]:MET:CE	2.31	0.94
1:A:890[B]:CYS:SG	1:A:900:LEU:HD23	2.08	0.94
1:A:862:ARG:NH1	1:A:868:VAL:HG13	1.97	0.80
1:A:768:ARG:H	1:A:852:HIS:HD2	1.32	0.78
1:A:774[A]:LEU:HD12	1:A:787:VAL:HG22	1.66	0.77
1:A:803[B]:GLU:HG2	1:A:842[B]:MET:HE1	0.82	0.76
1:A:818[B]:THR:HG22	1:A:833:ASP:HA	1.67	0.74
1:A:768:ARG:H	1:A:852:HIS:CD2	2.08	0.71
1:A:774[A]:LEU:CD1	1:A:787:VAL:HG22	2.22	0.70
1:A:786:THR:HG21	3:A:1109:HOH:O	1.93	0.68
1:A:802:PRO:HB3	3:A:1090:HOH:O	2.00	0.61
1:A:803[B]:GLU:OE2	1:A:842[B]:MET:HE3	2.03	0.58
1:A:862:ARG:O	1:A:865:LEU:HB3	2.05	0.56
1:A:746:ALA:HB1	1:A:824:ARG:HE	1.70	0.55
1:A:862:ARG:HH12	1:A:868:VAL:HG13	1.71	0.54
1:A:809:TYR:HD1	1:A:810:ASN:ND2	2.09	0.50
1:A:836[B]:GLN:HG2	1:A:836[B]:GLN:O	2.11	0.50
1:A:836[B]:GLN:O	1:A:836[B]:GLN:CG	2.57	0.49
1:A:890[B]:CYS:SG	1:A:900:LEU:HD21	2.49	0.48
1:A:818[B]:THR:HG23	3:A:1014:HOH:O	2.13	0.48
1:A:895:ILE:HB	1:A:897[B]:TYR:CE2	2.50	0.46
1:A:787:VAL:HG21	1:A:828:LEU:HD11	1.97	0.46
1:A:818[B]:THR:CG2	1:A:833:ASP:HA	2.39	0.46
1:A:784:LYS:HE3	1:A:804[B]:THR:HG21	1.97	0.45
1:A:862:ARG:O	1:A:865:LEU:CB	2.65	0.45
1:A:852:HIS:HE1	3:A:1101:HOH:O	1.99	0.44
1:A:880[B]:THR:HG22	3:A:1098:HOH:O	2.17	0.44
1:A:862:ARG:HA	1:A:865:LEU:HB2	2.02	0.42
1:A:862:ARG:NH2	3:A:1087:HOH:O	2.51	0.41
1:A:829:LYS:CG	1:A:837:ALA:HB1	2.51	0.41
1:A:859:ILE:O	1:A:862:ARG:CZ	2.69	0.41
1:A:739:PRO:HD3	1:A:897[A]:TYR:CE1	2.56	0.40
1:A:895:ILE:HD12	1:A:897[B]:TYR:CE1	2.55	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	196/182 (108%)	194 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/156 (108%)	168 (99%)	1 (1%)	90	66

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	757	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	743	HIS
1	A	810	ASN
1	A	852	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.



## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	175/182 (96%)	1.19	34 (19%) 1 5	8, 13, 20, 25	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	906	PHE	11.2
1	A	723	ALA	8.7
1	A	809	TYR	5.4
1	A	843	ALA	4.8
1	A	762	LEU	3.9
1	A	905	GLY	3.8
1	A	866	SER	3.7
1	A	764	ALA	3.6
1	A	865	LEU	3.4
1	A	845	ASP	3.3
1	A	776	LEU	3.3
1	A	761	ILE	3.2
1	A	774[A]	LEU	3.2
1	A	844	GLY	3.2
1	A	763[A]	MET	3.1
1	A	868	VAL	3.1
1	A	863	ARG	3.0
1	A	867	SER	2.9
1	A	751[A]	LEU	2.9
1	A	739	PRO	2.8
1	A	841	GLN	2.7
1	A	810	ASN	2.6
1	A	856	THR	2.6
1	A	854	ILE	2.5
1	A	736	ILE	2.4
1	A	802	PRO	2.4
1	A	824	ARG	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	785	LEU	2.4
1	A	765	THR	2.3
1	A	862	ARG	2.2
1	A	898[A]	CYS	2.2
1	A	896	ASP	2.1
1	A	800[A]	LYS	2.1
1	A	876	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CA	A	1000	1/1	0.99	0.06	-2.28	6,6,6,6	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.