



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:24 PM GMT

PDB ID : 3R79  
Title : Crystal structure of an uncharacterized protein from Agrobacterium tumefaciens  
Authors : Eswaramoorthy, S.; Almo, S.C.; Swaminathan, S.; New York Structural Genomics Research Consortium (NYSGRG)  
Deposited on : 2011-03-22  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	244	<div> <div>2%</div> <div> <div></div> <div>81%</div> <div>13%</div> <div>7%</div> </div> </div>
1	B	244	<div> <div>2%</div> <div> <div></div> <div>81%</div> <div>10%</div> <div>9%</div> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3606 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	228	Total	C	N	O	S	Se	0	0	0
			1738	1088	308	335	4	3			
1	B	221	Total	C	N	O	S	Se	0	0	0
			1683	1056	298	322	4	3			

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MSE	-	EXPRESSION TAG	UNP A9CHE9
A	0	VAL	-	EXPRESSION TAG	UNP A9CHE9
A	1	VAL	-	EXPRESSION TAG	UNP A9CHE9
A	221	ALA	-	EXPRESSION TAG	UNP A9CHE9
A	222	GLU	-	EXPRESSION TAG	UNP A9CHE9
A	223	ASN	-	EXPRESSION TAG	UNP A9CHE9
A	224	LEU	-	EXPRESSION TAG	UNP A9CHE9
A	225	TYR	-	EXPRESSION TAG	UNP A9CHE9
A	226	PHE	-	EXPRESSION TAG	UNP A9CHE9
A	227	GLN	-	EXPRESSION TAG	UNP A9CHE9
A	228	SER	-	EXPRESSION TAG	UNP A9CHE9
A	229	HIS	-	EXPRESSION TAG	UNP A9CHE9
A	230	HIS	-	EXPRESSION TAG	UNP A9CHE9
A	231	HIS	-	EXPRESSION TAG	UNP A9CHE9
A	232	HIS	-	EXPRESSION TAG	UNP A9CHE9
A	233	HIS	-	EXPRESSION TAG	UNP A9CHE9
A	234	HIS	-	EXPRESSION TAG	UNP A9CHE9
A	235	TRP	-	EXPRESSION TAG	UNP A9CHE9
A	236	SER	-	EXPRESSION TAG	UNP A9CHE9
A	237	HIS	-	EXPRESSION TAG	UNP A9CHE9
A	238	PRO	-	EXPRESSION TAG	UNP A9CHE9
A	239	GLN	-	EXPRESSION TAG	UNP A9CHE9
A	240	PHE	-	EXPRESSION TAG	UNP A9CHE9
A	241	GLU	-	EXPRESSION TAG	UNP A9CHE9
A	242	LYS	-	EXPRESSION TAG	UNP A9CHE9

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	MSE	-	EXPRESSION TAG	UNP A9CHE9
B	0	VAL	-	EXPRESSION TAG	UNP A9CHE9
B	1	VAL	-	EXPRESSION TAG	UNP A9CHE9
B	221	ALA	-	EXPRESSION TAG	UNP A9CHE9
B	222	GLU	-	EXPRESSION TAG	UNP A9CHE9
B	223	ASN	-	EXPRESSION TAG	UNP A9CHE9
B	224	LEU	-	EXPRESSION TAG	UNP A9CHE9
B	225	TYR	-	EXPRESSION TAG	UNP A9CHE9
B	226	PHE	-	EXPRESSION TAG	UNP A9CHE9
B	227	GLN	-	EXPRESSION TAG	UNP A9CHE9
B	228	SER	-	EXPRESSION TAG	UNP A9CHE9
B	229	HIS	-	EXPRESSION TAG	UNP A9CHE9
B	230	HIS	-	EXPRESSION TAG	UNP A9CHE9
B	231	HIS	-	EXPRESSION TAG	UNP A9CHE9
B	232	HIS	-	EXPRESSION TAG	UNP A9CHE9
B	233	HIS	-	EXPRESSION TAG	UNP A9CHE9
B	234	HIS	-	EXPRESSION TAG	UNP A9CHE9
B	235	TRP	-	EXPRESSION TAG	UNP A9CHE9
B	236	SER	-	EXPRESSION TAG	UNP A9CHE9
B	237	HIS	-	EXPRESSION TAG	UNP A9CHE9
B	238	PRO	-	EXPRESSION TAG	UNP A9CHE9
B	239	GLN	-	EXPRESSION TAG	UNP A9CHE9
B	240	PHE	-	EXPRESSION TAG	UNP A9CHE9
B	241	GLU	-	EXPRESSION TAG	UNP A9CHE9
B	242	LYS	-	EXPRESSION TAG	UNP A9CHE9

- Molecule 2 is PRASEODYMIUM ION (three-letter code: PR) (formula: Pr).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Pr 1 1	0	0
2	A	1	Total Pr 1 1	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula:  $C_8H_{10}NO_6P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	88	Total 88	O 88	0	0
5	B	76	Total 76	O 76	0	0



- Molecule 1: uncharacterized protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.69 Å 89.13 Å 109.73 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.90 47.50 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.3 (50.00-1.90) 97.9 (47.50-1.90)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.99 (at 1.91 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.219 , 0.245 0.216 , 0.242	Depositor DCC
$R_{free}$ test set	2028 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.7	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 40.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 41191 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3606	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PR, ACT, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.32	0/1762	0.61	0/2373
1	B	0.33	0/1703	0.59	0/2289
All	All	0.32	0/3465	0.60	0/4662

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1738	0	1730	22	0
1	B	1683	0	1685	14	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	B	4	0	3	1	0
4	B	15	0	7	0	0
5	A	88	0	0	2	0
5	B	76	0	0	0	0
All	All	3606	0	3425	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:165:ILE:HG23	1:A:195:MSE:HE1	1.21	1.09
1:B:67:PRO:O	1:B:71:GLU:HG3	1.88	0.72
1:A:36:LYS:HD3	1:A:57:ASN:HB2	1.72	0.71
1:A:86:SER:HB2	1:A:106:LYS:HE2	1.73	0.69
1:A:106:LYS:HD3	5:A:326:HOH:O	1.96	0.66
1:B:66:TRP:HZ3	1:B:76:ILE:HD13	1.64	0.63
1:A:117:ARG:HB3	1:A:117:ARG:NH1	2.15	0.62
1:A:100:GLU:HA	1:A:125:TYR:HB2	1.83	0.60
1:B:100:GLU:HA	1:B:125:TYR:HB2	1.87	0.56
1:A:36:LYS:HG2	1:A:37:THR:HG23	1.87	0.55
1:B:135:GLN:HG3	1:B:136:LYS:CD	2.38	0.53
1:A:165:ILE:HG23	1:A:195:MSE:CE	2.15	0.53
1:B:25:LYS:HG2	1:B:28:ASP:OD2	2.09	0.52
1:B:212:VAL:CG1	1:B:215:ALA:HB3	2.42	0.50
1:A:117:ARG:CB	1:A:117:ARG:HH11	2.26	0.49
1:B:46:VAL:HG12	1:B:51:GLN:HB3	1.95	0.49
1:A:3:GLU:O	1:A:7:ARG:HG3	2.12	0.49
1:B:44:GLN:N	1:B:45:PRO:HD2	2.28	0.49
1:A:224:LEU:HD11	5:A:330:HOH:O	2.14	0.48
1:A:117:ARG:CB	1:A:117:ARG:NH1	2.77	0.47
1:B:135:GLN:HG3	1:B:136:LYS:HD3	1.97	0.46
1:A:223:ASN:HA	1:A:226:PHE:CD2	2.51	0.45
1:B:58:ARG:HH12	1:B:60:GLN:HB2	1.81	0.44
1:B:75:ASP:N	1:B:75:ASP:OD1	2.49	0.43
1:B:135:GLN:HG3	1:B:136:LYS:HD2	2.00	0.43
1:A:176:PHE:CE1	1:A:193:MSE:HB2	2.53	0.43
1:B:195:MSE:HE3	3:B:244:ACT:H3	2.00	0.43
1:A:5:GLU:O	1:A:9:GLU:HG3	2.18	0.43
1:B:195:MSE:HA	1:B:211:ARG:O	2.19	0.42
1:A:75:ASP:N	1:A:75:ASP:OD1	2.48	0.42
1:A:224:LEU:HD12	1:A:224:LEU:C	2.40	0.42
1:A:140:ASP:HB3	1:A:143:GLU:HG3	2.03	0.41
1:A:46:VAL:HG12	1:A:51:GLN:HB3	2.03	0.41
1:A:171:ASN:HA	1:A:172:PRO:HD3	1.83	0.41
1:A:199:PHE:O	1:A:203:VAL:HG23	2.21	0.40
1:A:183:ALA:HB2	1:A:191:LEU:HD21	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	226/244 (93%)	218 (96%)	8 (4%)	0	100	100
1	B	215/244 (88%)	208 (97%)	7 (3%)	0	100	100
All	All	441/488 (90%)	426 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/192 (94%)	179 (99%)	1 (1%)	90	90
1	B	174/192 (91%)	171 (98%)	3 (2%)	68	64
All	All	354/384 (92%)	350 (99%)	4 (1%)	80	79

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115	CYS
1	B	3	GLU
1	B	85	GLN
1	B	140	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	44	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ACT	B	244	2	1,3,3	1.32	0	0,3,3	0.00	-
4	PLP	B	258	2	15,15,16	1.40	2 (13%)	21,22,23	1.52	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACT	B	244	2	-	0/0/0/0	0/0/0/0
4	PLP	B	258	2	-	0/6/6/8	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	258	PLP	C3-C2	-2.47	1.39	1.40
4	B	258	PLP	C2A-C2	2.43	1.55	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	258	PLP	O4P-P-O1P	-4.21	96.43	107.14
4	B	258	PLP	O3P-P-O2P	3.74	121.62	107.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	244	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	224/244 (91%)	0.10	6 (2%) 58 61	9, 16, 30, 40	0
1	B	218/244 (89%)	0.03	4 (1%) 71 74	10, 17, 30, 38	0
All	All	442/488 (90%)	0.06	10 (2%) 64 67	9, 17, 30, 40	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	169	GLU	3.3
1	A	229	HIS	3.2
1	A	228	SER	3.1
1	A	224	LEU	3.1
1	B	135	GLN	2.8
1	A	227	GLN	2.8
1	A	226	PHE	2.4
1	B	136	LYS	2.2
1	B	20	GLU	2.1
1	B	74	SER	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	PLP	B	258	15/16	0.94	0.12	0.52	16,18,19,21	0
2	PR	A	243	1/1	0.98	0.08	-	29,29,29,29	0
3	ACT	B	244	4/4	0.79	0.28	-	27,28,30,31	0
2	PR	B	243	1/1	0.99	0.18	-	47,47,47,47	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.