



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:14 PM GMT

PDB ID : 4RAC
Title : Aza-acyclic nucleoside phosphonates containing a second phosphonate group as inhibitors of the human, Plasmodium falciparum and vivax 6-oxopurine phosphoribosyltransferases and their pro-drugs as antimalarial agents
Authors : Keough, D.T.; Hockova, D.; Janeba, Z.; Wang, T.-H.; Naesens, L.; Edstein, M.D.; Chavchich, M.; Guddat, L.W.
Deposited on : 2014-09-10
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1192 (2.04-2.04)
Clashscore	102246	1269 (2.04-2.04)
Ramachandran outliers	100387	1258 (2.04-2.04)
Sidechain outliers	100360	1258 (2.04-2.04)
RSRZ outliers	91569	1194 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	217	 77% 16% • 6%
1	B	217	 75% 15% • 8%
1	C	217	 74% 18% • 6%
1	D	217	 77% 13% • 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	3L4	D	301[A]	-	-	-	X
2	3L4	D	301[B]	-	-	-	X

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6732 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypoxanthine-guanine phosphoribosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	204	Total	C	N	O	S	0	0	0
			1615	1039	271	298	7			
1	B	200	Total	C	N	O	S	0	0	0
			1588	1025	264	292	7			
1	C	203	Total	C	N	O	S	0	0	0
			1606	1032	269	298	7			
1	D	198	Total	C	N	O	S	0	0	0
			1554	1000	260	287	7			

- Molecule 2 is [(2-{[2-(2-AMINO-6-OXO-1,6-DIHYDRO-9H-PURIN-9-YL)ETHYL]](E)-2-PHOSPHONOETHENYL]AMINO}ETHOXY)METHYL]PHOSPHONIC ACID (three-letter code: 3L4) (formula: C₁₂H₂₀N₆O₈P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			28	12	6	8	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			28	12	6	8	2		
2	C	1	Total	C	N	O	P	0	0
			28	12	6	8	2		
2	D	1	Total	C	N	O	P	0	1
			56	24	12	16	4		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Mg	0	0
			2	2		
3	A	2	Total	Mg	0	0
			2	2		
3	D	1	Total	Mg	0	0
			1	1		
3	C	2	Total	Mg	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	75	Total	O	0	0
			75	75		
4	B	57	Total	O	0	0
			57	57		
4	C	58	Total	O	0	0
			58	58		
4	D	32	Total	O	0	0
			32	32		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hypoxanthine-guanine phosphoribosyltransferase




- Molecule 1: Hypoxanthine-guanine phosphoribosyltransferase




- Molecule 1: Hypoxanthine-guanine phosphoribosyltransferase



- Molecule 1: Hypoxanthine-guanine phosphoribosyltransferase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.82Å 114.80Å 67.66Å 90.00° 98.33° 90.00°	Depositor
Resolution (Å)	41.44 – 2.05 57.40 – 2.05	Depositor EDS
% Data completeness (in resolution range)	95.9 (41.44-2.05) 95.9 (57.40-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 2.05Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.194 , 0.244 0.200 , 0.240	Depositor DCC
R_{free} test set	2000 reflections (4.35%)	DCC
Wilson B-factor (Å ²)	31.0	Xtriage
Anisotropy	0.250	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 48000 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6732	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 3L4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/1646	0.58	0/2223
1	B	0.39	0/1617	0.58	0/2183
1	C	0.38	0/1636	0.58	0/2210
1	D	0.34	0/1581	0.56	0/2132
All	All	0.39	0/6480	0.57	0/8748

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	170	SER	Peptide
1	C	120	ASP	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1615	0	1632	18	0
1	B	1588	0	1607	16	0
1	C	1606	0	1613	21	0
1	D	1554	0	1562	19	0
2	A	28	0	17	1	0
2	B	28	0	17	2	0
2	C	28	0	18	0	0
2	D	56	0	34	5	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	1	0	0	0	0
4	A	75	0	0	2	0
4	B	57	0	0	3	0
4	C	58	0	0	1	0
4	D	32	0	0	0	0
All	All	6732	0	6500	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:301:3L4:H6	4:B:446:HOH:O	1.83	0.77
1:B:53:MET:HA	1:B:92:ILE:HD11	1.71	0.73
1:D:175:PRO:HG2	1:D:178:VAL:HG22	1.76	0.68
1:A:50:ARG:NH1	1:D:86:ARG:O	2.28	0.66
1:B:171:VAL:H	1:B:172:GLY:HA2	1.62	0.64
1:C:6:GLY:N	4:C:445:HOH:O	2.33	0.61
1:B:171:VAL:N	1:B:172:GLY:HA2	2.16	0.59
1:A:175:PRO:HG2	1:A:178:VAL:HG22	1.84	0.59
1:C:20:LEU:HD13	1:D:7:VAL:HG22	1.85	0.58
1:C:114:LYS:HE3	1:C:116:ILE:HD11	1.85	0.58
1:C:121:LEU:HG	1:C:124:LEU:HD22	1.86	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:143:GLN:HE21	1:B:143:GLN:HA	1.71	0.56
1:A:96:VAL:HG22	1:C:199:ARG:HD2	1.88	0.56
1:A:155:LYS:NZ	4:A:474:HOH:O	2.30	0.54
1:C:121:LEU:HB3	1:C:152:TYR:CE2	2.43	0.54
1:B:4:SER:N	4:B:437:HOH:O	2.40	0.54
1:A:57:GLY:HA2	1:A:92:ILE:HG12	1.89	0.53
1:B:124:LEU:HD23	1:B:149:VAL:HG13	1.90	0.53
1:A:113:ILE:N	4:A:469:HOH:O	2.40	0.53
1:B:165:LYS:HD3	1:B:184:ASP:HA	1.91	0.53
1:D:131:ILE:HB	1:D:159:VAL:HG22	1.91	0.52
1:B:120:ASP:N	4:B:449:HOH:O	2.42	0.51
1:D:141:THR:OG1	2:D:301[B]:3L4:OAE	2.29	0.51
1:C:53:MET:HA	1:C:92:ILE:HD11	1.92	0.51
1:A:175:PRO:HG2	1:A:178:VAL:CG2	2.41	0.50
1:D:146:LEU:O	1:D:150:ARG:HG3	2.12	0.49
1:D:129:VAL:HB	1:D:157:VAL:HG22	1.95	0.48
1:A:53:MET:CG	1:A:90:ARG:HD2	2.44	0.48
1:A:53:MET:HA	1:A:92:ILE:HD11	1.95	0.48
1:A:121:LEU:HA	1:A:124:LEU:HD22	1.96	0.48
1:A:100:ARG:NH1	1:C:120:ASP:OD1	2.47	0.47
1:C:207:ILE:HG23	1:C:212:LYS:HE3	1.97	0.47
1:C:101:LEU:HD23	1:C:113:ILE:HG21	1.96	0.47
1:A:23:ILE:HD11	1:A:28:ALA:HA	1.97	0.47
1:C:121:LEU:HB3	1:C:152:TYR:CZ	2.50	0.46
1:D:141:THR:OG1	2:D:301[A]:3L4:OAE	2.32	0.46
1:C:139:GLY:O	1:C:143:GLN:HG2	2.17	0.45
1:D:100:ARG:HB2	1:D:116:ILE:HB	1.97	0.45
1:D:124:LEU:HD23	1:D:149:VAL:HG13	1.98	0.45
1:B:57:GLY:HA2	1:B:92:ILE:HG12	1.99	0.45
2:A:301:3L4:H6	2:A:301:3L4:H13	1.25	0.45
1:B:67:LEU:HD22	1:B:98:PHE:HB3	1.99	0.44
1:A:138:THR:O	1:A:169:ARG:HG2	2.17	0.44
1:D:48:LEU:HA	1:D:48:LEU:HD12	1.83	0.44
1:A:56:MET:O	1:A:59:HIS:HB2	2.18	0.44
1:D:137:ASP:HB3	2:D:301[B]:3L4:OAC	2.17	0.43
1:B:41:ILE:HD11	1:B:204:VAL:HG23	2.00	0.43
1:D:138:THR:O	1:D:171:VAL:HG22	2.19	0.43
1:D:137:ASP:HB3	2:D:301[A]:3L4:OAC	2.18	0.43
1:C:174:LYS:HA	1:C:175:PRO:HD3	1.79	0.43
1:D:146:LEU:C	1:D:148:LEU:H	2.22	0.43
1:B:48:LEU:HD12	1:B:48:LEU:HA	1.86	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:142:MET:HG2	1:C:173:TYR:CE2	2.54	0.42
1:C:207:ILE:HD11	1:C:211:GLY:C	2.40	0.42
1:A:199:ARG:HD2	1:C:96:VAL:HG22	2.01	0.42
1:C:65:CYS:HB2	1:C:74:PHE:CD1	2.55	0.42
1:D:65:CYS:HB2	1:D:74:PHE:CD1	2.54	0.42
1:C:57:GLY:HA2	1:C:92:ILE:HG12	2.02	0.42
1:B:97:ASP:OD1	1:D:100:ARG:NH2	2.51	0.42
1:B:23:ILE:HG23	1:B:24:PRO:HD2	2.01	0.42
2:D:301[A]:3L4:H16	2:D:301[A]:3L4:H8	1.76	0.41
1:D:65:CYS:HB3	1:D:98:PHE:CD2	2.56	0.41
1:B:56:MET:HB3	1:B:61:ILE:HD11	2.03	0.41
2:B:301:3L4:H5	2:B:301:3L4:H13	1.80	0.41
1:C:118:GLY:HA3	1:C:121:LEU:HD22	2.03	0.41
1:A:146:LEU:O	1:A:150:ARG:HG2	2.21	0.40
1:A:134:ASP:OD1	1:A:134:ASP:N	2.52	0.40
1:D:101:LEU:HD12	1:D:101:LEU:HA	1.74	0.40
1:C:101:LEU:HG	1:C:115:VAL:HG22	2.03	0.40
1:B:101:LEU:HD12	1:B:141:THR:HG23	2.03	0.40
1:C:6:GLY:HA2	1:C:44:ARG:NH2	2.37	0.40
1:A:58:GLY:O	1:A:59:HIS:ND1	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	200/217 (92%)	195 (98%)	5 (2%)	0	100	100
1	B	192/217 (88%)	181 (94%)	11 (6%)	0	100	100
1	C	199/217 (92%)	192 (96%)	7 (4%)	0	100	100
1	D	190/217 (88%)	180 (95%)	10 (5%)	0	100	100
All	All	781/868 (90%)	748 (96%)	33 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	177/191 (93%)	163 (92%)	14 (8%)	15	7
1	B	176/191 (92%)	160 (91%)	16 (9%)	12	5
1	C	175/191 (92%)	159 (91%)	16 (9%)	12	5
1	D	169/191 (88%)	161 (95%)	8 (5%)	32	23
All	All	697/764 (91%)	643 (92%)	54 (8%)	16	7

All (54) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	ILE
1	A	40	LEU
1	A	47	ARG
1	A	48	LEU
1	A	67	LEU
1	A	71	TYR
1	A	88	SER
1	A	91	SER
1	A	97	ASP
1	A	119	ASP
1	A	124	LEU
1	A	130	LEU
1	A	148	LEU
1	A	210	THR
1	B	23	ILE
1	B	29	GLU
1	B	35	PHE
1	B	40	LEU
1	B	48	LEU
1	B	55	GLU
1	B	67	LEU
1	B	71	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	91	SER
1	B	100	ARG
1	B	130	LEU
1	B	143	GLN
1	B	148	LEU
1	B	150	ARG
1	B	210	THR
1	B	216	LYS
1	C	29	GLU
1	C	40	LEU
1	C	47	ARG
1	C	48	LEU
1	C	53	MET
1	C	59	HIS
1	C	67	LEU
1	C	71	TYR
1	C	89	ASP
1	C	112	ASP
1	C	119	ASP
1	C	121	LEU
1	C	122	SER
1	C	123	THR
1	C	130	LEU
1	C	210	THR
1	D	40	LEU
1	D	47	ARG
1	D	48	LEU
1	D	67	LEU
1	D	71	TYR
1	D	101	LEU
1	D	130	LEU
1	D	214	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 7 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	3L4	A	301	3	22,29,29	2.94	9 (40%)	25,42,42	2.75	11 (44%)
2	3L4	B	301	3	22,29,29	3.01	9 (40%)	25,42,42	2.17	8 (32%)
2	3L4	C	301	3	22,29,29	2.91	9 (40%)	25,42,42	2.38	10 (40%)
2	3L4	D	301[A]	-	22,29,29	3.09	9 (40%)	25,42,42	2.50	11 (44%)
2	3L4	D	301[B]	-	22,29,29	3.04	10 (45%)	25,42,42	2.29	12 (48%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3L4	A	301	3	-	1/12/19/19	0/2/2/2
2	3L4	B	301	3	-	0/12/19/19	0/2/2/2
2	3L4	C	301	3	-	1/12/19/19	0/2/2/2
2	3L4	D	301[A]	-	-	0/12/19/19	0/2/2/2
2	3L4	D	301[B]	-	-	0/12/19/19	0/2/2/2

All (46) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	301	3L4	CAN-N9	-3.30	1.41	1.50
2	B	301	3L4	CAN-N9	-3.23	1.41	1.50
2	A	301	3L4	CAN-N9	-3.22	1.41	1.50
2	D	301[A]	3L4	CAN-N9	-3.04	1.42	1.50
2	D	301[B]	3L4	CAN-N9	-2.99	1.42	1.50
2	D	301[A]	3L4	PBB-OAH	-2.99	1.47	1.54
2	D	301[B]	3L4	PBB-OAH	-2.98	1.47	1.54
2	A	301	3L4	CAK-NAY	-2.79	1.41	1.47
2	B	301	3L4	CAK-NAY	-2.27	1.42	1.47
2	C	301	3L4	CAK-NAY	-2.24	1.42	1.47
2	C	301	3L4	PBA-OAE	2.01	1.57	1.54
2	A	301	3L4	C4-N3	2.01	1.38	1.35
2	D	301[B]	3L4	PBA-OAC	2.02	1.52	1.48
2	A	301	3L4	PBA-OAE	2.05	1.57	1.54
2	C	301	3L4	C4-N3	2.37	1.39	1.35
2	B	301	3L4	C4-N3	2.47	1.39	1.35
2	D	301[B]	3L4	PBB-OAD	2.61	1.56	1.50
2	D	301[A]	3L4	PBB-OAD	2.70	1.56	1.50
2	A	301	3L4	CAL-NAY	2.78	1.52	1.47
2	D	301[B]	3L4	C4-N3	2.78	1.40	1.35
2	D	301[A]	3L4	C4-N3	2.79	1.40	1.35
2	C	301	3L4	CAL-NAY	2.94	1.52	1.47
2	B	301	3L4	CAL-NAY	3.15	1.52	1.47
2	D	301[A]	3L4	CAL-NAY	3.34	1.53	1.47
2	D	301[B]	3L4	CAL-NAY	3.51	1.53	1.47
2	B	301	3L4	PBA-OAC	3.74	1.55	1.48
2	B	301	3L4	C2-N2	4.76	1.43	1.34
2	D	301[B]	3L4	C2-N2	4.94	1.44	1.34
2	D	301[A]	3L4	C2-N2	5.03	1.44	1.34
2	A	301	3L4	PBB-CAP	5.06	1.85	1.79
2	A	301	3L4	C2-N2	5.07	1.44	1.34
2	D	301[B]	3L4	PBB-CAP	5.30	1.86	1.79
2	C	301	3L4	C2-N2	5.32	1.44	1.34
2	C	301	3L4	PBB-CAP	5.35	1.86	1.79
2	B	301	3L4	PBB-CAP	6.06	1.87	1.79
2	D	301[A]	3L4	PBB-CAP	6.09	1.87	1.79
2	C	301	3L4	O6-C6	6.14	1.39	1.24
2	B	301	3L4	O6-C6	6.21	1.39	1.24
2	D	301[B]	3L4	O6-C6	6.24	1.39	1.24
2	D	301[A]	3L4	O6-C6	6.27	1.39	1.24
2	C	301	3L4	CAM-NAY	6.48	1.51	1.33
2	B	301	3L4	CAM-NAY	6.54	1.51	1.33
2	A	301	3L4	O6-C6	6.60	1.40	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	3L4	CAM-NAY	6.65	1.52	1.33
2	D	301[A]	3L4	CAM-NAY	6.66	1.52	1.33
2	D	301[B]	3L4	CAM-NAY	6.81	1.52	1.33

All (52) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	3L4	CAL-NAY-CAK	-5.88	107.19	117.63
2	D	301[A]	3L4	CAL-NAY-CAK	-5.47	107.92	117.63
2	A	301	3L4	N3-C2-N1	-4.81	120.12	127.44
2	C	301	3L4	CAL-NAY-CAK	-4.74	109.21	117.63
2	A	301	3L4	C5-C6-N1	-4.71	117.14	123.59
2	A	301	3L4	CAJ-CAK-NAY	-4.62	105.33	113.75
2	C	301	3L4	N3-C2-N1	-4.58	120.47	127.44
2	B	301	3L4	C5-C6-N1	-4.28	117.73	123.59
2	D	301[A]	3L4	C5-C6-N1	-4.23	117.81	123.59
2	D	301[B]	3L4	C5-C6-N1	-4.18	117.88	123.59
2	D	301[A]	3L4	N3-C2-N1	-4.14	121.14	127.44
2	A	301	3L4	C4-C5-N7	-4.12	105.69	109.48
2	B	301	3L4	N3-C2-N1	-4.11	121.18	127.44
2	D	301[B]	3L4	N3-C2-N1	-4.03	121.30	127.44
2	C	301	3L4	C5-C6-N1	-3.85	118.32	123.59
2	D	301[B]	3L4	CAL-NAY-CAK	-3.72	111.03	117.63
2	D	301[A]	3L4	CAK-NAY-CAM	-3.70	107.69	121.41
2	C	301	3L4	CAO-CAM-NAY	-3.31	120.89	127.03
2	B	301	3L4	CAO-CAM-NAY	-3.29	120.94	127.03
2	D	301[B]	3L4	CAK-NAY-CAM	-3.17	109.65	121.41
2	C	301	3L4	CAL-NAY-CAM	-3.13	109.83	121.41
2	D	301[A]	3L4	CAL-NAY-CAM	-3.07	110.02	121.41
2	D	301[A]	3L4	CAO-CAM-NAY	-2.84	121.77	127.03
2	D	301[A]	3L4	C4-C5-N7	-2.64	107.05	109.48
2	B	301	3L4	CAL-NAY-CAK	-2.62	112.98	117.63
2	A	301	3L4	OAC-PBA-CAO	-2.61	108.27	114.41
2	D	301[B]	3L4	CAO-CAM-NAY	-2.60	122.22	127.03
2	D	301[B]	3L4	C4-C5-N7	-2.59	107.10	109.48
2	B	301	3L4	CAL-NAY-CAM	-2.56	111.91	121.41
2	C	301	3L4	OAC-PBA-CAO	-2.56	108.40	114.41
2	D	301[B]	3L4	CAL-NAY-CAM	-2.53	112.04	121.41
2	C	301	3L4	CAK-NAY-CAM	-2.38	112.59	121.41
2	C	301	3L4	C4-C5-N7	-2.32	107.34	109.48
2	A	301	3L4	CAL-NAY-CAM	-2.21	113.20	121.41
2	D	301[B]	3L4	OAC-PBA-CAO	-2.19	109.25	114.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	301[A]	3L4	OAC-PBA-CAO	-2.17	109.30	114.41
2	B	301	3L4	C4-C5-N7	-2.08	107.57	109.48
2	A	301	3L4	C6-C5-C4	-2.06	118.44	120.90
2	D	301[B]	3L4	OAG-PBB-CAP	2.02	111.36	106.64
2	D	301[B]	3L4	CAN-CAL-NAY	2.18	115.34	112.27
2	D	301[A]	3L4	OAT-CAJ-CAK	2.23	117.75	109.52
2	B	301	3L4	OAT-CAJ-CAK	2.79	119.80	109.52
2	C	301	3L4	OAT-CAJ-CAK	2.80	119.84	109.52
2	A	301	3L4	N2-C2-N1	2.81	121.85	117.20
2	A	301	3L4	OAT-CAJ-CAK	2.81	119.90	109.52
2	D	301[B]	3L4	CAJ-CAK-NAY	2.82	118.88	113.75
2	D	301[A]	3L4	CAJ-CAK-NAY	2.96	119.14	113.75
2	D	301[B]	3L4	C6-N1-C2	4.26	121.86	115.94
2	D	301[A]	3L4	C6-N1-C2	4.42	122.07	115.94
2	C	301	3L4	C6-N1-C2	4.52	122.21	115.94
2	B	301	3L4	C6-N1-C2	4.66	122.41	115.94
2	A	301	3L4	C6-N1-C2	5.35	123.37	115.94

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	301	3L4	CAO-CAM-NAY-CAL
2	A	301	3L4	CAO-CAM-NAY-CAL

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	3L4	1	0
2	B	301	3L4	2	0
2	D	301[A]	3L4	3	0
2	D	301[B]	3L4	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	204/217 (94%)	-0.12	1 (0%) 91 93	22, 38, 71, 103	0
1	B	200/217 (92%)	0.06	8 (4%) 42 48	24, 44, 78, 105	0
1	C	203/217 (93%)	0.14	11 (5%) 29 34	24, 45, 88, 118	0
1	D	198/217 (91%)	0.33	16 (8%) 15 17	29, 54, 100, 120	0
All	All	805/868 (92%)	0.10	36 (4%) 37 42	22, 44, 89, 120	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	171	VAL	7.7
1	D	172	GLY	6.8
1	D	147	SER	5.7
1	B	171	VAL	5.4
1	D	171	VAL	5.4
1	C	170	SER	4.9
1	D	123	THR	4.5
1	C	172	GLY	4.2
1	B	169	ARG	4.1
1	C	173	TYR	3.7
1	B	170	SER	3.7
1	D	121	LEU	3.6
1	D	217	ALA	3.5
1	C	167	THR	3.2
1	D	170	SER	3.1
1	B	4	SER	3.1
1	C	119	ASP	3.0
1	C	120	ASP	3.0
1	C	169	ARG	2.9
1	D	90	ARG	2.8
1	D	153	ASN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	127	LYS	2.7
1	D	143	GLN	2.5
1	D	122	SER	2.5
1	D	169	ARG	2.5
1	B	168	PRO	2.4
1	C	168	PRO	2.3
1	D	89	ASP	2.3
1	C	58	GLY	2.2
1	B	172	GLY	2.2
1	D	117	GLY	2.1
1	D	59	HIS	2.0
1	A	121	LEU	2.0
1	B	143	GLN	2.0
1	B	58	GLY	2.0
1	C	118	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	3L4	D	301[B]	28/28	0.80	0.21	2.58	33,48,84,125	28
2	3L4	D	301[A]	28/28	0.80	0.21	2.27	43,51,85,125	28
2	3L4	A	301	28/28	0.97	0.11	0.49	25,33,42,46	0
2	3L4	C	301	28/28	0.96	0.10	-0.40	31,38,49,50	0
2	3L4	B	301	28/28	0.95	0.10	-0.45	29,41,49,54	0
3	MG	C	303	1/1	0.99	0.11	-	47,47,47,47	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	MG	B	303	1/1	0.97	0.03	-	36,36,36,36	0
3	MG	D	302	1/1	0.97	0.07	-	41,41,41,41	0
3	MG	A	302	1/1	0.98	0.09	-	32,32,32,32	0
3	MG	A	303	1/1	0.97	0.07	-	30,30,30,30	0
3	MG	C	302	1/1	0.97	0.04	-	34,34,34,34	0
3	MG	B	302	1/1	0.98	0.05	-	46,46,46,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.