



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:37 AM GMT

PDB ID : 2RFE
Title : Crystal structure of the complex between the EGFR kinase domain and a Mig6 peptide
Authors : Zhang, X.; Pickin, K.A.; Bose, R.; Jura, N.; Cole, P.A.; Kuriyan, J.
Deposited on : 2007-09-28
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	<div> <div>2%</div> <div> <div></div> <div>45%</div> <div>40%</div> <div>•</div> <div>12%</div> </div> </div>
1	B	324	<div> <div>3%</div> <div> <div></div> <div>45%</div> <div>35%</div> <div>• •</div> <div>16%</div> </div> </div>
1	C	324	<div> <div>8%</div> <div> <div></div> <div>43%</div> <div>35%</div> <div>•</div> <div>19%</div> </div> </div>
1	D	324	<div> <div>7%</div> <div> <div></div> <div>49%</div> <div>35%</div> <div>•</div> <div>14%</div> </div> </div>
2	E	40	<div> <div>5%</div> <div> <div></div> <div>38%</div> <div>23%</div> <div>5%</div> <div>35%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	40	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>3%20%35%5%40%</div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8889 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Epidermal growth factor receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	286	Total	C	N	O	S	0	0	0
			2210	1426	371	398	15			
1	B	273	Total	C	N	O	S	0	0	0
			2116	1367	358	376	15			
1	C	262	Total	C	N	O	S	0	0	0
			2028	1309	344	360	15			
1	D	278	Total	C	N	O	S	0	0	0
			2130	1376	358	381	15			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	675	GLY	-	EXPRESSION TAG	UNP P00533
A	676	ALA	-	EXPRESSION TAG	UNP P00533
A	677	MET	-	EXPRESSION TAG	UNP P00533
A	799	GLU	LYS	ENGINEERED	UNP P00533
B	675	GLY	-	EXPRESSION TAG	UNP P00533
B	676	ALA	-	EXPRESSION TAG	UNP P00533
B	677	MET	-	EXPRESSION TAG	UNP P00533
B	799	GLU	LYS	ENGINEERED	UNP P00533
C	675	GLY	-	EXPRESSION TAG	UNP P00533
C	676	ALA	-	EXPRESSION TAG	UNP P00533
C	677	MET	-	EXPRESSION TAG	UNP P00533
C	799	GLU	LYS	ENGINEERED	UNP P00533
D	675	GLY	-	EXPRESSION TAG	UNP P00533
D	676	ALA	-	EXPRESSION TAG	UNP P00533
D	677	MET	-	EXPRESSION TAG	UNP P00533
D	799	GLU	LYS	ENGINEERED	UNP P00533


- Molecule 2 is a protein called ERBB receptor feedback inhibitor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	26	Total 190	C 122	N 29	O 38	S 1	0	0	0
2	F	24	Total 179	C 116	N 27	O 35	S 1	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	12	Total 12	O 12	0	0
3	B	7	Total 7	O 7	0	0
3	C	9	Total 9	O 9	0	0
3	D	7	Total 7	O 7	0	0
3	E	1	Total 1	O 1	0	0


- Molecule 1: Epidermal growth factor receptor




- Molecule 1: Epidermal growth factor receptor



- Molecule 2: ERBB receptor feedback inhibitor 1



- Molecule 2: ERBB receptor feedback inhibitor 1



LEU	SER	PRO	SER	ASN	SER	ARG	THR	PRO	SER	PRO	LYS	S337	L338	P339	S340	Y341	L342	N343	G344	V345	M346	P347	P348	T349	Q350	A353	P354	K357	Y358	V359	S360	SER	LYS	ALA	LEU
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	90.51Å 98.42Å 101.50Å 90.00° 112.59° 90.00°	Depositor
Resolution (Å)	49.85 – 2.90 49.84 – 2.91	Depositor EDS
% Data completeness (in resolution range)	82.6 (49.85-2.90) 83.4 (49.84-2.91)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.83 (at 2.91Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.229 , 0.271 0.227 , 0.266	Depositor DCC
R_{free} test set	1514 reflections (5.01%)	DCC
Wilson B-factor (Å ²)	46.2	Xtriage
Anisotropy	0.864	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 58.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 32539 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	8889	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/2258	0.67	1/3069 (0.0%)
1	B	0.43	0/2161	0.68	0/2935
1	C	0.45	0/2071	0.68	1/2812 (0.0%)
1	D	0.41	0/2175	0.64	1/2959 (0.0%)
2	E	0.53	0/197	0.82	0/271
2	F	0.49	0/186	0.67	0/256
All	All	0.44	0/9048	0.67	3/12302 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	729	PRO	N-CA-CB	5.83	110.30	103.30
1	A	729	PRO	N-CA-CB	5.71	110.15	103.30
1	D	729	PRO	N-CA-CB	5.57	109.98	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2210	0	2171	117	0
1	B	2116	0	2108	131	0
1	C	2028	0	2000	119	0
1	D	2130	0	2078	111	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	190	0	174	9	0
2	F	179	0	167	19	0
3	A	12	0	0	0	0
3	B	7	0	0	0	0
3	C	9	0	0	0	0
3	D	7	0	0	0	0
3	E	1	0	0	0	0
All	All	8889	0	8698	483	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 483 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:949:ARG:HG2	1:B:826:HIS:HB2	1.41	1.03
1:D:769:MET:HE1	1:D:822:LYS:HB2	1.45	0.98
1:A:716:ILE:H	1:A:716:ILE:HD13	1.31	0.94
1:D:716:ILE:HG22	1:D:987:VAL:O	1.69	0.93
1:D:707:TRP:HB3	1:D:716:ILE:HG13	1.54	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	280/324 (86%)	229 (82%)	45 (16%)	6 (2%)	9	32
1	B	267/324 (82%)	224 (84%)	36 (14%)	7 (3%)	7	26
1	C	254/324 (78%)	222 (87%)	27 (11%)	5 (2%)	9	33
1	D	270/324 (83%)	231 (86%)	31 (12%)	8 (3%)	5	22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	E	24/40 (60%)	17 (71%)	4 (17%)	3 (12%)	0	1
2	F	22/40 (55%)	17 (77%)	3 (14%)	2 (9%)	1	2
All	All	1117/1376 (81%)	940 (84%)	146 (13%)	31 (3%)	6	24

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	729	PRO
1	B	685	GLU
1	B	890	PRO
1	C	685	GLU
1	C	698	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/284 (81%)	215 (94%)	15 (6%)	21	52
1	B	223/284 (78%)	209 (94%)	14 (6%)	22	54
1	C	211/284 (74%)	200 (95%)	11 (5%)	29	64
1	D	220/284 (78%)	211 (96%)	9 (4%)	37	73
2	E	22/37 (60%)	21 (96%)	1 (4%)	34	70
2	F	21/37 (57%)	19 (90%)	2 (10%)	11	31
All	All	927/1210 (77%)	875 (94%)	52 (6%)	26	60

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	825	GLN
1	C	732	ASN
1	D	988	ASP
1	B	834	LEU
1	B	898	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	792	ASN
1	B	869	HIS
1	D	788	GLN
1	B	788	GLN
1	D	792	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	286/324 (88%)	0.23	7 (2%) 62 57	28, 50, 96, 104	0
1	B	273/324 (84%)	0.27	9 (3%) 50 42	30, 55, 93, 107	0
1	C	262/324 (80%)	0.47	26 (9%) 9 5	22, 56, 115, 125	0
1	D	278/324 (85%)	0.48	22 (7%) 15 10	29, 66, 103, 118	0
2	E	26/40 (65%)	0.47	2 (7%) 16 11	48, 61, 87, 90	0
2	F	24/40 (60%)	0.56	1 (4%) 40 33	66, 71, 88, 90	0
All	All	1149/1376 (83%)	0.37	67 (5%) 26 20	22, 57, 104, 125	0

The worst 5 of 67 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	723	LEU	4.7
1	B	714	VAL	4.7
1	D	699	PHE	4.3
1	C	683	LEU	4.1
1	D	683	LEU	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.