



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:38 PM GMT

PDB ID : 3RK1  
Title : 'X-ray crystal Structure of the putative N-type ATP pyrophosphatase (PF0828) in complex with ATP from Pyrococcus furiosus, Northeast Structural Genomics Consortium Target PfR23  
Authors : Forouhar, F.; Seetharaman, J.; Janjua, J.; Xiao, R.; Cunningham, K.; Ma, L.; Shastry, R.; Everett, J.K.; Nair, R.; Acton, T.B.; Rost, B.; Montelione, G.T.; Hunt, J.F.; Tong, L.; Northeast Structural Genomics Consortium (NESG)  
Deposited on : 2011-04-16  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	
1	B	237	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	A	302	-	-	-	X
3	PO4	B	302	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3755 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-type ATP pyrophosphatase superfamily.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	221	Total	C	N	O	S	Se	0	0	0
			1750	1129	285	330	1	5			
1	B	223	Total	C	N	O	S	Se	0	0	0
			1766	1141	287	332	1	5			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	230	LEU	-	EXPRESSION TAG	UNP Q8U2K6
A	231	GLU	-	EXPRESSION TAG	UNP Q8U2K6
A	232	HIS	-	EXPRESSION TAG	UNP Q8U2K6
A	233	HIS	-	EXPRESSION TAG	UNP Q8U2K6
A	234	HIS	-	EXPRESSION TAG	UNP Q8U2K6
A	235	HIS	-	EXPRESSION TAG	UNP Q8U2K6
A	236	HIS	-	EXPRESSION TAG	UNP Q8U2K6
A	237	HIS	-	EXPRESSION TAG	UNP Q8U2K6
B	230	LEU	-	EXPRESSION TAG	UNP Q8U2K6
B	231	GLU	-	EXPRESSION TAG	UNP Q8U2K6
B	232	HIS	-	EXPRESSION TAG	UNP Q8U2K6
B	233	HIS	-	EXPRESSION TAG	UNP Q8U2K6
B	234	HIS	-	EXPRESSION TAG	UNP Q8U2K6
B	235	HIS	-	EXPRESSION TAG	UNP Q8U2K6
B	236	HIS	-	EXPRESSION TAG	UNP Q8U2K6
B	237	HIS	-	EXPRESSION TAG	UNP Q8U2K6

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	94	Total	O	0	0
			94	94		
4	B	68	Total	O	0	0
			68	68		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.98 Å 64.07 Å 140.13 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.88 – 2.30 29.49 – 2.29	Depositor EDS
% Data completeness (in resolution range)	83.0 (19.88-2.30) 96.9 (29.49-2.29)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.06 (at 2.29 Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.203 , 0.253 0.223 , 0.265	Depositor DCC
$R_{free}$ test set	2343 reflections (11.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	25.5	Xtriage
Anisotropy	0.402	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 48.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 45049 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	3755	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/1774	0.58	0/2378
1	B	0.36	0/1790	0.54	0/2400
All	All	0.37	0/3564	0.56	0/4778

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1750	0	1788	42	0
1	B	1766	0	1810	47	0
2	A	31	0	12	0	0
2	B	31	0	12	3	0
3	A	10	0	0	0	0
3	B	5	0	0	0	0
4	A	94	0	0	4	0
4	B	68	0	0	2	0
All	All	3755	0	3622	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:MSE:HG2	1:A:195:PRO:HD2	1.39	1.00
1:A:132:MSE:HE2	1:A:132:MSE:HA	1.65	0.78
1:A:71:THR:HG22	1:A:73:GLY:H	1.47	0.77
1:A:92:GLN:HB3	4:A:292:HOH:O	1.91	0.71
1:B:194:MSE:HE3	1:B:196:LEU:HB2	1.72	0.71
1:A:154:GLU:HG3	1:A:157:LEU:HD12	1.73	0.70
1:A:131:TYR:CD2	1:A:132:MSE:HE3	2.27	0.70
1:B:76:GLU:O	1:B:79:VAL:HG22	1.95	0.67
1:B:4:LEU:HD12	1:B:92:GLN:HB3	1.76	0.66
1:B:194:MSE:HG3	1:B:195:PRO:HD2	1.79	0.65
1:B:48:HIS:HE1	4:B:246:HOH:O	1.80	0.64
1:B:9:VAL:HG22	1:B:95:VAL:HB	1.79	0.64
1:A:41:ASN:HB2	1:A:73:GLY:HA3	1.80	0.63
1:A:98:ALA:HB2	1:A:122:PRO:HG2	1.80	0.63
1:B:205:LYS:HB3	1:B:223:GLU:HB2	1.81	0.62
1:B:74:GLU:O	1:B:78:GLU:HG3	2.00	0.61
1:B:158:GLY:HA2	1:B:219:LEU:HD22	1.83	0.60
1:B:92:GLN:HG2	4:B:263:HOH:O	2.01	0.60
1:B:163:GLU:O	1:B:167:GLU:HG3	2.02	0.59
1:A:23:TRP:CG	1:A:122:PRO:HB3	2.38	0.59
1:B:194:MSE:HE2	1:B:197:PHE:CE2	2.38	0.58
1:B:4:LEU:HA	1:B:92:GLN:HG3	1.83	0.58
1:B:54:LEU:HD21	1:B:221:ILE:HB	1.85	0.58
1:B:102:LYS:HD2	1:B:103:TYR:N	2.19	0.57
1:A:34:LEU:HD13	1:A:64:ILE:HG21	1.84	0.57
1:A:143:MSE:HE3	1:A:192:LEU:HD11	1.88	0.55
1:B:22:TYR:OH	1:B:26:LYS:HD3	2.07	0.55
1:A:71:THR:HG23	1:A:81:ASP:OD2	2.07	0.55
1:B:166:LEU:O	1:B:170:ILE:HG12	2.07	0.54
1:B:170:ILE:O	1:B:174:GLU:HG3	2.07	0.53
1:B:104:GLN:O	1:B:108:ILE:HG12	2.09	0.52
1:A:179:HIS:CD2	1:A:186:GLU:HG3	2.45	0.52
1:A:104:GLN:O	1:A:108:ILE:HG12	2.10	0.52
1:A:98:ALA:HA	1:A:121:THR:CG2	2.41	0.51
1:B:203:VAL:HG13	1:B:221:ILE:HG23	1.93	0.50
1:B:50:ILE:O	1:B:51:ASN:ND2	2.45	0.50
1:B:81:ASP:O	1:B:85:VAL:HG23	2.12	0.49
1:A:159:ARG:HH12	1:A:168:GLU:CD	2.15	0.49
1:A:211:GLU:HG3	4:A:329:HOH:O	2.13	0.49
1:A:23:TRP:CD1	1:A:122:PRO:HB3	2.47	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:51:ASN:HD21	1:A:221:ILE:H	1.60	0.48
1:A:79:VAL:HG23	1:A:80:GLU:N	2.28	0.48
1:A:141:LYS:HB3	4:A:273:HOH:O	2.13	0.48
1:A:102:LYS:O	1:A:106:LYS:HG2	2.13	0.48
1:A:179:HIS:HD2	1:A:186:GLU:HG3	1.79	0.48
1:A:197:PHE:HB3	1:A:199:TYR:O	2.14	0.48
1:B:51:ASN:C	1:B:51:ASN:HD22	2.18	0.47
1:B:110:LYS:O	1:B:114:GLU:HG3	2.14	0.47
1:B:127:ASP:HB3	1:B:130:GLU:HG2	1.95	0.47
1:B:102:LYS:C	1:B:102:LYS:HD2	2.35	0.47
1:B:22:TYR:CZ	1:B:26:LYS:HD3	2.50	0.47
1:B:222:GLU:O	1:B:223:GLU:HG3	2.14	0.47
1:A:175:LYS:HE2	1:A:176:TYR:CZ	2.50	0.47
1:B:182:GLY:HA2	1:B:186:GLU:HG2	1.97	0.46
1:A:107:ARG:O	1:A:111:VAL:HG23	2.15	0.46
1:B:194:MSE:HG2	1:B:196:LEU:H	1.81	0.46
1:A:69:GLY:HA3	1:A:85:VAL:HG21	1.98	0.46
1:B:194:MSE:HE2	1:B:197:PHE:HE2	1.81	0.46
1:B:202:VAL:HG21	1:B:227:GLU:OE2	2.16	0.45
1:A:161:LEU:HD21	1:A:169:LEU:HD11	1.98	0.45
1:B:4:LEU:HA	1:B:92:GLN:CG	2.46	0.45
1:B:74:GLU:O	1:B:75:LYS:C	2.54	0.45
1:A:141:LYS:CG	1:A:193:ASP:HB3	2.46	0.45
1:A:119:VAL:O	1:A:119:VAL:HG13	2.16	0.44
1:B:128:ALA:HB1	1:B:180:VAL:HB	1.99	0.44
1:A:132:MSE:HE1	1:A:135:LEU:HD12	2.00	0.44
1:A:24:ALA:O	1:A:29:PHE:HB2	2.18	0.44
1:B:197:PHE:HB3	1:B:199:TYR:O	2.17	0.44
1:A:75:LYS:HA	1:A:78:GLU:HG3	2.00	0.44
1:B:24:ALA:O	1:B:29:PHE:HB2	2.18	0.44
1:B:47:TYR:CE1	1:B:78:GLU:HB3	2.54	0.43
1:B:10:LEU:HB3	2:B:301:ATP:N3	2.34	0.43
1:A:53:ASN:HB2	4:A:261:HOH:O	2.17	0.43
1:B:186:GLU:HG2	1:B:186:GLU:H	1.57	0.43
1:A:15:LYS:HG3	1:A:16:ASP:N	2.34	0.43
1:B:179:HIS:HB3	1:B:183:GLU:OE2	2.19	0.43
1:A:131:TYR:CE2	1:A:132:MSE:HE3	2.54	0.42
1:B:4:LEU:HA	1:B:92:GLN:CB	2.49	0.42
1:B:154:GLU:HG3	1:B:157:LEU:HD12	2.02	0.42
1:A:200:LYS:HE3	1:A:229:LYS:HB3	2.02	0.42
1:A:107:ARG:HH21	1:A:107:ARG:HG2	1.85	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23:TRP:CE2	1:B:122:PRO:HB3	2.55	0.42
1:B:12:SER:HB3	2:B:301:ATP:C5	2.55	0.42
1:A:158:GLY:HA2	1:A:219:LEU:HD22	2.01	0.42
1:A:176:TYR:O	1:A:177:LYS:HB2	2.20	0.41
1:A:129:LYS:O	1:A:133:ARG:HG3	2.20	0.41
1:B:97:GLY:HA3	2:B:301:ATP:O3'	2.20	0.41
1:B:157:LEU:HD13	1:B:208:LYS:HB2	2.03	0.40
1:A:46:MSE:HE3	1:A:47:TYR:CZ	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/237 (92%)	212 (98%)	5 (2%)	0	100	100
1	B	219/237 (92%)	207 (94%)	10 (5%)	2 (1%)	21	24
All	All	436/474 (92%)	419 (96%)	15 (3%)	2 (0%)	34	41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	75	LYS
1	B	100	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/194 (96%)	181 (97%)	6 (3%)	46	62
1	B	189/194 (97%)	180 (95%)	9 (5%)	31	42
All	All	376/388 (97%)	361 (96%)	15 (4%)	38	52

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	LYS
1	A	32	LYS
1	A	41	ASN
1	A	159	ARG
1	A	166	LEU
1	A	211	GLU
1	B	34	LEU
1	B	48	HIS
1	B	51	ASN
1	B	72	GLN
1	B	102	LYS
1	B	121	THR
1	B	162	ASP
1	B	186	GLU
1	B	215	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	ASN
1	A	51	ASN
1	A	137	ASN
1	A	179	HIS
1	B	48	HIS
1	B	51	ASN
1	B	53	ASN
1	B	92	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ATP	A	301	-	24,33,33	1.48	3 (12%)	31,52,52	2.49	5 (16%)
3	PO4	A	302	-	4,4,4	1.04	0	6,6,6	0.27	0
3	PO4	A	303	-	4,4,4	1.01	0	6,6,6	0.27	0
2	ATP	B	301	-	24,33,33	1.48	3 (12%)	31,52,52	2.48	5 (16%)
3	PO4	B	302	-	4,4,4	1.11	0	6,6,6	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	A	301	-	-	0/18/38/38	0/3/3/3
3	PO4	A	302	-	-	0/0/0/0	0/0/0/0
3	PO4	A	303	-	-	0/0/0/0	0/0/0/0
2	ATP	B	301	-	-	0/18/38/38	0/3/3/3
3	PO4	B	302	-	-	0/0/0/0	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	ATP	PG-O2G	-2.64	1.45	1.54
2	A	301	ATP	PG-O2G	-2.62	1.45	1.54
2	B	301	ATP	C2-N1	2.32	1.38	1.33
2	A	301	ATP	C2-N1	2.34	1.38	1.33
2	A	301	ATP	C2-N3	4.16	1.39	1.32
2	B	301	ATP	C2-N3	4.18	1.39	1.32

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	ATP	N3-C2-N1	-11.40	120.17	128.89
2	B	301	ATP	N3-C2-N1	-11.35	120.21	128.89
2	A	301	ATP	C4-C5-N7	-4.86	105.00	109.48
2	B	301	ATP	C4-C5-N7	-4.84	105.03	109.48
2	A	301	ATP	PB-O3B-PG	-2.35	124.79	132.67
2	B	301	ATP	PB-O3B-PG	-2.34	124.81	132.67
2	A	301	ATP	N6-C6-N1	-2.27	114.32	119.20
2	B	301	ATP	N6-C6-N1	-2.24	114.39	119.20
2	A	301	ATP	C2'-C1'-N9	2.29	117.80	114.29
2	B	301	ATP	C2'-C1'-N9	2.30	117.81	114.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	ATP	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	216/237 (91%)	0.06	2 (0%) 85 89	9, 23, 45, 57	0
1	B	218/237 (91%)	0.38	14 (6%) 23 31	10, 30, 47, 56	0
All	All	434/474 (91%)	0.22	16 (3%) 45 54	9, 26, 47, 57	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	230	LEU	3.3
1	B	102	LYS	3.2
1	B	212	PRO	2.9
1	B	106	LYS	2.9
1	B	48	HIS	2.8
1	B	45	TYR	2.7
1	B	75	LYS	2.6
1	B	4	LEU	2.4
1	B	98	ALA	2.3
1	B	50	ILE	2.2
1	A	100	ALA	2.1
1	B	108	ILE	2.1
1	A	76	GLU	2.1
1	B	105	ARG	2.1
1	B	209	VAL	2.1
1	B	223	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PO4	B	302	5/5	0.94	0.54	7.08	32,32,32,32	5
3	PO4	A	302	5/5	0.97	0.22	4.06	25,25,26,26	5
2	ATP	B	301	31/31	0.92	0.21	0.90	20,25,32,34	0
3	PO4	A	303	5/5	0.96	0.17	0.49	37,37,39,40	0
2	ATP	A	301	31/31	0.95	0.13	-1.17	15,24,36,38	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.