



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 10:01 PM GMT

PDB ID : 1RQH
Title : Propionibacterium shermanii transcarboxylase 5S subunit bound to pyruvic acid
Authors : Hall, P.R.; Zheng, R.; Antony, L.; Pusztai-Carey, M.; Carey, P.R.; Yee, V.C.
Deposited on : 2003-12-05
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	540	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PYR	A	602	-	-	X	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4098 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called transcarboxylase 5S subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	471	3676	2308	637	702	29	0	3	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	CLONING ARTIFACT	GB 38304072
A	-9	ALA	-	CLONING ARTIFACT	GB 38304072
A	-8	ILE	-	CLONING ARTIFACT	GB 38304072
A	-7	SER	-	CLONING ARTIFACT	GB 38304072
A	-6	ARG	-	CLONING ARTIFACT	GB 38304072
A	-5	GLU	-	CLONING ARTIFACT	GB 38304072
A	-4	LEU	-	CLONING ARTIFACT	GB 38304072
A	-3	VAL	-	CLONING ARTIFACT	GB 38304072
A	-2	ASP	-	CLONING ARTIFACT	GB 38304072
A	-1	PRO	-	CLONING ARTIFACT	GB 38304072
A	0	ASN	-	CLONING ARTIFACT	GB 38304072
A	1	SER	-	CLONING ARTIFACT	GB 38304072
A	184	KCX	LYS	MICROHETEROGENEITY	GB 38304072
A	506	THR	-	CLONING ARTIFACT	GB 38304072
A	507	ARG	-	CLONING ARTIFACT	GB 38304072
A	508	ALA	-	CLONING ARTIFACT	GB 38304072
A	509	SER	-	CLONING ARTIFACT	GB 38304072
A	510	GLN	-	CLONING ARTIFACT	GB 38304072
A	511	PRO	-	CLONING ARTIFACT	GB 38304072
A	512	GLU	-	CLONING ARTIFACT	GB 38304072
A	513	LEU	-	CLONING ARTIFACT	GB 38304072
A	514	ALA	-	CLONING ARTIFACT	GB 38304072
A	515	PRO	-	CLONING ARTIFACT	GB 38304072
A	516	GLU	-	CLONING ARTIFACT	GB 38304072
A	517	ASP	-	CLONING ARTIFACT	GB 38304072
A	518	PRO	-	CLONING ARTIFACT	GB 38304072
A	519	GLU	-	CLONING ARTIFACT	GB 38304072

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	520	ASP	-	CLONING ARTIFACT	GB 38304072
A	521	LEU	-	CLONING ARTIFACT	GB 38304072
A	522	GLU	-	CLONING ARTIFACT	GB 38304072
A	523	HIS	-	CLONING ARTIFACT	GB 38304072
A	524	HIS	-	CLONING ARTIFACT	GB 38304072
A	525	HIS	-	CLONING ARTIFACT	GB 38304072
A	526	HIS	-	CLONING ARTIFACT	GB 38304072
A	527	HIS	-	CLONING ARTIFACT	GB 38304072
A	528	HIS	-	CLONING ARTIFACT	GB 38304072

- Molecule 2 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Co	0	0
			1	1		

- Molecule 3 is PYRUVIC ACID (three-letter code: PYR) (formula: C₃H₄O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	415	Total 415	O 415	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: transcarboxylase 5S subunit



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	96.03Å 145.72Å 78.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.09 – 2.00 40.99 – 1.90	Depositor EDS
% Data completeness (in resolution range)	92.6 (28.09-2.00) 92.7 (40.99-1.90)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.00 (at 1.91Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.164 , 0.197 0.165 , 0.196	Depositor DCC
R_{free} test set	1657 reflections (4.75%)	DCC
Wilson B-factor (Å ²)	23.5	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 55.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 43392 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4098	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PYR, CO, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.28	41/3744 (1.1%)	0.65	2/5058 (0.0%)

All (41) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	184[B]	LYS	CG-CD	12.08	1.93	1.52
1	A	184[B]	LYS	CD-CE	-9.88	1.26	1.51
1	A	184[B]	LYS	CB-CG	9.71	1.78	1.52
1	A	184[B]	LYS	CA-CB	-7.84	1.36	1.53
1	A	184[B]	LYS	CE-NZ	-7.84	1.29	1.49
1	A	5	GLU	CB-CG	-6.73	1.39	1.52
1	A	420	GLU	CB-CG	-6.72	1.39	1.52
1	A	13	GLU	CB-CG	-6.72	1.39	1.52
1	A	367	GLU	CB-CG	-6.71	1.39	1.52
1	A	399	GLU	CB-CG	-6.71	1.39	1.52
1	A	70	GLU	CB-CG	-6.70	1.39	1.52
1	A	164	GLU	CB-CG	-6.70	1.39	1.52
1	A	334	GLU	CB-CG	-6.69	1.39	1.52
1	A	374	GLU	CB-CG	-6.69	1.39	1.52
1	A	325	GLU	CB-CG	-6.68	1.39	1.52
1	A	316	GLU	CB-CG	-6.67	1.39	1.52
1	A	473	GLU	CB-CG	-6.67	1.39	1.52
1	A	106	GLU	CB-CG	-6.66	1.39	1.52
1	A	232	GLU	CB-CG	-6.66	1.39	1.52
1	A	255	GLU	CB-CG	-6.66	1.39	1.52
1	A	36	GLU	CB-CG	-6.66	1.39	1.52
1	A	400	GLU	CB-CG	-6.66	1.39	1.52
1	A	117	GLU	CB-CG	-6.66	1.39	1.52
1	A	262	GLU	CB-CG	-6.66	1.39	1.52
1	A	330	GLU	CB-CG	-6.65	1.39	1.52
1	A	425	GLU	CB-CG	-6.65	1.39	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	293	GLU	CB-CG	-6.65	1.39	1.52
1	A	54	GLU	CB-CG	-6.65	1.39	1.52
1	A	259	GLU	CB-CG	-6.64	1.39	1.52
1	A	7	GLU	CB-CG	-6.64	1.39	1.52
1	A	454	GLU	CB-CG	-6.64	1.39	1.52
1	A	18	GLU	CB-CG	-6.64	1.39	1.52
1	A	437	GLU	CB-CG	-6.64	1.39	1.52
1	A	10	GLU	CB-CG	-6.63	1.39	1.52
1	A	74	GLU	CB-CG	-6.62	1.39	1.52
1	A	224	GLU	CB-CG	-6.61	1.39	1.52
1	A	418	GLU	CB-CG	-6.60	1.39	1.52
1	A	458	GLU	CB-CG	-6.60	1.39	1.52
1	A	184[B]	LYS	CA-C	6.14	1.69	1.52
1	A	370	ARG	CG-CD	-5.02	1.39	1.51
1	A	320	ARG	CG-CD	-5.02	1.39	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	184[B]	LYS	CD-CE-NZ	10.38	135.56	111.70
1	A	184[B]	LYS	CB-CG-CD	-9.35	87.28	111.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3676	0	3639	66	0
2	A	1	0	0	0	0
3	A	6	0	3	5	0
4	A	415	0	0	5	2
All	All	4098	0	3642	66	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:353:GLN:H	1:A:353:GLN:HE21	0.94	0.93
1:A:290:LYS:HE3	4:A:963:HOH:O	1.71	0.90
1:A:327:LYS:N	1:A:327:LYS:HD2	1.90	0.87
1:A:186[A]:MET:SD	4:A:945:HOH:O	2.32	0.86
1:A:186[B]:MET:HE3	3:A:602:PYR:H31	1.57	0.85
1:A:353:GLN:HE21	1:A:353:GLN:N	1.75	0.84
1:A:324:ALA:HB1	1:A:327:LYS:HD3	1.61	0.82
1:A:353:GLN:H	1:A:353:GLN:NE2	1.78	0.79
1:A:378:ILE:HG12	1:A:383:TYR:CD2	2.21	0.76
1:A:432:PHE:CE1	1:A:434:GLY:HA2	2.22	0.74
1:A:335:VAL:HB	1:A:336:PRO:HD3	1.74	0.70
1:A:391:ASP:OD2	1:A:394:VAL:HG23	1.96	0.64
1:A:349:THR:HA	1:A:353:GLN:HE22	1.63	0.64
1:A:458:GLU:OE2	4:A:1016:HOH:O	2.14	0.64
1:A:153:ILE:HG12	1:A:169:LEU:HD13	1.82	0.62
1:A:378:ILE:HG12	1:A:383:TYR:HD2	1.66	0.61
1:A:186[B]:MET:HE3	3:A:602:PYR:C3	2.30	0.61
1:A:327:LYS:N	1:A:327:LYS:CD	2.63	0.60
1:A:186[B]:MET:CE	3:A:602:PYR:C3	2.80	0.59
1:A:186[B]:MET:CE	3:A:602:PYR:H31	2.30	0.58
1:A:139:MET:HE1	1:A:149:ALA:HB1	1.86	0.58
1:A:305:SER:O	1:A:306:GLN:HB2	2.03	0.58
1:A:193:GLN:HB3	1:A:194:PRO:HD3	1.85	0.58
1:A:319:LEU:HD11	1:A:331:VAL:HG21	1.87	0.57
1:A:37:ASP:HB3	1:A:285:ILE:HD13	1.88	0.56
1:A:104:ASN:OD1	1:A:106:GLU:HG2	2.06	0.56
1:A:354:ILE:HD12	1:A:378:ILE:HD11	1.87	0.56
1:A:22:ARG:C	1:A:22:ARG:HD2	2.27	0.54
1:A:354:ILE:CD1	1:A:378:ILE:HD11	2.37	0.54
1:A:362:ASN:HA	1:A:366:GLY:O	2.08	0.53
1:A:365:MET:SD	1:A:370:ARG:HD3	2.49	0.53
1:A:401:GLN:O	1:A:401:GLN:HG2	2.08	0.52
1:A:18:GLU:HB2	1:A:50:TYR:CE2	2.45	0.51
1:A:314:ASN:O	1:A:318:GLN:HG3	2.11	0.49
1:A:391:ASP:O	1:A:395:VAL:HG23	2.12	0.48
1:A:130:ASN:O	1:A:132:PRO:HD3	2.13	0.47
1:A:446:PRO:HG3	4:A:805:HOH:O	2.14	0.47
1:A:467:ASP:O	1:A:471:LYS:HG3	2.15	0.46
1:A:139:MET:CE	1:A:149:ALA:HB1	2.44	0.46
1:A:371:MET:HE2	1:A:375:PHE:HD2	1.80	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:ASP:CB	1:A:285:ILE:HD13	2.45	0.46
1:A:229:LYS:HA	1:A:229:LYS:HD3	1.69	0.45
1:A:326:ASP:C	1:A:327:LYS:HD2	2.36	0.45
1:A:114:LYS:HE3	1:A:436:ASP:HB2	1.99	0.45
1:A:427:ALA:HA	1:A:432:PHE:CD1	2.52	0.45
1:A:374:GLU:CD	1:A:374:GLU:H	2.20	0.45
1:A:360:VAL:O	1:A:364:MET:HG2	2.16	0.45
1:A:213:ASN:HD22	1:A:213:ASN:N	2.14	0.45
1:A:418:GLU:O	1:A:422:GLN:HG2	2.18	0.44
1:A:37:ASP:HB3	1:A:285:ILE:HG21	1.99	0.44
1:A:18:GLU:HB2	1:A:50:TYR:CZ	2.53	0.43
1:A:186[B]:MET:HE1	3:A:602:PYR:H33	2.01	0.43
1:A:421:LYS:O	1:A:425:GLU:HG3	2.19	0.43
1:A:313:SER:O	1:A:316:GLU:HB3	2.18	0.42
1:A:374:GLU:N	1:A:374:GLU:CD	2.73	0.42
1:A:405:LYS:HE2	1:A:405:LYS:HB3	1.82	0.42
1:A:81:LYS:HB3	1:A:81:LYS:HE2	1.79	0.42
1:A:371:MET:HB3	1:A:371:MET:HE2	1.77	0.41
1:A:114:LYS:HD3	1:A:114:LYS:HA	1.76	0.41
1:A:129:MET:CE	1:A:345:PRO:HB3	2.50	0.41
1:A:19:LEU:HD11	1:A:239:ASP:HB3	2.01	0.41
1:A:466:THR:OG1	1:A:469:GLN:HG3	2.20	0.41
1:A:409:GLN:HG3	1:A:410:ARG:N	2.35	0.41
1:A:198:ILE:O	1:A:202:ILE:HG13	2.21	0.41
1:A:129:MET:HE3	1:A:345:PRO:HB3	2.02	0.40
1:A:405:LYS:NZ	4:A:830:HOH:O	2.54	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:712:HOH:O	4:A:712:HOH:O[4_555]	1.71	0.49
4:A:712:HOH:O	4:A:834:HOH:O[4_555]	2.14	0.06

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	470/540 (87%)	455 (97%)	13 (3%)	2 (0%)	39	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	364	MET
1	A	367	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	392/448 (88%)	382 (97%)	10 (3%)	54	54

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	GLU
1	A	29	MET
1	A	31	THR
1	A	56	TRP
1	A	169	LEU
1	A	213	ASN
1	A	294	SER
1	A	322	GLN
1	A	326	ASP
1	A	353	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	208	GLN
1	A	211	GLN
1	A	213	ASN
1	A	353	GLN
1	A	447	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	KCX	A	184[A]	1,2	7,11,12	1.97	2 (28%)	7,12,14	2.06	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	184[A]	1,2	-	0/6/10/12	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	184[A]	KCX	CB-CA	-2.02	1.51	1.53
1	A	184[A]	KCX	CA-N	4.50	1.62	1.47

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	184[A]	KCX	CB-CA-N	-4.42	97.96	110.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PYR	A	602	-	2,5,5	2.03	1 (50%)	2,6,6	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PYR	A	602	-	-	0/0/4/4	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	PYR	C3-C2	-2.86	1.40	1.49

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	PYR	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	470/540 (87%)	-0.33	23 (4%) 33 35	13, 25, 71, 105	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	324	ALA	6.5
1	A	370	ARG	5.3
1	A	323	GLY	5.0
1	A	326	ASP	4.6
1	A	327	LYS	4.6
1	A	363	VAL	4.5
1	A	320	ARG	4.2
1	A	322	GLN	3.9
1	A	364	MET	3.6
1	A	369	LYS	3.6
1	A	365	MET	3.4
1	A	474	ALA	3.4
1	A	321	ALA	3.2
1	A	319	LEU	2.9
1	A	424	LYS	2.5
1	A	360	VAL	2.4
1	A	316	GLU	2.4
1	A	472	ALA	2.4
1	A	331	VAL	2.3
1	A	368	TYR	2.3
1	A	325	GLU	2.2
1	A	471	LYS	2.2
1	A	314	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	KCX	A	184[A]	12/13	0.97	0.12	-	10,20,21,21	12

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	PYR	A	602	6/6	0.89	0.28	8.92	16,18,23,24	6
2	CO	A	601	1/1	1.00	0.04	-3.55	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.