



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:23 PM GMT

PDB ID : 4RR0  
Title : re-refined 1vcw, CRYSTAL STRUCTURE OF DEGS AFTER BACKSOAK-  
ING THE ACTIVATING PEPTIDE  
Authors : Sauer, R.T.; Grant, R.A.  
Deposited on : 2014-11-05  
Resolution : 3.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1191 (3.10-3.02)
Clashscore	102246	1303 (3.10-3.02)
Ramachandran outliers	100387	1254 (3.10-3.02)
Sidechain outliers	100360	1254 (3.10-3.02)
RSRZ outliers	91569	1197 (3.10-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	<div> <div>4%</div> <div>84%</div> <div>11%</div> </div>
1	B	314	<div> <div>6%</div> <div>83%</div> <div>15%</div> </div>
1	C	314	<div> <div>6%</div> <div>83%</div> <div>14%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 12160 atoms, of which 6139 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protease degS.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	279	Total	C	H	N	O	S	0	0	0
			4171	1295	2106	366	399	5			
1	B	268	Total	C	H	N	O	S	0	0	0
			3970	1237	2008	345	375	5			
1	C	271	Total	C	H	N	O	S	0	0	0
			4016	1249	2025	353	384	5			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	42	MET	-	EXPRESSION TAG	UNP H9UXC8
B	42	MET	-	EXPRESSION TAG	UNP H9UXC8
C	42	MET	-	EXPRESSION TAG	UNP H9UXC8


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	O	0	0
			2	2		
2	C	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Protease degS



#### • Molecule 1: Protease degS



#### • Molecule 1: Protease degS



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	206.99Å 142.73Å 41.23Å 90.00° 90.09° 90.00°	Depositor
Resolution (Å)	19.90 – 3.05 19.90 – 3.05	Depositor EDS
% Data completeness (in resolution range)	96.4 (19.90-3.05) 96.4 (19.90-3.05)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.04 (at 3.04Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1760)	Depositor
R, $R_{free}$	0.226 , 0.256 0.226 , 0.256	Depositor DCC
$R_{free}$ test set	1104 reflections (5.05%)	DCC
Wilson B-factor (Å <sup>2</sup> )	72.1	Xtriage
Anisotropy	0.282	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 70.1	EDS
Estimated twinning fraction	0.048 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 21871 reflections	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	12160	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	102.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/2090	0.47	0/2849
1	B	0.25	0/1984	0.47	0/2701
1	C	0.25	0/2014	0.47	0/2743
All	All	0.26	0/6088	0.47	0/8293

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2065	2106	2104	7	0
1	B	1962	2008	2004	4	0
1	C	1991	2025	2023	5	0
2	A	2	0	0	0	0
2	C	1	0	0	0	0
All	All	6021	6139	6131	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:48:ASN:OD1	1:B:52:ARG:NH1	2.45	0.49
1:C:94:ASN:OD1	1:C:202:GLY:N	2.47	0.48
1:C:85:MET:HG3	1:C:245:MET:SD	2.55	0.47
1:B:326:ARG:HG3	1:B:327:PRO:HD2	1.98	0.45
1:A:145:ALA:O	1:A:239:GLN:NE2	2.50	0.44
1:A:65:GLY:HA2	1:A:102:ASP:HB2	2.00	0.43
1:C:48:ASN:OD1	1:C:52:ARG:NH1	2.52	0.43
1:C:47:TYR:HB3	1:C:156:LEU:HD11	2.00	0.42
1:A:184:THR:HG21	1:A:188:ASN:ND2	2.34	0.42
1:A:47:TYR:HB3	1:A:156:LEU:HD11	2.02	0.42
1:B:261:ILE:HG22	1:B:262:GLY:N	2.34	0.41
1:A:69:ASN:OD1	1:A:72:ASN:N	2.53	0.41
1:B:184:THR:HG21	1:B:188:ASN:ND2	2.35	0.41
1:A:259:ILE:HG23	1:A:260:GLY:N	2.36	0.41
1:A:55:ALA:HB3	1:A:56:PRO:HD3	2.03	0.40
1:C:184:THR:HG21	1:C:188:ASN:ND2	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/314 (86%)	268 (99%)	3 (1%)	0	100	100
1	B	256/314 (82%)	252 (98%)	4 (2%)	0	100	100
1	C	261/314 (83%)	255 (98%)	6 (2%)	0	100	100
All	All	788/942 (84%)	775 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/253 (89%)	225 (100%)	0	100	100
1	B	210/253 (83%)	210 (100%)	0	100	100
1	C	214/253 (85%)	214 (100%)	0	100	100
All	All	649/759 (86%)	649 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.



## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	279/314 (88%)	0.07	14 (5%)	32 13	29, 62, 177, 199	0
1	B	268/314 (85%)	0.28	20 (7%)	17 6	31, 90, 179, 201	0
1	C	271/314 (86%)	0.25	18 (6%)	22 7	32, 80, 175, 193	0
All	All	818/942 (86%)	0.20	52 (6%)	23 8	29, 80, 178, 201	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	305	ILE	6.3
1	A	301	ASN	6.1
1	B	301	ASN	4.9
1	B	303	LEU	4.8
1	A	303	LEU	4.7
1	C	301	ASN	4.2
1	C	303	LEU	4.1
1	C	185	GLY	3.8
1	B	305	ILE	3.7
1	A	345	GLN	3.5
1	B	264	ARG	3.3
1	A	282	ILE	3.3
1	B	288	SER	3.3
1	C	219	SER	3.2
1	C	68	THR	3.1
1	A	283	VAL	3.1
1	C	313	ILE	3.0
1	B	280	GLN	3.0
1	A	341	GLN	3.0
1	C	69	ASN	2.9
1	A	297	GLY	2.8
1	B	290	ASP	2.8
1	A	290	ASP	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	77	ARG	2.7
1	C	314	SER	2.7
1	A	227	GLU	2.6
1	A	313	ILE	2.6
1	B	258	TYR	2.6
1	B	263	GLY	2.6
1	B	344	LEU	2.6
1	A	219	SER	2.6
1	C	333	VAL	2.5
1	B	185	GLY	2.5
1	C	73	GLN	2.5
1	C	183	PRO	2.5
1	C	330	VAL	2.4
1	B	316	LEU	2.4
1	B	286	GLU	2.4
1	C	283	VAL	2.4
1	B	180	GLY	2.3
1	B	345	GLN	2.3
1	C	309	ASN	2.3
1	B	75	GLU	2.2
1	B	262	GLY	2.2
1	A	287	VAL	2.2
1	A	352	PRO	2.2
1	B	352	PRO	2.2
1	B	314	SER	2.1
1	C	82	GLY	2.1
1	A	335	VAL	2.1
1	C	72	ASN	2.1
1	C	182	ASN	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.