



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:23 PM GMT

PDB ID : 4RR1
Title : re-refinement of entry 1sot, Crystal Structure of the DegS stress sensor
Authors : Sauer, R.T.; Grant, R.A.
Deposited on : 2014-11-05
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	<div> <div>13%</div> <div>88%</div> <div>12%</div> </div>
1	B	320	<div> <div>12%</div> <div>85%</div> <div>13%</div> </div>
1	C	320	<div> <div>15%</div> <div>81%</div> <div>16%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NI	B	401	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12732 atoms, of which 6216 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protease degS.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	282	Total	C	H	N	O	Se	0	3	0
			4225	1308	2145	373	393	6			
1	B	277	Total	C	H	N	O	Se	0	1	0
			4116	1281	2075	369	385	6			
1	C	268	Total	C	H	N	O	Se	0	2	0
			3938	1230	1996	342	365	5			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	42	MSE	-	EXPRESSION TAG	UNP H9UXC8
A	356	HIS	-	EXPRESSION TAG	UNP H9UXC8
A	357	HIS	-	EXPRESSION TAG	UNP H9UXC8
A	358	HIS	-	EXPRESSION TAG	UNP H9UXC8
A	359	HIS	-	EXPRESSION TAG	UNP H9UXC8
A	360	HIS	-	EXPRESSION TAG	UNP H9UXC8
A	361	HIS	-	EXPRESSION TAG	UNP H9UXC8
B	42	MSE	-	EXPRESSION TAG	UNP H9UXC8
B	356	HIS	-	EXPRESSION TAG	UNP H9UXC8
B	357	HIS	-	EXPRESSION TAG	UNP H9UXC8
B	358	HIS	-	EXPRESSION TAG	UNP H9UXC8
B	359	HIS	-	EXPRESSION TAG	UNP H9UXC8
B	360	HIS	-	EXPRESSION TAG	UNP H9UXC8
B	361	HIS	-	EXPRESSION TAG	UNP H9UXC8
C	42	MSE	-	EXPRESSION TAG	UNP H9UXC8
C	356	HIS	-	EXPRESSION TAG	UNP H9UXC8
C	357	HIS	-	EXPRESSION TAG	UNP H9UXC8
C	358	HIS	-	EXPRESSION TAG	UNP H9UXC8
C	359	HIS	-	EXPRESSION TAG	UNP H9UXC8
C	360	HIS	-	EXPRESSION TAG	UNP H9UXC8
C	361	HIS	-	EXPRESSION TAG	UNP H9UXC8

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Ni	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	178	Total	O	0	0
			178	178		
4	B	144	Total	O	0	0
			144	144		
4	C	125	Total	O	0	0
			125	125		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Protease degS



• Molecule 1: Protease degS



• Molecule 1: Protease degS



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	207.68Å 143.06Å 41.53Å 90.00° 90.07° 90.00°	Depositor
Resolution (Å)	15.00 – 2.30 15.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	96.4 (15.00-2.30) 93.7 (15.00-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.03 (at 1.95Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.184 , 0.217 0.184 , 0.217	Depositor DCC
R_{free} test set	2616 reflections (5.07%)	DCC
Wilson B-factor (Å ²)	37.9	Xtriage
Anisotropy	0.257	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 65.8	EDS
Estimated twinning fraction	0.032 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	3 of 82290 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12732	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NI, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/2105	0.49	0/2853
1	B	0.28	0/2063	0.47	0/2794
1	C	0.27	0/1964	0.46	0/2666
All	All	0.28	0/6132	0.48	0/8313

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2080	2145	2145	1	0
1	B	2041	2075	2075	3	0
1	C	1942	1996	1996	4	0
2	A	5	0	0	0	0
3	B	1	0	0	0	0
4	A	178	0	0	1	0
4	B	144	0	0	0	0
4	C	125	0	0	0	0
All	All	6516	6216	6216	8	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 1.

All (8) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:GLY:O	4:A:553:HOH:O	2.14	0.65
1:B:85:MSE:HG3	1:B:245:MSE:SE	2.54	0.57
1:C:47:TYR:HB3	1:C:156:LEU:HD11	1.99	0.43
1:C:159:GLY:HA3	1:C:200:ASN:HA	2.01	0.43
1:B:159:GLY:HA3	1:B:200:ASN:HA	2.01	0.42
1:B:258:TYR:CZ	1:B:260:GLY:HA2	2.54	0.42
1:C:85:MSE:HG3	1:C:245:MSE:SE	2.71	0.41
1:C:94:ASN:OD1	1:C:202:GLY:N	2.53	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/320 (86%)	274 (100%)	1 (0%)	0	100	100
1	B	266/320 (83%)	266 (100%)	0	0	100	100
1	C	258/320 (81%)	257 (100%)	1 (0%)	0	100	100
All	All	799/960 (83%)	797 (100%)	2 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/253 (88%)	224 (100%)	0	100	100
1	B	217/253 (86%)	217 (100%)	0	100	100
1	C	204/253 (81%)	204 (100%)	0	100	100
All	All	645/759 (85%)	645 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	163	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	401	-	4,4,4	0.48	0	6,6,6	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	401	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/320 (86%)	0.53	41 (14%) 3 5	18, 43, 128, 144	0
1	B	271/320 (84%)	0.61	39 (14%) 3 5	23, 53, 114, 133	0
1	C	263/320 (82%)	0.82	48 (18%) 2 2	25, 51, 138, 154	0
All	All	810/960 (84%)	0.65	128 (15%) 3 4	18, 50, 129, 154	0

All (128) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	354	THR	8.4
1	C	263	GLY	7.6
1	C	309	ASN	7.4
1	C	74	LEU	7.3
1	C	297	GLY	7.1
1	C	303	LEU	6.5
1	A	342	LEU	6.4
1	A	264	ARG	6.3
1	A	301	ASN	6.2
1	A	229	PRO	5.8
1	C	230	GLU	5.7
1	B	75	GLU	5.6
1	C	289	PRO	5.6
1	A	297	GLY	5.5
1	A	294	ALA	5.3
1	C	335	VAL	5.3
1	A	265	GLU	5.0
1	B	289	PRO	5.0
1	C	313	ILE	4.9
1	B	342	LEU	4.8
1	A	163	ASN	4.8
1	C	334	VAL	4.8
1	A	164	LEU	4.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	262	GLY	4.7
1	A	289	PRO	4.7
1	A	266	ILE	4.6
1	B	164	LEU	4.5
1	B	165	GLY	4.5
1	C	75	GLU	4.5
1	B	313	ILE	4.4
1	B	135	THR	4.4
1	B	77	ARG	4.4
1	C	330	VAL	4.4
1	B	163	ASN	4.4
1	C	353	ALA	4.4
1	C	100	ASP	4.3
1	C	290	ASP	4.3
1	C	305	ILE	4.3
1	C	316	LEU	4.1
1	C	347	THR	4.1
1	A	313	ILE	4.1
1	C	197	ASN	4.0
1	B	360	HIS	4.0
1	C	328	GLY	4.0
1	C	229	PRO	3.9
1	A	219	SER	3.8
1	B	230	GLU	3.8
1	B	355	ASN	3.7
1	C	325	ILE	3.6
1	B	309	ASN	3.6
1	C	296	ALA	3.6
1	A	311	PRO	3.5
1	C	295	ASN	3.5
1	A	262	GLY	3.5
1	A	300	VAL	3.5
1	A	305	ILE	3.5
1	B	258	TYR	3.5
1	C	164	LEU	3.4
1	C	301	ASN	3.4
1	C	299	GLN	3.4
1	C	283	VAL	3.3
1	B	100	ASP	3.3
1	C	298	ILE	3.3
1	B	286	GLU	3.2
1	C	258	TYR	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	335	VAL	3.2
1	C	300	VAL	3.2
1	C	314	SER	3.2
1	C	333	VAL	3.2
1	B	257	GLY	3.1
1	A	258	TYR	3.1
1	B	316	LEU	3.1
1	B	288	SER	3.1
1	B	290	ASP	3.1
1	B	361	HIS	3.1
1	C	184	THR	3.0
1	C	343	THR	3.0
1	C	198	HIS	3.0
1	A	79	LEU	3.0
1	B	101	ALA	2.9
1	C	163	ASN	2.9
1	C	352	PRO	2.9
1	A	100	ASP	2.9
1	A	160	ASN	2.8
1	B	296	ALA	2.8
1	A	290	ASP	2.7
1	A	310	LYS	2.7
1	B	341	GLN	2.7
1	A	286	GLU	2.7
1	B	354	THR	2.7
1	B	62	TYR	2.6
1	C	287	VAL	2.6
1	A	296	ALA	2.6
1	A	316	LEU	2.6
1	C	292	PRO	2.6
1	B	297	GLY	2.6
1	C	77	ARG	2.6
1	A	186	ARG	2.5
1	A	231	GLY	2.5
1	B	76	ILE	2.5
1	B	305	ILE	2.5
1	A	309	ASN	2.5
1	A	257	GLY	2.4
1	B	65	GLY	2.4
1	A	329	SER	2.4
1	B	160	ASN	2.4
1	B	335	VAL	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	311	PRO	2.4
1	B	109	GLN	2.4
1	B	198	HIS	2.4
1	C	345	GLN	2.4
1	B	102	ASP	2.3
1	B	262	GLY	2.3
1	A	184	THR	2.3
1	C	199	GLY	2.3
1	B	303	LEU	2.3
1	A	263	GLY	2.2
1	A	303	LEU	2.2
1	A	330	VAL	2.2
1	B	352	PRO	2.1
1	C	288	SER	2.1
1	A	299	GLN	2.1
1	A	347	THR	2.1
1	B	263	GLY	2.1
1	A	326	ARG	2.0
1	C	282	ILE	2.0
1	A	323	ALA	2.0
1	A	230	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NI	B	401	1/1	0.79	0.40	3.03	116,116,116,116	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	PO4	A	401	5/5	0.93	0.15	-	32,45,68,73	5

6.5 Other polymers [i](#)

There are no such residues in this entry.