



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 12:42 PM GMT

PDB ID : 3RSJ  
Title : Structure of HCRF in complex with Ganglioside GD1a  
Authors : Fu, Z.; Benson, M.A.; Barbieri, J.T.; Kim, J-J.P.; Baldwin, M.R.  
Deposited on : 2011-05-02  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	413	<div> <div>7%</div> <div>67%29%..</div> </div>
1	B	413	<div> <div>5%</div> <div>69%25%..</div> </div>
1	C	413	<div> <div>5%</div> <div>69%26%..</div> </div>
1	D	413	<div> <div>6%</div> <div>74%21%..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SIA	A	5	-	-	-	X
2	SIA	D	5	-	-	-	X
4	SIA	C	5	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 13922 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BoNT/F.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	406	Total	C	N	O	S	0	0	0
			3334	2114	577	635	8			
1	B	402	Total	C	N	O	S	0	0	0
			3307	2097	572	630	8			
1	C	402	Total	C	N	O	S	0	0	0
			3307	2097	572	630	8			
1	D	404	Total	C	N	O	S	0	0	0
			3324	2109	575	632	8			

- Molecule 2 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	3	Total	C	N	O	0	0
			46	25	2	19		
2	D	3	Total	C	N	O	0	0
			46	25	2	19		

- Molecule 3 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	5	Total	C	N	O	0	0
			77	42	3	32		

- Molecule 4 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	4	Total	C	N	O	0	0
			57	31	2	24		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	117	Total 117	O 117	0	0
5	B	98	Total 98	O 98	0	0
5	C	96	Total 96	O 96	0	0
5	D	113	Total 113	O 113	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: BoNT/F





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.47Å 84.27Å 117.60Å 72.55° 87.01° 67.13°	Depositor
Resolution (Å)	29.86 – 2.00 29.85 – 2.00	Depositor EDS
% Data completeness (in resolution range)	87.0 (29.86-2.00) 79.2 (29.85-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.86 (at 2.00Å)	Xtriage
Refinement program	CNS 1.3	Depositor
R, $R_{free}$	0.232 , 0.263 0.227 , 0.258	Depositor DCC
$R_{free}$ test set	12918 reflections (11.07%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.9	Xtriage
Anisotropy	0.592	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 44.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 129601 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	13922	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, GAL, NGA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/3401	0.65	0/4603
1	B	0.37	0/3374	0.64	0/4568
1	C	0.37	0/3374	0.64	0/4568
1	D	0.38	0/3391	0.66	0/4590
All	All	0.37	0/13540	0.65	0/18329

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3334	0	3249	104	0
1	B	3307	0	3217	108	0
1	C	3307	0	3217	102	0
1	D	3324	0	3241	84	0
2	A	46	0	39	2	0
2	D	46	0	39	2	0
3	B	77	0	63	5	0
4	C	57	0	48	3	0
5	A	117	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	98	0	0	1	0
5	C	96	0	0	3	0
5	D	113	0	0	0	0
All	All	13922	0	13113	396	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 396 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:953:ASN:HD21	1:D:1055:ASN:H	1.05	1.00
1:B:1229:ASN:HD21	1:B:1233:GLY:H	1.15	0.93
1:A:953:ASN:HD21	1:A:1055:ASN:H	1.17	0.91
1:D:945:TYR:H	1:D:997:GLN:HE22	1.18	0.90
1:C:1229:ASN:HD21	1:C:1233:GLY:H	1.22	0.88

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	400/413 (97%)	360 (90%)	37 (9%)	3 (1%)	24	15
1	B	396/413 (96%)	355 (90%)	37 (9%)	4 (1%)	19	11
1	C	396/413 (96%)	360 (91%)	33 (8%)	3 (1%)	24	15
1	D	398/413 (96%)	362 (91%)	32 (8%)	4 (1%)	19	11
All	All	1590/1652 (96%)	1437 (90%)	139 (9%)	14 (1%)	21	13

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	926	ILE
1	C	1257	LYS
1	D	1084	GLU
1	A	1257	LYS
1	D	926	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	375/382 (98%)	362 (96%)	13 (4%)	43	40
1	B	372/382 (97%)	358 (96%)	14 (4%)	40	36
1	C	372/382 (97%)	360 (97%)	12 (3%)	46	44
1	D	374/382 (98%)	357 (96%)	17 (4%)	34	29
All	All	1493/1528 (98%)	1437 (96%)	56 (4%)	40	36

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1126	ASN
1	C	1037	ASN
1	D	1126	ASN
1	B	1229	ASN
1	C	973	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 106 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	1229	ASN
1	C	982	GLN
1	D	1096	ASN
1	B	1231	ASN
1	C	928	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

15 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NGA	A	3	2	14,14,15	0.96	0	12,19,21	1.17	1 (8%)
2	GAL	A	4	2	11,11,12	1.19	1 (9%)	12,15,17	1.26	1 (8%)
2	SIA	A	5	2	17,21,21	2.70	3 (17%)	19,31,31	1.17	2 (10%)
3	GAL	B	2	3	10,10,12	1.21	1 (10%)	13,13,17	1.39	2 (15%)
3	NGA	B	3	3	14,14,15	0.93	0	12,19,21	1.30	2 (16%)
3	GAL	B	4	3	11,11,12	1.05	2 (18%)	12,15,17	1.29	1 (8%)
3	SIA	B	5	3	17,21,21	2.61	3 (17%)	19,31,31	1.26	1 (5%)
3	SIA	B	6	3	17,21,21	3.35	4 (23%)	19,31,31	1.32	2 (10%)
4	GAL	C	2	4	11,11,12	0.87	0	15,15,17	1.02	0
4	NGA	C	3	4	14,14,15	0.98	1 (7%)	12,19,21	1.41	3 (25%)
4	GAL	C	4	4	11,11,12	1.06	2 (18%)	12,15,17	1.21	1 (8%)
4	SIA	C	5	4	17,21,21	2.66	2 (11%)	19,31,31	1.26	1 (5%)
2	NGA	D	3	2	14,14,15	1.07	2 (14%)	12,19,21	1.27	2 (16%)
2	GAL	D	4	2	11,11,12	1.21	1 (9%)	12,15,17	1.29	1 (8%)
2	SIA	D	5	2	17,21,21	2.81	3 (17%)	19,31,31	1.12	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NGA	A	3	2	-	0/6/22/26	0/1/1/1
2	GAL	A	4	2	-	0/2/18/22	0/1/1/1
2	SIA	A	5	2	-	0/14/38/38	0/1/1/1
3	GAL	B	2	3	-	0/2/15/22	0/1/1/1
3	NGA	B	3	3	-	0/6/22/26	0/1/1/1
3	GAL	B	4	3	-	0/2/18/22	0/1/1/1
3	SIA	B	5	3	-	0/14/38/38	0/1/1/1
3	SIA	B	6	3	-	0/14/38/38	0/1/1/1
4	GAL	C	2	4	-	0/2/18/22	0/1/1/1
4	NGA	C	3	4	-	0/6/22/26	0/1/1/1
4	GAL	C	4	4	-	0/2/18/22	0/1/1/1
4	SIA	C	5	4	-	0/14/38/38	0/1/1/1
2	NGA	D	3	2	-	0/6/22/26	0/1/1/1
2	GAL	D	4	2	-	0/2/18/22	0/1/1/1
2	SIA	D	5	2	-	0/14/38/38	0/1/1/1

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	2	GAL	C1-C2	-2.86	1.50	1.52
4	C	3	NGA	C8-C7	-2.07	1.46	1.50
2	D	3	NGA	C8-C7	-2.06	1.46	1.50
2	D	5	SIA	O6-C6	2.06	1.47	1.44
2	D	3	NGA	C1-C2	2.06	1.55	1.53

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	6	SIA	C2-C3-C4	-3.89	102.31	110.69
3	B	5	SIA	O2-C2-C3	-3.89	104.55	109.41
4	C	5	SIA	O2-C2-C3	-3.76	104.72	109.41
3	B	4	GAL	C6-C5-C4	-3.63	108.66	113.29
2	A	4	GAL	C6-C5-C4	-3.40	108.95	113.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	4	GAL	2	0
3	B	2	GAL	1	0
3	B	3	NGA	1	0
3	B	4	GAL	3	0
3	B	6	SIA	2	0
4	C	2	GAL	1	0
4	C	4	GAL	2	0
2	D	4	GAL	1	0
2	D	5	SIA	1	0

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	406/413 (98%)	0.30	27 (6%) 21 22	23, 39, 58, 72	0
1	B	402/413 (97%)	0.24	19 (4%) 35 37	24, 40, 62, 81	0
1	C	402/413 (97%)	0.29	22 (5%) 29 30	26, 40, 61, 76	0
1	D	404/413 (97%)	0.24	23 (5%) 27 29	24, 38, 58, 77	0
All	All	1614/1652 (97%)	0.27	91 (5%) 28 29	23, 39, 60, 81	0

The worst 5 of 91 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	929	GLY	8.9
1	D	931	TYR	6.9
1	B	1162	THR	6.2
1	B	931	TYR	6.1
1	A	1162	THR	6.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SIA	D	5	21/21	0.76	0.24	4.36	73,76,79,79	0
2	SIA	A	5	21/21	0.79	0.27	3.57	64,66,72,74	0
4	SIA	C	5	21/21	0.87	0.21	2.00	60,62,65,67	0
4	GAL	C	4	11/12	0.81	0.15	1.62	57,60,63,65	0
2	GAL	D	4	11/12	0.86	0.17	1.53	68,71,72,73	0
2	GAL	A	4	11/12	0.81	0.15	1.31	59,63,65,66	0
3	SIA	B	5	21/21	0.87	0.15	0.22	49,56,60,61	0
3	GAL	B	4	11/12	0.92	0.11	-0.03	43,46,49,51	0
3	GAL	B	2	10/12	0.86	0.23	-	72,74,77,81	0
2	NGA	A	3	14/15	0.81	0.25	-	66,69,71,72	0
2	NGA	D	3	14/15	0.73	0.33	-	71,75,76,77	0
4	GAL	C	2	11/12	0.69	0.26	-	80,83,84,86	0
3	NGA	B	3	14/15	0.90	0.15	-	45,55,58,65	0
3	SIA	B	6	21/21	0.72	0.36	-	77,83,85,86	0
4	NGA	C	3	14/15	0.76	0.25	-	64,71,73,78	0

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.