



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:42 PM GMT

PDB ID : 3RU3  
Title : Crystal structure of tm0922, a fusion of a domain of unknown function and ADP/ATP-dependent NAD(P)H-hydrate dehydratase from *Thermotoga maritima* soaked with NADPH and ATP.  
Authors : Shumilin, I.A.; Cymborowski, M.; Lesley, S.A.; Minor, W.  
Deposited on : 2011-05-04  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	502	<div> <div>5%</div> <div>87%</div> <div>10%</div> <div>.</div> </div>
2	B	6	<div> <div>100%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	K	A	491	-	-	-	X
4	MG	A	492	-	-	-	X
6	ATP	A	494	-	-	-	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 3918 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	489	Total	C	N	O	S	0	0	0
			3721	2376	628	706	11			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	EXPRESSION TAG	UNP Q9X024
A	-10	GLY	-	EXPRESSION TAG	UNP Q9X024
A	-9	SER	-	EXPRESSION TAG	UNP Q9X024
A	-8	ASP	-	EXPRESSION TAG	UNP Q9X024
A	-7	LYS	-	EXPRESSION TAG	UNP Q9X024
A	-6	ILE	-	EXPRESSION TAG	UNP Q9X024
A	-5	HIS	-	EXPRESSION TAG	UNP Q9X024
A	-4	HIS	-	EXPRESSION TAG	UNP Q9X024
A	-3	HIS	-	EXPRESSION TAG	UNP Q9X024
A	-2	HIS	-	EXPRESSION TAG	UNP Q9X024
A	-1	HIS	-	EXPRESSION TAG	UNP Q9X024
A	0	HIS	-	EXPRESSION TAG	UNP Q9X024

- Molecule 2 is a protein called Unknown peptide, probably from expression host.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	6	Total	C	N	O	0	0	0
			52	37	7	8			


- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	K	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C<sub>21</sub>H<sub>30</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			40	15	6	16	3		

- Molecule 6 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 7 is BETA-6-HYDROXY-1,4,5,6-TETRAHYDRONICOTINAMIDE ADENINE DINUCLEOTIDE PHOSPHATE (three-letter code: NPW) (formula:  $C_{21}H_{32}N_7O_{18}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	P	0	0
			49	21	7	18	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	26	Total	O	0	0
			26	26		
8	B	1	Total	O	0	0
			1	1		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative uncharacterized protein



- Molecule 2: Unknown peptide, probably from expression host



There are no outlier residues recorded for this chain.

## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.94Å 120.94Å 154.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.60 42.76 – 2.60	Depositor EDS
% Data completeness (in resolution range)	93.8 (50.00-2.60) 93.8 (42.76-2.60)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.20 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.162 , 0.237 0.169 , 0.242	Depositor DCC
$R_{free}$ test set	869 reflections (5.44%)	DCC
Wilson B-factor (Å <sup>2</sup> )	53.1	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 45.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 16834 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3918	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NPW, NDP, MG, K, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.92	0/3779	0.87	2/5112 (0.0%)
2	B	1.36	0/54	1.07	0/73
All	All	0.93	0/3833	0.87	2/5185 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	243	LEU	CB-CG-CD1	-5.71	101.29	111.00
1	A	220	LEU	CB-CG-CD1	-5.11	102.31	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3721	0	3839	30	0
2	B	52	0	45	0	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	40	0	19	1	0
6	A	27	0	12	2	0
7	A	49	0	28	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	A	26	0	0	0	0
8	B	1	0	0	0	0
All	All	3918	0	3943	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:206:ILE:CG2	1:A:208:ARG:O	2.39	0.70
1:A:152:ILE:HD11	1:A:187:LEU:HD12	1.77	0.67
1:A:402:LYS:HB3	6:A:494:ATP:H3'	1.84	0.58
1:A:75:LEU:HD22	1:A:132:ILE:HD13	1.86	0.57
1:A:234:LYS:HG3	1:A:262:LEU:HB3	1.88	0.54
1:A:364:THR:O	1:A:402:LYS:HE2	2.09	0.53
1:A:24:GLY:HA3	1:A:58:VAL:CG1	2.40	0.52
1:A:106:LEU:HD13	1:A:139:SER:HB3	1.92	0.51
1:A:4:ILE:CD1	1:A:202:LEU:HD22	2.41	0.51
1:A:206:ILE:HG22	1:A:208:ARG:O	2.12	0.49
7:A:495:NPW:O3	7:A:495:NPW:O3A	2.18	0.49
1:A:347:ASN:OD1	1:A:372:ARG:NH1	2.46	0.49
1:A:311:VAL:HG11	1:A:438:ALA:O	2.14	0.48
1:A:75:LEU:CD2	1:A:132:ILE:HD13	2.44	0.48
1:A:235:VAL:HG23	1:A:260:THR:HG21	1.95	0.48
1:A:428:GLY:N	6:A:494:ATP:O3B	2.46	0.47
1:A:1:MET:CE	1:A:174:VAL:HG22	2.44	0.47
1:A:77:LYS:N	1:A:77:LYS:HD2	2.31	0.45
1:A:75:LEU:HD22	1:A:132:ILE:CD1	2.47	0.45
1:A:4:ILE:HD12	1:A:202:LEU:HB3	1.99	0.45
1:A:41:TYR:HB2	1:A:43:PHE:CE1	2.52	0.44
1:A:132:ILE:O	1:A:136:VAL:HG23	2.18	0.43
1:A:370:MET:O	1:A:374:VAL:HG22	2.18	0.43
1:A:466:GLU:CD	1:A:466:GLU:H	2.22	0.43
1:A:214:GLU:CD	1:A:214:GLU:H	2.21	0.43
1:A:129:TYR:CE2	5:A:493:NDP:H2A	2.53	0.42
1:A:125:ILE:HD13	1:A:133:ILE:CD1	2.49	0.42
1:A:249:VAL:HG22	1:A:275:ALA:HA	2.01	0.41
1:A:24:GLY:HA3	1:A:58:VAL:HG12	2.02	0.41
1:A:44:LEU:HD11	1:A:73:VAL:HG23	2.03	0.41
1:A:52:ASN:O	1:A:55:ASP:HB2	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	487/502 (97%)	473 (97%)	14 (3%)	0	100	100
2	B	4/6 (67%)	4 (100%)	0	0	100	100
All	All	491/508 (97%)	477 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	410/422 (97%)	398 (97%)	12 (3%)	50	77
2	B	4/4 (100%)	4 (100%)	0	100	100
All	All	414/426 (97%)	402 (97%)	12 (3%)	50	77

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	LYS
1	A	77	LYS
1	A	158	LYS
1	A	209	TYR
1	A	234	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	278	ARG
1	A	352	SER
1	A	394	GLU
1	A	396	ASP
1	A	403	SER
1	A	415	LEU
1	A	426	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	323	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	NDP	A	493	-	35,43,52	1.15	3 (8%)	44,67,80	2.21	6 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	ATP	A	494	4	22,29,33	1.33	2 (9%)	27,45,52	2.42	10 (37%)
7	NPW	A	495	-	42,53,53	1.35	5 (11%)	49,82,82	1.80	11 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NDP	A	493	-	-	0/23/59/77	0/4/4/5
6	ATP	A	494	4	-	0/12/32/38	0/3/3/3
7	NPW	A	495	-	-	0/30/80/80	0/5/5/5

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	495	NPW	C2C-C3C	-2.53	1.28	1.34
5	A	493	NDP	PN-O2N	-2.28	1.45	1.54
7	A	495	NPW	O5B-C5B	-2.09	1.36	1.44
7	A	495	NPW	O4D-C1A	2.15	1.43	1.41
5	A	493	NDP	C2A-N1A	2.63	1.38	1.33
6	A	494	ATP	C5-C4	2.94	1.47	1.40
7	A	495	NPW	C5A-C4D	3.02	1.47	1.40
5	A	493	NDP	C2A-N3A	3.88	1.39	1.32
6	A	494	ATP	O4'-C1'	4.10	1.46	1.41
7	A	495	NPW	C8A-N7A	5.82	1.45	1.34

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	493	NDP	N3A-C2A-N1A	-12.00	119.71	128.89
6	A	494	ATP	N3-C2-N1	-8.06	122.72	128.89
7	A	495	NPW	N3D-C2D-N1D	-7.35	123.27	128.89
6	A	494	ATP	C1'-N9-C4	-3.71	121.35	126.94
5	A	493	NDP	PN-O3-PA	-3.47	122.99	132.73
6	A	494	ATP	C4-C5-N7	-2.99	106.72	109.48
7	A	495	NPW	O2A-P-O2	-2.88	99.91	107.11
5	A	493	NDP	O5D-PN-O1N	-2.81	98.72	109.62
7	A	495	NPW	O7C-C7C-N7C	-2.39	116.82	122.76
6	A	494	ATP	C2'-C1'-N9	-2.37	110.67	114.29
7	A	495	NPW	O5B-PB-OB1	-2.35	100.51	109.62
5	A	493	NDP	C1B-N9A-C4A	-2.24	123.56	126.94

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	494	ATP	PA-O3A-PB	-2.11	125.58	132.67
7	A	495	NPW	P-O2A-C2A	-2.08	116.57	121.56
7	A	495	NPW	O4B-C1B-C2B	-2.05	101.82	106.58
7	A	495	NPW	OA1-PA-OA3	2.02	114.27	105.09
5	A	493	NDP	O2X-P2B-O1X	2.10	117.33	110.58
6	A	494	ATP	C2'-C3'-C4'	2.14	107.01	102.61
6	A	494	ATP	O5'-C5'-C4'	2.21	117.25	109.12
6	A	494	ATP	O2B-PB-O1B	2.56	118.81	110.58
7	A	495	NPW	C4D-C5A-N7A	2.67	111.93	109.48
7	A	495	NPW	N6D-C6D-N1D	2.68	124.96	119.20
7	A	495	NPW	O3-P-O2	2.71	119.31	110.58
7	A	495	NPW	O5E-C5E-C4E	2.82	119.53	109.12
5	A	493	NDP	O2N-PN-O1N	2.87	128.08	112.53
6	A	494	ATP	O2A-PA-O1A	3.00	128.78	112.53
6	A	494	ATP	O4'-C1'-N9	3.54	115.51	108.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	493	NDP	1	0
6	A	494	ATP	2	0
7	A	495	NPW	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	489/502 (97%)	0.23	24 (4%) 33 26	34, 48, 80, 101	0
2	B	6/6 (100%)	-0.46	0 100 100	48, 49, 64, 65	0
All	All	495/508 (97%)	0.22	24 (4%) 34 27	34, 49, 80, 101	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	101	PHE	4.2
1	A	278	ARG	3.8
1	A	276	THR	3.7
1	A	279	PHE	3.7
1	A	105	ILE	3.6
1	A	277	SER	3.5
1	A	358	LYS	3.4
1	A	288	ASP	3.2
1	A	290	GLU	3.2
1	A	280	PRO	3.0
1	A	275	ALA	3.0
1	A	274	ILE	2.8
1	A	100	GLN	2.7
1	A	249	VAL	2.6
1	A	253	MET	2.6
1	A	282	LEU	2.4
1	A	356	GLU	2.3
1	A	252	GLY	2.2
1	A	96	LYS	2.2
1	A	1	MET	2.2
1	A	107	ASN	2.1
1	A	108	GLU	2.0
1	A	102	GLU	2.0
1	A	246	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	K	A	491	1/1	0.98	0.23	3.92	70,70,70,70	0
6	ATP	A	494	27/31	0.90	0.27	3.74	67,85,91,95	0
4	MG	A	492	1/1	0.91	0.23	2.69	67,67,67,67	0
7	NPW	A	495	49/49	0.95	0.15	-0.70	39,48,73,78	0
5	NDP	A	493	40/48	0.95	0.14	-0.72	39,59,75,78	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.