



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 12:43 PM GMT

PDB ID : 3RZX  
Title : Mouse importin alpha-Ku70 NLS peptide complex  
Authors : Takeda, A.A.S.; Fontes, M.R.M.  
Deposited on : 2011-05-12  
Resolution : 2.61 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2700 (2.64-2.60)
Clashscore	102246	3065 (2.64-2.60)
Ramachandran outliers	100387	3015 (2.64-2.60)
Sidechain outliers	100360	3015 (2.64-2.60)
RSRZ outliers	91569	2706 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	510	
2	B	22	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3373 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Importin subunit alpha-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	427	3195	2037	541	607	10	0	0	0

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	MET	-	EXPRESSION TAG	UNP P52293
A	21	HIS	-	EXPRESSION TAG	UNP P52293
A	22	HIS	-	EXPRESSION TAG	UNP P52293
A	23	HIS	-	EXPRESSION TAG	UNP P52293
A	24	HIS	-	EXPRESSION TAG	UNP P52293
A	25	HIS	-	EXPRESSION TAG	UNP P52293
A	26	HIS	-	EXPRESSION TAG	UNP P52293
A	27	SER	-	EXPRESSION TAG	UNP P52293
A	28	SER	-	EXPRESSION TAG	UNP P52293
A	29	GLY	-	EXPRESSION TAG	UNP P52293
A	30	LEU	-	EXPRESSION TAG	UNP P52293
A	31	VAL	-	EXPRESSION TAG	UNP P52293
A	32	PRO	-	EXPRESSION TAG	UNP P52293
A	33	ARG	-	EXPRESSION TAG	UNP P52293
A	34	GLY	-	EXPRESSION TAG	UNP P52293
A	35	SER	-	EXPRESSION TAG	UNP P52293
A	36	GLY	-	EXPRESSION TAG	UNP P52293
A	37	MET	-	EXPRESSION TAG	UNP P52293
A	38	LYS	-	EXPRESSION TAG	UNP P52293
A	39	GLU	-	EXPRESSION TAG	UNP P52293
A	40	THR	-	EXPRESSION TAG	UNP P52293
A	41	ALA	-	EXPRESSION TAG	UNP P52293
A	42	ALA	-	EXPRESSION TAG	UNP P52293
A	43	ALA	-	EXPRESSION TAG	UNP P52293
A	44	LYS	-	EXPRESSION TAG	UNP P52293
A	45	PHE	-	EXPRESSION TAG	UNP P52293
A	46	GLU	-	EXPRESSION TAG	UNP P52293

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	ARG	-	EXPRESSION TAG	UNP P52293
A	48	GLN	-	EXPRESSION TAG	UNP P52293
A	49	HIS	-	EXPRESSION TAG	UNP P52293
A	50	MET	-	EXPRESSION TAG	UNP P52293
A	51	ASP	-	EXPRESSION TAG	UNP P52293
A	52	SER	-	EXPRESSION TAG	UNP P52293
A	53	PRO	-	EXPRESSION TAG	UNP P52293
A	54	ASP	-	EXPRESSION TAG	UNP P52293
A	55	LEU	-	EXPRESSION TAG	UNP P52293
A	56	GLY	-	EXPRESSION TAG	UNP P52293
A	57	THR	-	EXPRESSION TAG	UNP P52293
A	58	ASP	-	EXPRESSION TAG	UNP P52293
A	59	ASP	-	EXPRESSION TAG	UNP P52293
A	60	ASP	-	EXPRESSION TAG	UNP P52293
A	61	ASP	-	EXPRESSION TAG	UNP P52293
A	62	LYS	-	EXPRESSION TAG	UNP P52293
A	63	ALA	-	EXPRESSION TAG	UNP P52293
A	64	MET	-	EXPRESSION TAG	UNP P52293
A	65	ALA	-	EXPRESSION TAG	UNP P52293
A	66	ASP	-	EXPRESSION TAG	UNP P52293
A	67	ILE	-	EXPRESSION TAG	UNP P52293
A	68	GLY	-	EXPRESSION TAG	UNP P52293
A	69	SER	-	EXPRESSION TAG	UNP P52293

- Molecule 2 is a protein called Ku70 NLS peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	12	Total	C	N	O	0	0	0
			82	48	17	17			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	89	Total	O	0	0
			89	89		
3	B	7	Total	O	0	0
			7	7		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.52Å 90.00Å 100.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.07 – 2.61 29.07 – 2.61	Depositor EDS
% Data completeness (in resolution range)	99.2 (29.07-2.61) 99.2 (29.07-2.61)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.58 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.168 , 0.223 0.168 , 0.222	Depositor DCC
$R_{free}$ test set	1126 reflections (5.38%)	DCC
Wilson B-factor (Å <sup>2</sup> )	53.8	Xtriage
Anisotropy	0.070	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 50.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 22103 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3373	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.12	5/3253 (0.2%)	1.10	10/4445 (0.2%)
2	B	1.42	0/82	1.35	0/105
All	All	1.13	5/3335 (0.1%)	1.11	10/4550 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	223	CYS	CB-SG	-8.35	1.68	1.82
1	A	79	GLU	CG-CD	6.28	1.61	1.51
1	A	125	VAL	CB-CG1	5.96	1.65	1.52
1	A	237	CYS	CB-SG	-5.12	1.73	1.81
1	A	340	PHE	CB-CG	-5.11	1.42	1.51

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	456	GLU	N-CA-CB	-9.97	92.65	110.60
1	A	481	ASN	CB-CA-C	6.89	124.18	110.40
1	A	441	LEU	CA-CB-CG	-6.67	99.95	115.30
1	A	258	ARG	NE-CZ-NH1	-6.28	117.16	120.30
1	A	101	ARG	NE-CZ-NH2	-6.17	117.21	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	108	LYS	Peptide
1	A	455	GLY	Mainchain
1	A	496	PHE	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3195	0	3219	106	0
2	B	82	0	79	10	0
3	A	89	0	0	10	0
3	B	7	0	0	3	0
All	All	3373	0	3298	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 111 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:308:PRO:HA	2:B:547:ASN:CB	1.59	1.29
1:A:308:PRO:CA	2:B:547:ASN:CB	2.11	1.26
2:B:549:GLY:HA3	3:B:71:HOH:O	1.34	1.23
1:A:472:LYS:O	1:A:476:LEU:HD23	1.38	1.17
1:A:308:PRO:N	2:B:547:ASN:CB	2.10	1.14

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was



analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	425/510 (83%)	412 (97%)	13 (3%)	0	100	100
2	B	10/22 (46%)	8 (80%)	1 (10%)	1 (10%)	1	0
All	All	435/532 (82%)	420 (97%)	14 (3%)	1 (0%)	52	76

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	549	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/426 (80%)	313 (91%)	30 (9%)	13	24
2	B	8/19 (42%)	6 (75%)	2 (25%)	1	1
All	All	351/445 (79%)	319 (91%)	32 (9%)	12	22

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	350	ASN
1	A	392	LYS
1	A	476	LEU
1	A	390	ASP
1	A	395	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	352	GLN
1	A	375	ASN
1	A	477	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	346	ASN
1	A	449	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	427/510 (83%)	-0.33	12 (2%) 56 50	33, 48, 91, 100	0
2	B	12/22 (54%)	0.23	2 (16%) 2 1	36, 49, 70, 78	0
All	All	439/532 (82%)	-0.31	14 (3%) 51 44	33, 48, 91, 100	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	485	TYR	5.4
2	B	547	ASN	5.3
1	A	497	SER	4.4
1	A	74	VAL	3.4
1	A	482	GLU	3.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.