



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 12, 2016 – 07:23 PM EDT

PDB ID : 4S0U
Title : Crystal structure of NKG2D in complex with ULBP6
Authors : Mohammed, F.; Willcox, B.E.
Deposited on : 2015-01-06
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027107
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027107


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1352 (2.38-2.34)
Clashscore	102246	1456 (2.38-2.34)
Ramachandran outliers	100387	1435 (2.38-2.34)
Sidechain outliers	100360	1436 (2.38-2.34)
RSRZ outliers	91569	1358 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	126	<div> <div>4%</div> <div>71%</div> <div>25%</div> <div>..</div> </div>
1	B	126	<div> <div>18%</div> <div>77%</div> <div>23%</div> </div>
2	C	175	<div> <div>11%</div> <div>79%</div> <div>21%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3516 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NKG2-D type II integral membrane protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	124	Total	C	N	O	S	7	0	0
			1004	637	161	194	12			
1	B	126	Total	C	N	O	S	14	0	0
			1019	648	163	196	12			

- Molecule 2 is a protein called Retinoic acid early transcript 1L protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	175	Total	C	N	O	S	28	0	0
			1405	893	236	264	12			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	34	SER	CYS	ENGINEERED MUTATION	UNP Q5VY80


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	44	Total	O	0	0
			44	44		
3	B	15	Total	O	0	0
			15	15		
3	C	29	Total	O	0	0
			29	29		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NKG2-D type II integral membrane protein



- Molecule 1: NKG2-D type II integral membrane protein



- Molecule 2: Retinoic acid early transcript 1L protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	82.30Å 80.00Å 73.20Å 90.00° 106.50° 90.00°	Depositor
Resolution (Å)	18.73 – 2.35 18.30 – 2.35	Depositor EDS
% Data completeness (in resolution range)	94.6 (18.73-2.35) 94.8 (18.30-2.35)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.33 (at 2.35Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.238 , 0.286 0.238 , 0.287	Depositor DCC
R_{free} test set	876 reflections (5.10%)	DCC
Wilson B-factor (Å ²)	37.8	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 30.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 18065 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3516	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/1032	0.63	0/1398
1	B	0.53	1/1048 (0.1%)	0.56	0/1420
2	C	0.53	1/1442 (0.1%)	0.56	0/1953
All	All	0.55	2/3522 (0.1%)	0.58	0/4771

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	154	TRP	CD2-CE2	5.33	1.47	1.41
2	C	158	TRP	CD2-CE2	5.08	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1004	0	940	22	0
1	B	1019	0	959	16	0
2	C	1405	0	1364	25	0
3	A	44	0	0	2	0
3	B	15	0	0	0	0
3	C	29	0	0	1	0
All	All	3516	0	3263	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:PRO:HB2	1:B:93:GLU:HG2	1.78	0.65
1:A:145:LEU:HD23	1:B:104:ILE:HD11	1.78	0.64
1:A:115:ASP:O	1:A:117:SER:N	2.34	0.61
1:A:149:VAL:O	1:A:194:SER:HB2	2.01	0.60
2:C:118:PRO:O	2:C:120:THR:HG23	2.05	0.56
2:C:33:LEU:HB2	2:C:194:LEU:HD13	1.86	0.56
1:B:106:TYR:CD2	1:B:145:LEU:HD22	2.41	0.56
1:B:163:ASN:O	1:B:165:SER:N	2.39	0.55
1:A:116:GLU:O	1:A:118:LYS:HG3	2.08	0.54
1:B:113:PHE:CD2	1:B:210:ILE:HD12	2.43	0.54
1:B:106:TYR:HD2	1:B:145:LEU:HD22	1.73	0.53
1:A:163:ASN:HD22	1:A:165:SER:H	1.56	0.53
1:A:167:GLN:NE2	1:A:171:GLY:HA2	2.24	0.52
2:C:108:ILE:HD11	2:C:168:MET:HA	1.90	0.52
2:C:198:LEU:HA	2:C:201:MET:HE3	1.91	0.52
1:A:124:GLN:HG3	1:A:169:GLU:OE2	2.10	0.52
2:C:132:GLU:HG2	2:C:133:GLY:N	2.24	0.52
2:C:132:GLU:HG2	2:C:133:GLY:H	1.76	0.50
2:C:38:THR:O	2:C:50:CYS:HA	2.11	0.50
1:A:111:TYR:OH	1:A:143:GLN:HG2	2.11	0.50
1:A:118:LYS:HB3	1:A:122:GLU:HB2	1.94	0.49
1:A:99:CYS:O	1:B:93:GLU:HG3	2.12	0.49
1:A:162:THR:HG22	3:A:329:HOH:O	2.13	0.48
2:C:127:CYS:HB2	2:C:191:ILE:HG13	1.95	0.48
1:A:195:SER:HB3	3:A:328:HOH:O	2.13	0.48
1:B:104:ILE:O	1:B:110:CYS:HA	2.15	0.47
1:B:137:VAL:HG23	1:B:155:MET:HE1	1.96	0.47
1:B:153:HIS:O	1:B:192:TYR:N	2.46	0.46
2:C:36:ASP:O	2:C:52:VAL:HA	2.14	0.46
1:A:180:THR:HG21	3:C:326:HOH:O	2.14	0.46
2:C:139:TRP:HB2	2:C:151:PHE:HB3	1.96	0.46
1:A:154:TRP:HB2	1:A:209:TYR:HB3	1.97	0.45
1:B:157:LEU:HB2	1:B:168:TRP:CZ3	2.51	0.45
2:C:40:ILE:HG23	2:C:120:THR:HG22	1.98	0.45
2:C:132:GLU:HG2	2:C:134:HIS:H	1.82	0.44
2:C:42:LYS:HD3	2:C:118:PRO:HB3	2.00	0.44
2:C:172:TRP:N	2:C:172:TRP:CD1	2.86	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:180:MET:HG2	2:C:184:TYR:CE1	2.53	0.44
1:B:187:GLY:HA3	1:B:202:ASN:HA	1.99	0.44
2:C:140:GLN:HG2	2:C:150:LEU:HD22	1.99	0.44
1:A:159:HIS:O	1:A:188:ASP:HB3	2.18	0.43
1:A:184:MET:HE3	2:C:187:MET:HB2	2.00	0.43
2:C:105:LEU:HD11	2:C:121:LEU:CB	2.49	0.43
1:B:157:LEU:HB3	1:B:190:ALA:HB3	2.00	0.42
2:C:180:MET:HG2	2:C:184:TYR:HE1	1.84	0.42
1:B:113:PHE:CD2	1:B:210:ILE:CD1	3.02	0.42
1:B:184:MET:HG3	2:C:106:LEU:HD11	2.02	0.42
2:C:85:MET:HB3	2:C:196:ASP:OD2	2.20	0.42
2:C:52:VAL:HB	2:C:64:TYR:HB3	2.02	0.41
2:C:187:MET:HA	2:C:191:ILE:HD12	2.02	0.41
1:A:157:LEU:HB2	1:A:168:TRP:CZ3	2.56	0.41
1:A:119:ASN:HB2	1:A:204:SER:HA	2.03	0.41
1:A:113:PHE:CD2	1:A:210:ILE:HG12	2.56	0.41
1:B:160:ILE:HA	1:B:161:PRO:HD2	1.94	0.41
1:A:214:ARG:O	1:A:215:THR:C	2.59	0.40
2:C:125:MET:HG3	2:C:139:TRP:CZ2	2.57	0.40
2:C:151:PHE:CE2	2:C:153:SER:HA	2.56	0.40
1:A:163:ASN:ND2	1:A:165:SER:H	2.19	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	122/126 (97%)	117 (96%)	4 (3%)	1 (1%)	24	26
1	B	124/126 (98%)	112 (90%)	9 (7%)	3 (2%)	7	5
2	C	173/175 (99%)	162 (94%)	11 (6%)	0	100	100
All	All	419/427 (98%)	391 (93%)	24 (6%)	4 (1%)	19	20

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	116	GLU
1	B	164	GLY
1	B	132	ALA
1	B	176	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/116 (98%)	112 (98%)	2 (2%)	66	81
1	B	116/116 (100%)	112 (97%)	4 (3%)	44	57
2	C	157/157 (100%)	157 (100%)	0	100	100
All	All	387/389 (100%)	381 (98%)	6 (2%)	70	83

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	145	LEU
1	A	163	ASN
1	B	115	ASP
1	B	151	SER
1	B	162	THR
1	B	205	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	163	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	124/126 (98%)	0.31	5 (4%) 42 56	18, 27, 42, 73	2 (1%)
1	B	126/126 (100%)	1.02	23 (18%) 2 3	30, 48, 64, 70	5 (3%)
2	C	175/175 (100%)	0.66	19 (10%) 7 13	23, 39, 69, 83	8 (4%)
All	All	425/427 (99%)	0.66	47 (11%) 7 12	18, 38, 65, 83	15 (3%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	131	ASN	4.7
2	C	133	GLY	4.4
2	C	30	PRO	4.4
2	C	29	ASP	4.1
2	C	203	SER	3.6
2	C	174	ASN	3.6
1	A	92	THR	3.6
2	C	107	ASP	3.5
2	C	131	ALA	3.5
2	C	166	ARG	3.4
1	A	215	THR	3.4
1	B	125	ALA	3.4
1	B	128	MET	3.3
2	C	132	GLU	3.2
2	C	114	THR	3.2
1	B	163	ASN	3.0
1	B	215	THR	3.0
2	C	130	LYS	2.9
1	B	205	THR	2.8
2	C	111	GLU	2.7
2	C	113	TYR	2.7
2	C	110	LEU	2.6
1	B	108	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	123	SER	2.5
1	B	214	ARG	2.5
1	B	134	LEU	2.5
1	B	154	TRP	2.5
2	C	202	ASP	2.4
2	C	112	ASN	2.4
1	B	206	PRO	2.4
1	B	210	ILE	2.3
1	A	214	ARG	2.3
1	B	116	GLU	2.3
1	B	124	GLN	2.3
1	B	122	GLU	2.2
1	B	155	MET	2.2
1	B	176	PRO	2.2
1	B	91	LEU	2.2
1	A	117	SER	2.1
1	A	134	LEU	2.1
2	C	67	GLY	2.1
2	C	76	PRO	2.1
1	B	173	ILE	2.1
1	B	152	TYR	2.1
2	C	43	PHE	2.1
1	B	121	TYR	2.0
1	B	164	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.