



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:00 PM GMT

PDB ID : 3SGB
Title : STRUCTURE OF THE COMPLEX OF STREPTOMYCES GRISEUS PROTEASE B AND THE THIRD DOMAIN OF THE TURKEY OVOMUCOID INHIBITOR AT 1.8 ANGSTROMS RESOLUTION
Authors : Read, R.J.; Fujinaga, M.; Sielecki, A.R.; James, M.N.G.
Deposited on : 1983-01-21
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	E	185	 82% 16% ..
2	I	56	 70% 16% .. 11%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1872 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEINASE B (SGPB).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	185	Total	C	N	O	S	0	0	0
			1310	801	228	275	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	235A	VAL	SER	CONFLICT	UNP P00777


- Molecule 2 is a protein called TURKEY OVOMUCOID INHIBITOR (OMTKY3).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	50	Total	C	N	O	S	0	0	0
			380	233	64	77	6			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	135	Total	O	0	0
			135	135		
3	I	47	Total	O	0	0
			47	47		

Note EDS was not executed.

- Chain E: 

- Chain I:

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	45.35Å 54.52Å 45.65Å 90.00° 119.20° 90.00°	Depositor
Resolution (Å)	6.00 – 1.80	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-1.80)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	unknown	Depositor
R, R_{free}	0.125 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1872	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	1.15	1/1335 (0.1%)	1.76	23/1820 (1.3%)
2	I	1.00	0/388	1.69	6/523 (1.1%)
All	All	1.12	1/1723 (0.1%)	1.75	29/2343 (1.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	58	CYS	CB-SG	5.02	1.90	1.82

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	138	ARG	NE-CZ-NH1	12.26	126.43	120.30
1	E	48(A)	ARG	NE-CZ-NH2	-9.12	115.74	120.30
1	E	48(A)	ARG	NE-CZ-NH1	8.83	124.72	120.30
2	I	47	THR	N-CA-CB	-8.16	94.79	110.30
1	E	182	ARG	CD-NE-CZ	8.00	134.80	123.60
1	E	41	ARG	NE-CZ-NH1	7.83	124.22	120.30
1	E	233	GLU	CA-CB-CG	7.58	130.08	113.40
1	E	116	ASP	CB-CG-OD2	-6.51	112.44	118.30
2	I	47	THR	OG1-CB-CG2	6.40	124.72	110.00
2	I	15	ALA	N-CA-CB	6.34	118.98	110.10
2	I	19	GLU	CG-CD-OE1	6.26	130.82	118.30
1	E	107	ARG	NE-CZ-NH2	-6.24	117.18	120.30
2	I	31	TYR	CG-CD1-CE1	-6.09	116.42	121.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	162	VAL	O-C-N	6.02	132.33	122.70
1	E	41	ARG	O-C-N	5.95	132.22	122.70
1	E	90	GLY	O-C-N	5.84	132.05	122.70
1	E	138	ARG	CD-NE-CZ	5.62	131.47	123.60
1	E	81	ARG	NE-CZ-NH1	5.53	123.07	120.30
1	E	51	TYR	CA-CB-CG	5.51	123.88	113.40
1	E	105	ILE	O-C-N	5.50	131.50	122.70
1	E	52	PHE	CB-CG-CD2	-5.44	116.99	120.80
1	E	129	ASN	O-C-N	5.31	131.20	122.70
1	E	120	GLY	N-CA-C	-5.30	99.86	113.10
1	E	51	TYR	O-C-N	5.22	131.06	122.70
1	E	171	TYR	CB-CG-CD2	5.22	124.14	121.00
1	E	52	PHE	N-CA-CB	5.13	119.84	110.60
2	I	37	PHE	CB-CG-CD1	-5.11	117.22	120.80
1	E	160	GLY	O-C-N	5.09	130.85	122.70
1	E	171	TYR	CB-CG-CD1	-5.08	117.95	121.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	81	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1310	0	1231	16	0
2	I	380	0	349	10	0
3	E	135	0	0	1	0
3	I	47	0	0	1	0
All	All	1872	0	1580	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:29:LYS:HE2	3:I:83:HOH:O	1.59	1.01
2:I:21:ARG:HB3	2:I:21:ARG:HH21	1.37	0.86
1:E:110:ASN:C	1:E:110:ASN:HD22	1.84	0.81
1:E:166:ASN:HD22	1:E:179:GLY:HA2	1.45	0.81
2:I:44:SER:HB2	2:I:47:THR:HG22	1.70	0.74
2:I:11:TYR:CZ	2:I:13:LYS:HE3	2.28	0.68
2:I:21:ARG:HH21	2:I:21:ARG:CB	2.07	0.67
2:I:11:TYR:HB3	2:I:12:PRO:HA	1.77	0.64
1:E:166:ASN:ND2	1:E:179:GLY:HA2	2.18	0.58
1:E:110:ASN:ND2	1:E:113:ILE:H	2.06	0.54
1:E:107:ARG:NH1	3:E:374:HOH:O	2.23	0.52
2:I:21:ARG:HH21	2:I:21:ARG:CG	2.23	0.51
2:I:11:TYR:CE1	2:I:13:LYS:HE3	2.47	0.50
1:E:110:ASN:HD22	1:E:111:THR:N	2.09	0.49
2:I:44:SER:O	2:I:47:THR:HB	2.12	0.49
1:E:33:SER:HB3	1:E:66:TRP:CH2	2.48	0.48
1:E:110:ASN:HD21	1:E:113:ILE:H	1.63	0.46
1:E:108:TYR:CD2	1:E:115:LYS:HE3	2.50	0.46
1:E:67:TRP:CE3	1:E:81:ARG:HG3	2.52	0.45
1:E:199:LEU:HD13	1:E:228:PHE:CZ	2.53	0.43
1:E:110:ASN:ND2	1:E:110:ASN:C	2.59	0.43
1:E:166:ASN:HA	1:E:179:GLY:HA2	2.00	0.42
1:E:33:SER:HB3	1:E:66:TRP:CZ3	2.55	0.42
2:I:21:ARG:NH2	2:I:21:ARG:CG	2.82	0.41
1:E:113:ILE:HA	1:E:114:PRO:HD3	1.88	0.41
1:E:181:ILE:O	1:E:227:PHE:HA	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	183/185 (99%)	173 (94%)	10 (6%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	I	48/56 (86%)	46 (96%)	2 (4%)	0	100	100
All	All	231/241 (96%)	219 (95%)	12 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	138/138 (100%)	135 (98%)	3 (2%)	60	45
2	I	44/48 (92%)	42 (96%)	2 (4%)	34	16
All	All	182/186 (98%)	177 (97%)	5 (3%)	52	36

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	34	SER
1	E	48(A)	ARG
1	E	110	ASN
2	I	21	ARG
2	I	47	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	110	ASN
1	E	166	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.