



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:13 PM GMT

PDB ID : 3SJL  
Title : Crystal Structure of the P107S-MauG/pre-Methylamine Dehydrogenase Complex  
Authors : Jensen, L.M.R.; Wilmot, C.M.  
Deposited on : 2011-06-21  
Resolution : 1.63 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1953 (1.66-1.62)
Clashscore	102246	2091 (1.66-1.62)
Ramachandran outliers	100387	2052 (1.66-1.62)
Sidechain outliers	100360	2052 (1.66-1.62)
RSRZ outliers	91569	1955 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	373	<div> <div>2%</div> <div>78% 12% • 9%</div> </div>
1	B	373	<div> <div>2%</div> <div>80% 11% • 8%</div> </div>
2	C	137	<div> <div>2%</div> <div>81% 14% • •</div> </div>
2	E	137	<div> <div>%</div> <div>78% 12% • 9%</div> </div>
3	D	386	<div> <div>2%</div> <div>88% 9% • •</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	386	<div> <div></div> <div>%</div> <div>84%</div> <div>13%</div> <div></div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NA	A	402	-	-	-	X
7	PEG	B	374	-	-	X	X
8	EDO	D	387	-	X	X	X
8	EDO	F	388	-	-	-	X
9	MES	F	387	-	-	-	X

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 15480 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methylamine utilization protein mauG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	338	Total	C	N	O	S	0	6	0
			2659	1669	475	503	12			
1	B	343	Total	C	N	O	S	0	9	0
			2726	1705	495	514	12			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	107	SER	PRO	ENGINEERED MUTATION	UNP Q51658
A	368	HIS	-	EXPRESSION TAG	UNP Q51658
A	369	HIS	-	EXPRESSION TAG	UNP Q51658
A	370	HIS	-	EXPRESSION TAG	UNP Q51658
A	371	HIS	-	EXPRESSION TAG	UNP Q51658
A	372	HIS	-	EXPRESSION TAG	UNP Q51658
A	373	HIS	-	EXPRESSION TAG	UNP Q51658
B	107	SER	PRO	ENGINEERED MUTATION	UNP Q51658
B	368	HIS	-	EXPRESSION TAG	UNP Q51658
B	369	HIS	-	EXPRESSION TAG	UNP Q51658
B	370	HIS	-	EXPRESSION TAG	UNP Q51658
B	371	HIS	-	EXPRESSION TAG	UNP Q51658
B	372	HIS	-	EXPRESSION TAG	UNP Q51658
B	373	HIS	-	EXPRESSION TAG	UNP Q51658

- Molecule 2 is a protein called Methylamine dehydrogenase light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	131	Total	C	N	O	S	0	3	0
			1027	634	179	200	14			
2	E	125	Total	C	N	O	S	0	4	0
			978	610	162	192	14			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	132	HIS	-	EXPRESSION TAG	UNP P22619
C	133	HIS	-	EXPRESSION TAG	UNP P22619
C	134	HIS	-	EXPRESSION TAG	UNP P22619
C	135	HIS	-	EXPRESSION TAG	UNP P22619
C	136	HIS	-	EXPRESSION TAG	UNP P22619
C	137	HIS	-	EXPRESSION TAG	UNP P22619
E	132	HIS	-	EXPRESSION TAG	UNP P22619
E	133	HIS	-	EXPRESSION TAG	UNP P22619
E	134	HIS	-	EXPRESSION TAG	UNP P22619
E	135	HIS	-	EXPRESSION TAG	UNP P22619
E	136	HIS	-	EXPRESSION TAG	UNP P22619
E	137	HIS	-	EXPRESSION TAG	UNP P22619

- Molecule 3 is a protein called Methylamine dehydrogenase heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	376	Total	C	N	O	S	0	3	0
			2937	1865	502	562	8			
3	F	376	Total	C	N	O	S	0	7	0
			2964	1884	507	565	8			


- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		
4	A	1	Total	Ca	0	0
			1	1		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total	Na	0	0
			2	2		
5	A	2	Total	Na	0	0
			2	2		

- Molecule 6 is HEME C (three-letter code: HEC) (formula: C<sub>34</sub>H<sub>34</sub>FeN<sub>4</sub>O<sub>4</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
6	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
6	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
6	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	4	3		
7	B	1	Total	C	O	0	0
			7	4	3		
7	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 8 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			4	2	2		
8	D	1	Total	C	O	0	0
			4	2	2		
8	F	1	Total	C	O	0	0
			4	2	2		

- Molecule 9 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula:  $C_6H_{13}NO_4S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	F	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	400	Total	O	0	0
			400	400		
10	B	383	Total	O	0	0
			383	383		
10	C	127	Total	O	0	0
			127	127		
10	D	385	Total	O	0	0
			385	385		
10	E	137	Total	O	0	0
			137	137		
10	F	534	Total	O	0	0
			534	534		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Methylamine utilization protein mauG




- Molecule 1: Methylamine utilization protein mauG





- Molecule 2: Methylamine dehydrogenase light chain



- Molecule 2: Methylamine dehydrogenase light chain



Chain D:  2% 88% 9% ..



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.61Å 89.00Å 104.81Å 67.05° 79.51° 79.72°	Depositor
Resolution (Å)	29.39 – 1.63 29.39 – 1.63	Depositor EDS
% Data completeness (in resolution range)	92.4 (29.39-1.63) 82.8 (29.39-1.63)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.05 (at 1.63Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.142 , 0.180 0.161 , 0.189	Depositor DCC
$R_{free}$ test set	10468 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.0	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 36.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 208545 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	15480	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 0AF, NA, CA, EDO, MES, HEC, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.39	13/2738 (0.5%)	1.30	23/3712 (0.6%)
1	B	1.31	7/2812 (0.2%)	1.19	15/3808 (0.4%)
2	C	1.37	1/1050 (0.1%)	1.17	6/1430 (0.4%)
2	E	1.33	2/1001 (0.2%)	1.20	3/1368 (0.2%)
3	D	1.24	3/3023 (0.1%)	1.13	9/4120 (0.2%)
3	F	1.44	13/3062 (0.4%)	1.25	20/4170 (0.5%)
All	All	1.35	39/13686 (0.3%)	1.21	76/18608 (0.4%)

The worst 5 of 39 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	213	GLU	CD-OE1	9.55	1.36	1.25
1	A	112	VAL	CB-CG2	-7.53	1.37	1.52
1	B	162	TYR	CD2-CE2	6.98	1.49	1.39
1	A	93	TRP	CE3-CZ3	6.93	1.50	1.38
3	F	213	GLU	CG-CD	6.71	1.62	1.51

The worst 5 of 76 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	39[A]	ARG	NE-CZ-NH1	-14.86	112.87	120.30
1	A	39[B]	ARG	NE-CZ-NH1	-14.86	112.87	120.30
1	A	177	ARG	NE-CZ-NH1	9.70	125.15	120.30
3	D	104	ARG	NE-CZ-NH2	-9.50	115.55	120.30
3	F	278	ARG	NE-CZ-NH2	-9.33	115.63	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2659	0	2566	15	0
1	B	2726	0	2635	24	0
2	C	1027	0	916	15	0
2	E	978	0	890	13	0
3	D	2937	0	2832	12	0
3	F	2964	0	2874	19	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	86	0	60	3	0
6	B	86	0	60	1	0
7	A	7	0	10	0	0
7	B	14	0	20	5	0
8	B	4	0	6	1	0
8	D	4	0	6	6	0
8	F	4	0	6	1	0
9	F	12	0	12	0	0
10	A	400	0	0	3	0
10	B	383	0	0	8	0
10	C	127	0	0	5	0
10	D	385	0	0	5	0
10	E	137	0	0	5	0
10	F	534	0	0	5	0
All	All	15480	0	12893	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 97 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:153:ASP:OD1	7:B:374:PEG:H11	1.50	1.09
3:F:372[B]:GLN:NE2	10:F:1487:HOH:O	1.87	1.08
3:F:372[A]:GLN:OE1	10:F:2086:HOH:O	1.73	1.04

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:64:ASP:HB2	10:B:423:HOH:O	1.58	1.03
8:F:388:EDO:H12	10:F:1692:HOH:O	1.63	0.96

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/373 (91%)	331 (97%)	9 (3%)	0	100	100
1	B	348/373 (93%)	336 (97%)	12 (3%)	0	100	100
2	C	131/137 (96%)	126 (96%)	5 (4%)	0	100	100
2	E	126/137 (92%)	124 (98%)	2 (2%)	0	100	100
3	D	377/386 (98%)	366 (97%)	10 (3%)	1 (0%)	46	23
3	F	381/386 (99%)	370 (97%)	10 (3%)	1 (0%)	46	23
All	All	1703/1792 (95%)	1653 (97%)	48 (3%)	2 (0%)	56	31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	D	102	ILE
3	F	102	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/292 (93%)	266 (98%)	5 (2%)	66	41
1	B	277/292 (95%)	269 (97%)	8 (3%)	50	18
2	C	113/112 (101%)	111 (98%)	2 (2%)	66	41
2	E	108/112 (96%)	105 (97%)	3 (3%)	51	20
3	D	307/311 (99%)	297 (97%)	10 (3%)	45	14
3	F	311/311 (100%)	306 (98%)	5 (2%)	70	46
All	All	1387/1430 (97%)	1354 (98%)	33 (2%)	57	26

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	29	CYS
3	D	218	GLU
3	F	293	ARG
3	D	94	HIS
3	D	208	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	210	GLN
2	C	134	HIS
3	F	14	GLN
1	B	106	GLN
3	D	30	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	0AF	C	57	2	13,16,17	1.49	3 (23%)	10,22,24	1.78	2 (20%)
2	0AF	E	57	2	13,16,17	2.11	4 (30%)	10,22,24	1.64	3 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	0AF	C	57	2	-	0/3/6/8	0/2/2/2
2	0AF	E	57	2	-	0/3/6/8	0/2/2/2

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	57	0AF	CZ2-CE2	-3.69	1.37	1.42
2	C	57	0AF	CB-CA	-3.43	1.46	1.53
2	C	57	0AF	CZ2-CE2	-2.52	1.39	1.42
2	C	57	0AF	CZ3-CE3	2.31	1.42	1.36
2	E	57	0AF	CB-CG	2.69	1.59	1.51

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	57	0AF	O-C-CA	-3.68	115.90	125.49
2	E	57	0AF	O-C-CA	-3.15	117.28	125.49
2	C	57	0AF	CB-CG-CD1	-2.77	124.54	127.97
2	E	57	0AF	CE3-CZ3-CH2	-2.38	117.01	120.96
2	E	57	0AF	CE3-CD2-CE2	2.48	122.14	117.03

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	57	0AF	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	57	0AF	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 6 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	PEG	A	374	-	6,6,6	0.83	0	5,5,5	1.25	1 (20%)
6	HEC	A	500	1	24,50,50	1.61	6 (25%)	19,82,82	3.23	7 (36%)
6	HEC	A	600	1	24,50,50	1.35	3 (12%)	19,82,82	3.56	10 (52%)
7	PEG	B	374	-	6,6,6	1.26	1 (16%)	5,5,5	1.27	0
7	PEG	B	375	-	6,6,6	0.77	0	5,5,5	0.97	0
8	EDO	B	376	-	3,3,3	0.52	0	2,2,2	0.78	0
6	HEC	B	500	1	24,50,50	1.75	6 (25%)	19,82,82	2.95	6 (31%)
6	HEC	B	600	1	24,50,50	1.18	3 (12%)	19,82,82	3.22	8 (42%)
8	EDO	D	387	-	3,3,3	2.07	2 (66%)	2,2,2	3.13	2 (100%)
9	MES	F	387	-	11,12,12	0.84	0	14,16,16	2.91	4 (28%)
8	EDO	F	388	-	3,3,3	0.97	0	2,2,2	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	PEG	A	374	-	-	0/4/4/4	0/0/0/0
6	HEC	A	500	1	-	0/6/54/54	0/0/8/8

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	HEC	A	600	1	-	0/6/54/54	0/0/8/8
7	PEG	B	374	-	-	0/4/4/4	0/0/0/0
7	PEG	B	375	-	-	0/4/4/4	0/0/0/0
8	EDO	B	376	-	-	0/1/1/1	0/0/0/0
6	HEC	B	500	1	-	0/6/54/54	0/0/8/8
6	HEC	B	600	1	-	0/6/54/54	0/0/8/8
8	EDO	D	387	-	-	0/1/1/1	0/0/0/0
9	MES	F	387	-	-	0/6/14/14	0/1/1/1
8	EDO	F	388	-	-	0/1/1/1	0/0/0/0

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	500	HEC	C3C-C2C	-3.91	1.36	1.40
6	B	500	HEC	C3B-C2B	-3.61	1.37	1.40
6	A	500	HEC	C4B-NB	-2.82	1.32	1.36
6	B	500	HEC	C4B-NB	-2.70	1.33	1.36
6	B	600	HEC	C3B-C2B	-2.68	1.38	1.40

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	600	HEC	CBB-CAB-C3B	-10.58	103.84	127.35
6	A	500	HEC	CBB-CAB-C3B	-8.90	107.58	127.35
6	B	600	HEC	CBB-CAB-C3B	-8.49	108.49	127.35
6	B	500	HEC	CBB-CAB-C3B	-8.06	109.44	127.35
6	A	500	HEC	CBC-CAC-C3C	-7.10	111.57	127.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	500	HEC	2	0
6	A	600	HEC	1	0
7	B	374	PEG	4	0
7	B	375	PEG	1	0
8	B	376	EDO	1	0
6	B	500	HEC	1	0
8	D	387	EDO	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	F	388	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	338/373 (90%)	-0.14	9 (2%) 58 56	15, 21, 39, 61	0
1	B	343/373 (91%)	-0.21	6 (1%) 73 73	13, 21, 38, 61	0
2	C	130/137 (94%)	-0.18	3 (2%) 64 63	14, 19, 32, 49	0
2	E	124/137 (90%)	-0.31	1 (0%) 87 88	13, 17, 27, 55	0
3	D	376/386 (97%)	-0.11	9 (2%) 62 61	15, 26, 43, 66	0
3	F	376/386 (97%)	-0.31	2 (0%) 91 92	12, 17, 28, 53	0
All	All	1687/1792 (94%)	-0.20	30 (1%) 71 72	12, 20, 39, 66	0

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5	PRO	10.9
1	B	49	ALA	6.1
1	A	359	GLU	5.5
3	D	208	THR	5.5
1	A	6	ALA	5.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	0AF	E	57	15/16	0.96	0.08	-	16,18,22,24	0
2	0AF	C	57	15/16	0.97	0.08	-	17,20,22,25	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
8	EDO	F	388	4/4	0.86	0.17	13.45	33,33,34,40	0
9	MES	F	387	12/12	0.97	0.12	4.71	23,39,44,45	0
7	PEG	B	374	7/7	0.79	0.21	3.70	24,37,45,47	0
5	NA	A	402	1/1	0.98	0.09	2.08	26,26,26,26	0
8	EDO	D	387	4/4	0.90	0.12	2.02	25,28,31,32	0
7	PEG	A	374	7/7	0.89	0.14	1.52	26,31,37,37	0
8	EDO	B	376	4/4	0.88	0.09	0.91	36,42,47,48	0
6	HEC	B	600	43/43	0.99	0.11	0.34	9,13,16,18	0
6	HEC	A	500	43/43	0.99	0.10	0.21	13,15,18,19	0
6	HEC	A	600	43/43	0.99	0.10	0.11	13,16,18,19	0
6	HEC	B	500	43/43	0.98	0.09	-0.12	12,16,19,22	0
4	CA	B	400	1/1	1.00	0.07	-1.17	16,16,16,16	0
5	NA	B	402	1/1	0.99	0.04	-1.41	24,24,24,24	0
4	CA	A	400	1/1	1.00	0.04	-2.35	19,19,19,19	0
5	NA	B	401	1/1	0.96	0.23	-	30,30,30,30	0
7	PEG	B	375	7/7	0.85	0.10	-	43,48,52,53	0
5	NA	A	401	1/1	0.95	0.14	-	31,31,31,31	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.