



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:03 PM GMT

PDB ID : 3SJO
Title : structure of EV71 3C in complex with Rupintrivir (AG7088)
Authors : Lu, G.; Qi, J.; Chen, Z.; Xu, X.; Gao, F.; Lin, D.; Qian, W.; Liu, H.; Jiang, H.; Yan, J.; Gao, G.F.
Deposited on : 2011-06-21
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	190	<div> <div>3%</div> <div>77%</div> <div>16%</div> <div>6%</div> </div>
1	B	190	<div> <div>17%</div> <div>59%</div> <div>33%</div> <div>6%</div> </div>
1	C	190	<div> <div>2%</div> <div>76%</div> <div>15%</div> <div>6%</div> </div>
1	D	190	<div> <div>3%</div> <div>81%</div> <div>11%</div> <div>6%</div> </div>
1	E	190	<div> <div>5%</div> <div>73%</div> <div>21%</div> <div>6%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	190	
1	G	190	
1	H	190	

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 12391 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3C protease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	179	Total	C	N	O	S	0	0	0
			1391	881	247	256	7			
1	B	179	Total	C	N	O	S	0	0	0
			1391	881	247	256	7			
1	C	179	Total	C	N	O	S	0	0	0
			1391	881	247	256	7			
1	D	179	Total	C	N	O	S	0	0	0
			1391	881	247	256	7			
1	E	179	Total	C	N	O	S	0	0	0
			1391	881	247	256	7			
1	F	179	Total	C	N	O	S	0	0	0
			1391	881	247	256	7			
1	G	179	Total	C	N	O	S	0	0	0
			1391	881	247	256	7			
1	H	179	Total	C	N	O	S	0	0	0
			1391	881	247	256	7			

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	INITIATING METHIONINE	UNP E0WWC7
A	184	HIS	-	EXPRESSION TAG	UNP E0WWC7
A	185	HIS	-	EXPRESSION TAG	UNP E0WWC7
A	186	HIS	-	EXPRESSION TAG	UNP E0WWC7
A	187	HIS	-	EXPRESSION TAG	UNP E0WWC7
A	188	HIS	-	EXPRESSION TAG	UNP E0WWC7
A	189	HIS	-	EXPRESSION TAG	UNP E0WWC7
B	0	MET	-	INITIATING METHIONINE	UNP E0WWC7
B	184	HIS	-	EXPRESSION TAG	UNP E0WWC7
B	185	HIS	-	EXPRESSION TAG	UNP E0WWC7
B	186	HIS	-	EXPRESSION TAG	UNP E0WWC7
B	187	HIS	-	EXPRESSION TAG	UNP E0WWC7
B	188	HIS	-	EXPRESSION TAG	UNP E0WWC7

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	189	HIS	-	EXPRESSION TAG	UNP E0WWC7
C	0	MET	-	INITIATING METHIONINE	UNP E0WWC7
C	184	HIS	-	EXPRESSION TAG	UNP E0WWC7
C	185	HIS	-	EXPRESSION TAG	UNP E0WWC7
C	186	HIS	-	EXPRESSION TAG	UNP E0WWC7
C	187	HIS	-	EXPRESSION TAG	UNP E0WWC7
C	188	HIS	-	EXPRESSION TAG	UNP E0WWC7
C	189	HIS	-	EXPRESSION TAG	UNP E0WWC7
D	0	MET	-	INITIATING METHIONINE	UNP E0WWC7
D	184	HIS	-	EXPRESSION TAG	UNP E0WWC7
D	185	HIS	-	EXPRESSION TAG	UNP E0WWC7
D	186	HIS	-	EXPRESSION TAG	UNP E0WWC7
D	187	HIS	-	EXPRESSION TAG	UNP E0WWC7
D	188	HIS	-	EXPRESSION TAG	UNP E0WWC7
D	189	HIS	-	EXPRESSION TAG	UNP E0WWC7
E	0	MET	-	INITIATING METHIONINE	UNP E0WWC7
E	184	HIS	-	EXPRESSION TAG	UNP E0WWC7
E	185	HIS	-	EXPRESSION TAG	UNP E0WWC7
E	186	HIS	-	EXPRESSION TAG	UNP E0WWC7
E	187	HIS	-	EXPRESSION TAG	UNP E0WWC7
E	188	HIS	-	EXPRESSION TAG	UNP E0WWC7
E	189	HIS	-	EXPRESSION TAG	UNP E0WWC7
F	0	MET	-	INITIATING METHIONINE	UNP E0WWC7
F	184	HIS	-	EXPRESSION TAG	UNP E0WWC7
F	185	HIS	-	EXPRESSION TAG	UNP E0WWC7
F	186	HIS	-	EXPRESSION TAG	UNP E0WWC7
F	187	HIS	-	EXPRESSION TAG	UNP E0WWC7
F	188	HIS	-	EXPRESSION TAG	UNP E0WWC7
F	189	HIS	-	EXPRESSION TAG	UNP E0WWC7
G	0	MET	-	INITIATING METHIONINE	UNP E0WWC7
G	184	HIS	-	EXPRESSION TAG	UNP E0WWC7
G	185	HIS	-	EXPRESSION TAG	UNP E0WWC7
G	186	HIS	-	EXPRESSION TAG	UNP E0WWC7
G	187	HIS	-	EXPRESSION TAG	UNP E0WWC7
G	188	HIS	-	EXPRESSION TAG	UNP E0WWC7
G	189	HIS	-	EXPRESSION TAG	UNP E0WWC7
H	0	MET	-	INITIATING METHIONINE	UNP E0WWC7
H	184	HIS	-	EXPRESSION TAG	UNP E0WWC7
H	185	HIS	-	EXPRESSION TAG	UNP E0WWC7
H	186	HIS	-	EXPRESSION TAG	UNP E0WWC7
H	187	HIS	-	EXPRESSION TAG	UNP E0WWC7
H	188	HIS	-	EXPRESSION TAG	UNP E0WWC7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	189	HIS	-	EXPRESSION TAG	UNP E0WWC7

- Molecule 2 is 4-{2-(4-FLUORO-BENZYL)-6-METHYL-5-[(5-METHYL-ISOXAZOLE-3-CARBONYL)-AMINO]-4-OXO-HEPTANOYLAMINO}-5-(2-OXO-PYRROLIDIN-3-YL)-PENTANOIC ACID ETHYL ESTER (three-letter code: AG7) (formula: C₃₁H₄₁FN₄O₇).







Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			43	31	1	4	7		
2	B	1	Total	C	F	N	O	0	0
			43	31	1	4	7		
2	C	1	Total	C	F	N	O	0	0
			43	31	1	4	7		
2	D	1	Total	C	F	N	O	0	0
			43	31	1	4	7		
2	E	1	Total	C	F	N	O	0	0
			43	31	1	4	7		
2	F	1	Total	C	F	N	O	0	0
			43	31	1	4	7		
2	G	1	Total	C	F	N	O	0	0
			43	31	1	4	7		
2	H	1	Total	C	F	N	O	0	0
			43	31	1	4	7		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	124	Total 124	O 124	0	0
3	B	97	Total 97	O 97	0	0
3	C	144	Total 144	O 144	0	0
3	D	137	Total 137	O 137	0	0
3	E	146	Total 146	O 146	0	0
3	F	106	Total 106	O 106	0	0
3	G	73	Total 73	O 73	0	0
3	H	92	Total 92	O 92	0	0

- Molecule 1: 3C protease







• Molecule 1: 3C protease




• Molecule 1: 3C protease



• Molecule 1: 3C protease



• Molecule 1: 3C protease



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	41.26Å 82.17Å 97.95Å 89.96° 89.99° 90.13°	Depositor
Resolution (Å)	34.54 – 1.70 34.54 – 1.70	Depositor EDS
% Data completeness (in resolution range)	93.2 (34.54-1.70) 96.5 (34.54-1.70)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.99 (at 1.70Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.5_2)	Depositor
R, R_{free}	0.205 , 0.222 0.203 , 0.219	Depositor DCC
R_{free} test set	6839 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	20.2	Xtriage
Anisotropy	0.129	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 41.0	EDS
Estimated twinning fraction	0.021 for h,-k,-l 0.019 for -h,k,-l 0.478 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 136404 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12391	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 34.48 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 6.7811e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AG7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/1417	0.58	0/1918
1	B	0.50	3/1417 (0.2%)	0.58	0/1918
1	C	0.48	0/1417	0.58	0/1918
1	D	0.35	0/1417	0.55	0/1918
1	E	0.31	0/1417	0.53	0/1918
1	F	0.34	0/1417	0.54	0/1918
1	G	0.36	0/1417	0.51	0/1918
1	H	0.32	0/1417	0.52	0/1918
All	All	0.40	3/11336 (0.0%)	0.55	0/15344

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	92	GLU	CB-CG	-6.18	1.40	1.52
1	B	100	ALA	N-CA	-5.12	1.36	1.46
1	B	100	ALA	CA-CB	-5.06	1.41	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1391	0	1404	33	0
1	B	1391	0	1404	90	0
1	C	1391	0	1404	37	0
1	D	1391	0	1404	32	0
1	E	1391	0	1404	54	0
1	F	1391	0	1404	81	1
1	G	1391	0	1404	70	0
1	H	1391	0	1404	68	0
2	A	43	0	40	0	0
2	B	43	0	40	7	0
2	C	43	0	40	2	0
2	D	43	0	40	0	0
2	E	43	0	40	0	0
2	F	43	0	40	3	0
2	G	43	0	40	2	0
2	H	43	0	40	1	0
3	A	124	0	0	21	0
3	B	97	0	0	79	0
3	C	144	0	0	25	0
3	D	137	0	0	22	1
3	E	146	0	0	37	0
3	F	106	0	0	66	1
3	G	73	0	0	49	0
3	H	92	0	0	48	2
All	All	12391	0	11552	473	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 473 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:145:GLY:HA2	3:B:481:HOH:O	1.20	1.34
1:B:103:VAL:HG23	3:B:736:HOH:O	1.10	1.28
1:H:101:THR:HA	3:H:502:HOH:O	1.13	1.26
1:G:11:LEU:HD21	3:G:440:HOH:O	1.14	1.25
1:H:14:ASN:HB3	3:H:831:HOH:O	1.14	1.25

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:182:GLU:OE1	3:H:273:HOH:O[1_556]	1.95	0.25
3:D:550:HOH:O	3:D:634:HOH:O[1_455]	2.08	0.12
3:F:884:HOH:O	3:H:776:HOH:O[1_556]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/190 (93%)	170 (96%)	6 (3%)	1 (1%)	30	12
1	B	177/190 (93%)	163 (92%)	13 (7%)	1 (1%)	30	12
1	C	177/190 (93%)	171 (97%)	5 (3%)	1 (1%)	30	12
1	D	177/190 (93%)	172 (97%)	4 (2%)	1 (1%)	30	12
1	E	177/190 (93%)	171 (97%)	5 (3%)	1 (1%)	30	12
1	F	177/190 (93%)	166 (94%)	10 (6%)	1 (1%)	30	12
1	G	177/190 (93%)	171 (97%)	5 (3%)	1 (1%)	30	12
1	H	177/190 (93%)	170 (96%)	6 (3%)	1 (1%)	30	12
All	All	1416/1520 (93%)	1354 (96%)	54 (4%)	8 (1%)	30	12

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	154	VAL
1	D	154	VAL
1	E	154	VAL
1	F	107	GLU
1	G	154	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	154/164 (94%)	150 (97%)	4 (3%)	54	32
1	B	154/164 (94%)	147 (96%)	7 (4%)	34	13
1	C	154/164 (94%)	148 (96%)	6 (4%)	39	16
1	D	154/164 (94%)	147 (96%)	7 (4%)	34	13
1	E	154/164 (94%)	149 (97%)	5 (3%)	46	24
1	F	154/164 (94%)	149 (97%)	5 (3%)	46	24
1	G	154/164 (94%)	147 (96%)	7 (4%)	34	13
1	H	154/164 (94%)	149 (97%)	5 (3%)	46	24
All	All	1232/1312 (94%)	1186 (96%)	46 (4%)	41	18

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	138	TYR
1	E	66	GLN
1	H	39	ARG
1	D	139	ASN
1	D	175	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 47 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	40	HIS
1	D	126	ASN
1	H	55	ASN
1	D	66	GLN
1	E	22	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	AG7	A	501	1	42,45,45	3.58	12 (28%)	45,61,61	1.89	11 (24%)
2	AG7	B	501	1	42,45,45	3.63	12 (28%)	45,61,61	1.92	10 (22%)
2	AG7	C	501	1	42,45,45	3.61	11 (26%)	45,61,61	1.93	13 (28%)
2	AG7	D	501	1	42,45,45	3.59	12 (28%)	45,61,61	1.87	11 (24%)
2	AG7	E	501	1	42,45,45	3.57	11 (26%)	45,61,61	1.98	14 (31%)
2	AG7	F	501	1	42,45,45	3.63	12 (28%)	45,61,61	1.92	10 (22%)
2	AG7	G	501	1	42,45,45	3.66	12 (28%)	45,61,61	1.78	9 (20%)
2	AG7	H	501	1	42,45,45	3.62	12 (28%)	45,61,61	1.84	11 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AG7	A	501	1	-	0/40/54/54	0/2/3/3
2	AG7	B	501	1	-	0/40/54/54	0/2/3/3
2	AG7	C	501	1	-	0/40/54/54	0/2/3/3
2	AG7	D	501	1	-	0/40/54/54	0/2/3/3
2	AG7	E	501	1	-	0/40/54/54	0/2/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AG7	F	501	1	-	0/40/54/54	0/2/3/3
2	AG7	G	501	1	-	0/40/54/54	0/2/3/3
2	AG7	H	501	1	-	0/40/54/54	0/2/3/3

The worst 5 of 94 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	501	AG7	C83-N17	2.12	1.50	1.46
2	F	501	AG7	C1-N5	2.15	1.37	1.33
2	B	501	AG7	C83-N17	2.20	1.51	1.46
2	A	501	AG7	C83-N17	2.20	1.51	1.46
2	F	501	AG7	C83-N17	2.20	1.51	1.46

The worst 5 of 89 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	AG7	O18-C16-N17	-6.18	119.16	125.96
2	E	501	AG7	O18-C16-N17	-5.68	119.70	125.96
2	D	501	AG7	O18-C16-N17	-5.65	119.74	125.96
2	F	501	AG7	O18-C16-N17	-5.54	119.86	125.96
2	B	501	AG7	O18-C16-N17	-5.51	119.90	125.96

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	AG7	7	0
2	C	501	AG7	2	0
2	F	501	AG7	3	0
2	G	501	AG7	2	0
2	H	501	AG7	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	179/190 (94%)	-0.09	5 (2%) 56 61	11, 19, 33, 252	0
1	B	179/190 (94%)	0.87	33 (18%) 2 2	14, 27, 105, 176	0
1	C	179/190 (94%)	0.10	3 (1%) 73 77	10, 21, 41, 204	0
1	D	179/190 (94%)	-0.05	6 (3%) 49 53	10, 19, 35, 132	0
1	E	179/190 (94%)	0.16	9 (5%) 32 35	11, 20, 48, 205	0
1	F	179/190 (94%)	1.01	26 (14%) 3 4	14, 25, 128, 215	0
1	G	179/190 (94%)	0.77	26 (14%) 3 4	14, 27, 108, 176	0
1	H	179/190 (94%)	0.61	16 (8%) 12 13	15, 26, 85, 290	0
All	All	1432/1520 (94%)	0.42	124 (8%) 13 14	10, 23, 87, 290	0

The worst 5 of 124 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	145	GLY	15.1
1	E	4	LEU	15.0
1	H	182	GLU	10.9
1	H	4	LEU	10.7
1	F	4	LEU	10.4

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	AG7	A	501	43/43	0.92	0.10	0.36	14,21,29,36	0
2	AG7	H	501	43/43	0.90	0.13	0.22	20,27,34,37	0
2	AG7	G	501	43/43	0.89	0.12	-0.00	19,27,35,38	0
2	AG7	E	501	43/43	0.91	0.12	-0.00	17,24,32,39	0
2	AG7	B	501	43/43	0.87	0.16	-0.01	23,30,39,41	0
2	AG7	C	501	43/43	0.93	0.10	-0.11	16,24,29,33	0
2	AG7	F	501	43/43	0.85	0.14	-0.12	21,29,37,37	0
2	AG7	D	501	43/43	0.93	0.10	-0.15	12,17,24,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.